



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 9, 2020 – 02:03 PM BST

PDB ID : 6CU0  
Title : Crystal structure of 4-1BBL/4-1BB (C121S) complex in P21 space group  
Authors : Aruna, B.; Zajonc, D.M.; Doukov, T.  
Deposited on : 2018-03-23  
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	165	<div> <div style="width: 100%; height: 10px; background-color: red;"></div> <div style="width: 100%; height: 10px; background-color: green;"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>81%</span> <span>10%</span> <span>8%</span> </div> </div>
1	B	165	<div> <div style="width: 2%; height: 10px; background-color: red;"></div> <div style="width: 100%; height: 10px; background-color: green;"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>81%</span> <span>8%</span> <span>11%</span> </div> </div>
1	C	165	<div> <div style="width: 2%; height: 10px; background-color: red;"></div> <div style="width: 100%; height: 10px; background-color: green;"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>81%</span> <span>7%</span> <span>11%</span> </div> </div>
1	D	165	<div> <div style="width: 100%; height: 10px; background-color: green;"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>79%</span> <span>10%</span> <span>11%</span> </div> </div>
1	E	165	<div> <div style="width: 4%; height: 10px; background-color: red;"></div> <div style="width: 100%; height: 10px; background-color: green;"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>78%</span> <span>9%</span> <span>13%</span> </div> </div>
1	F	165	<div> <div style="width: 5%; height: 10px; background-color: red;"></div> <div style="width: 100%; height: 10px; background-color: green;"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>78%</span> <span>13%</span> <span>8%</span> </div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	138	<div><div></div><div>7%</div><div>87%</div><div>8%</div><div>5%</div></div>
2	H	138	<div><div></div><div>13%</div><div>90%</div><div>6%</div><div></div></div>
2	I	138	<div><div></div><div>10%</div><div>90%</div><div>5%</div><div>5%</div></div>
2	J	138	<div><div></div><div>9%</div><div>87%</div><div>7%</div><div>7%</div></div>
2	K	138	<div><div></div><div>25%</div><div>91%</div><div></div><div></div></div>
2	L	138	<div><div></div><div>8%</div><div>87%</div><div>7%</div><div>7%</div></div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11476 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tumor necrosis factor ligand superfamily member 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	152	Total	C	N	O		0	0	0
			1072	692	185	195				
1	B	147	Total	C	N	O	S	0	0	0
			1033	666	179	187	1			
1	C	147	Total	C	N	O		0	0	0
			999	644	169	186				
1	D	147	Total	C	N	O	S	0	0	0
			1023	656	177	189	1			
1	E	144	Total	C	N	O	S	0	0	0
			987	631	171	184	1			
1	F	151	Total	C	N	O	S	0	0	0
			1046	672	182	191	1			

- Molecule 2 is a protein called Tumor necrosis factor receptor superfamily member 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	G	131	Total	C	N	O	S	0	0	0
			880	517	158	184	21			
2	H	132	Total	C	N	O	S	0	0	0
			899	527	161	190	21			
2	J	129	Total	C	N	O	S	0	0	0
			865	507	154	183	21			
2	K	132	Total	C	N	O	S	0	0	0
			868	507	158	182	21			
2	L	129	Total	C	N	O	S	0	0	0
			863	506	153	184	20			
2	I	131	Total	C	N	O	S	0	0	0
			869	511	151	186	21			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
G	121	SER	CYS	engineered mutation	UNP Q07011

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
H	121	SER	CYS	engineered mutation	UNP Q07011
J	121	SER	CYS	engineered mutation	UNP Q07011
K	121	SER	CYS	engineered mutation	UNP Q07011
L	121	SER	CYS	engineered mutation	UNP Q07011
I	121	SER	CYS	engineered mutation	UNP Q07011

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	G	1	Total	C	N	O	0	0
			14	8	1	5		
3	H	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	5	Total	O	0	0
			5	5		
4	B	2	Total	O	0	0
			2	2		
4	C	1	Total	O	0	0
			1	1		
4	E	1	Total	O	0	0
			1	1		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	F	1	Total 1	O 1	0	0
4	G	7	Total 7	O 7	0	0
4	H	3	Total 3	O 3	0	0
4	J	5	Total 5	O 5	0	0
4	K	13	Total 13	O 13	0	0
4	L	2	Total 2	O 2	0	0
4	I	4	Total 4	O 4	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tumor necrosis factor ligand superfamily member 9




- Molecule 1: Tumor necrosis factor ligand superfamily member 9



- Molecule 1: Tumor necrosis factor ligand superfamily member 9




- Molecule 1: Tumor necrosis factor ligand superfamily member 9




- Molecule 1: Tumor necrosis factor ligand superfamily member 9









- Molecule 1: Tumor necrosis factor ligand superfamily member 9





- Molecule 2: Tumor necrosis factor receptor superfamily member 9





- Molecule 2: Tumor necrosis factor receptor superfamily member 9



- Molecule 2: Tumor necrosis factor receptor superfamily member 9





- Molecule 2: Tumor necrosis factor receptor superfamily member 9




- Molecule 2: Tumor necrosis factor receptor superfamily member 9



Chain L: 



• Molecule 2: Tumor necrosis factor receptor superfamily member 9

Chain I: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	72.96Å 114.75Å 126.04Å 90.00° 101.24° 90.00°	Depositor
Resolution (Å)	50.00 – 3.20 43.83 – 3.20	Depositor EDS
% Data completeness (in resolution range)	94.6 (50.00-3.20) 94.7 (43.83-3.20)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.29 (at 3.19Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, $R_{free}$	0.247 , 0.296 0.258 , 0.302	Depositor DCC
$R_{free}$ test set	1668 reflections (5.22%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	126.3	Xtriage
Anisotropy	0.329	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.26 , 117.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	11476	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	160.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.12% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.47	0/1095	0.73	0/1501
1	B	0.41	0/1054	0.66	0/1443
1	C	0.41	0/1017	0.64	0/1398
1	D	0.46	0/1043	0.66	0/1428
1	E	0.44	0/1006	0.64	0/1377
1	F	0.40	0/1068	0.64	0/1464
2	G	0.48	0/894	0.64	0/1214
2	H	0.45	0/913	0.63	0/1239
2	I	0.46	0/883	0.58	0/1204
2	J	0.41	0/879	0.62	0/1193
2	K	0.40	0/880	0.59	0/1195
2	L	0.49	0/875	0.65	0/1187
All	All	0.44	0/11607	0.64	0/15843

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1072	0	1029	11	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1033	0	990	7	0
1	C	999	0	943	5	0
1	D	1023	0	956	9	0
1	E	987	0	909	7	0
1	F	1046	0	986	12	0
2	G	880	0	701	5	0
2	H	899	0	717	3	0
2	I	869	0	669	3	0
2	J	865	0	675	3	0
2	K	868	0	677	5	0
2	L	863	0	676	4	0
3	G	14	0	13	0	0
3	H	14	0	13	0	0
4	A	5	0	0	1	0
4	B	2	0	0	0	0
4	C	1	0	0	0	0
4	E	1	0	0	0	0
4	F	1	0	0	0	0
4	G	7	0	0	1	0
4	H	3	0	0	1	0
4	I	4	0	0	0	0
4	J	5	0	0	1	0
4	K	13	0	0	3	0
4	L	2	0	0	0	0
All	All	11476	0	9954	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:J:149:ASN:OD1	2:J:150:GLY:N	1.94	1.00
2:L:146:VAL:HA	2:L:158:CYS:SG	2.01	1.00
1:A:189:SER:O	1:A:192:ALA:HB2	1.87	0.73
1:D:139:GLY:HA3	1:D:241:THR:HG21	1.69	0.73
1:D:151:ARG:HE	1:D:194:ASN:HD22	1.37	0.73
2:H:99:CYS:SG	4:H:601:HOH:O	2.48	0.72
1:E:114:GLY:O	2:H:61:THR:HB	1.90	0.71
2:J:99:CYS:SG	4:J:204:HOH:O	2.51	0.69
1:F:191:GLU:OE1	1:F:191:GLU:N	2.24	0.68

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:K:124:THR:HA	4:K:201:HOH:O	1.96	0.64
1:D:115:LEU:HD22	2:I:64:ILE:HD13	1.78	0.64
1:C:112:ASP:HB3	1:C:115:LEU:HD22	1.79	0.63
1:B:112:ASP:HB3	1:B:115:LEU:HD22	1.79	0.63
1:A:115:LEU:HD22	2:G:64:ILE:HD13	1.81	0.61
2:K:136:TRP:N	4:K:201:HOH:O	2.36	0.58
2:K:136:TRP:CD1	4:K:201:HOH:O	2.53	0.55
1:F:193:ARG:O	1:F:194:ASN:OD1	2.25	0.55
1:F:170:LEU:O	1:F:170:LEU:HD23	2.07	0.55
1:F:170:LEU:HD12	1:F:173:ALA:C	2.28	0.54
1:A:190:SER:HA	4:A:304:HOH:O	2.07	0.54
1:C:149:LEU:HD13	1:C:185:LEU:HD21	1.90	0.54
1:B:149:LEU:HD13	1:B:185:LEU:HD21	1.88	0.53
2:L:146:VAL:CA	2:L:158:CYS:SG	2.89	0.53
1:F:168:GLN:HA	1:F:169:PRO:C	2.29	0.53
1:E:149:LEU:HD13	1:E:185:LEU:HD21	1.90	0.52
1:A:149:LEU:HD13	1:A:185:LEU:HD21	1.91	0.52
1:B:150:ARG:NH2	2:K:100:SER:O	2.43	0.52
1:F:149:LEU:HD13	1:F:185:LEU:HD21	1.90	0.52
1:E:146:GLN:HE22	1:F:202:ARG:HD2	1.74	0.51
1:A:150:ARG:NH2	2:G:100:SER:O	2.47	0.48
1:E:94:GLN:HB2	1:F:203:LEU:HD12	1.95	0.47
2:L:150:GLY:O	2:L:151:THR:HB	2.14	0.47
1:B:94:GLN:HB2	1:D:203:LEU:HD12	1.99	0.45
1:D:230:GLN:HE22	2:I:67:GLN:H	1.65	0.45
1:F:170:LEU:CD1	1:F:173:ALA:C	2.85	0.45
1:A:182:THR:HB	1:D:197:PHE:CE2	2.53	0.44
1:B:112:ASP:HB3	1:B:115:LEU:CD2	2.47	0.44
1:E:123:GLY:HA3	1:E:136:ALA:HB3	2.00	0.43
1:C:112:ASP:HB3	1:C:115:LEU:CD2	2.46	0.43
1:A:123:GLY:HA3	1:A:136:ALA:HB3	2.01	0.43
1:F:155:GLY:HA2	1:F:188:ALA:HB1	2.00	0.43
1:F:99:ASN:HD21	1:F:108:SER:N	2.17	0.43
1:A:115:LEU:HD21	2:G:52:SER:HB3	2.01	0.43
1:C:99:ASN:HD21	1:C:108:SER:N	2.17	0.43
1:B:99:ASN:HD21	1:B:108:SER:N	2.17	0.42
1:A:99:ASN:HD21	1:A:108:SER:N	2.17	0.42
1:C:123:GLY:HA3	1:C:136:ALA:HB3	2.02	0.42
1:F:123:GLY:HA3	1:F:136:ALA:HB3	2.02	0.42
2:L:53:PHE:CZ	2:L:83:ASN:HB2	2.55	0.42
1:D:99:ASN:HD21	1:D:108:SER:N	2.17	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:I:53:PHE:CZ	2:I:83:ASN:HB2	2.54	0.42
1:B:123:GLY:HA3	1:B:136:ALA:HB3	2.01	0.42
2:H:53:PHE:CZ	2:H:83:ASN:HB2	2.55	0.41
2:G:53:PHE:CZ	2:G:83:ASN:HB2	2.55	0.41
2:K:53:PHE:CZ	2:K:83:ASN:HB2	2.55	0.41
1:E:99:ASN:HD21	1:E:108:SER:N	2.17	0.41
2:J:53:PHE:CZ	2:J:83:ASN:HB2	2.56	0.41
2:G:76:LYS:NZ	4:G:601:HOH:O	2.53	0.41
1:A:147:LEU:HD22	1:A:163:LEU:HG	2.03	0.40
1:A:203:LEU:HD12	1:D:94:GLN:HB2	2.04	0.40
1:D:123:GLY:HA3	1:D:136:ALA:HB3	2.01	0.40
1:E:135:VAL:HG21	1:E:206:LEU:HD13	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	150/165 (91%)	142 (95%)	7 (5%)	1 (1%)	22	61
1	B	143/165 (87%)	139 (97%)	4 (3%)	0	100	100
1	C	143/165 (87%)	136 (95%)	7 (5%)	0	100	100
1	D	143/165 (87%)	136 (95%)	6 (4%)	1 (1%)	22	61
1	E	138/165 (84%)	133 (96%)	5 (4%)	0	100	100
1	F	147/165 (89%)	143 (97%)	4 (3%)	0	100	100
2	G	127/138 (92%)	116 (91%)	11 (9%)	0	100	100
2	H	128/138 (93%)	117 (91%)	11 (9%)	0	100	100
2	I	127/138 (92%)	116 (91%)	11 (9%)	0	100	100
2	J	125/138 (91%)	116 (93%)	9 (7%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	K	126/138 (91%)	115 (91%)	11 (9%)	0	100	100
2	L	125/138 (91%)	114 (91%)	11 (9%)	0	100	100
All	All	1622/1818 (89%)	1523 (94%)	97 (6%)	2 (0%)	51	83

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	191	GLU
1	A	192	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	98/125 (78%)	93 (95%)	5 (5%)	24	60
1	B	94/125 (75%)	89 (95%)	5 (5%)	22	58
1	C	89/125 (71%)	82 (92%)	7 (8%)	12	43
1	D	91/125 (73%)	87 (96%)	4 (4%)	28	64
1	E	86/125 (69%)	80 (93%)	6 (7%)	15	48
1	F	94/125 (75%)	86 (92%)	8 (8%)	10	38
2	G	91/120 (76%)	86 (94%)	5 (6%)	21	57
2	H	94/120 (78%)	90 (96%)	4 (4%)	29	64
2	I	89/120 (74%)	86 (97%)	3 (3%)	37	70
2	J	88/120 (73%)	84 (96%)	4 (4%)	27	63
2	K	87/120 (72%)	86 (99%)	1 (1%)	73	88
2	L	87/120 (72%)	84 (97%)	3 (3%)	37	70
All	All	1088/1470 (74%)	1033 (95%)	55 (5%)	24	60

All (55) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	108	SER
1	A	145	PHE
1	A	181	LEU
1	A	210	GLN
1	A	218	THR
1	B	115	LEU
1	B	145	PHE
1	B	162	SER
1	B	181	LEU
1	B	218	THR
1	C	108	SER
1	C	115	LEU
1	C	145	PHE
1	C	162	SER
1	C	181	LEU
1	C	210	GLN
1	C	218	THR
1	D	108	SER
1	D	145	PHE
1	D	181	LEU
1	D	218	THR
1	E	108	SER
1	E	162	SER
1	E	181	LEU
1	E	210	GLN
1	E	218	THR
1	E	221	ARG
1	F	108	SER
1	F	145	PHE
1	F	162	SER
1	F	170	LEU
1	F	181	LEU
1	F	210	GLN
1	F	215	HIS
1	F	218	THR
2	G	26	ASP
2	G	67	GLN
2	G	139	CYS
2	G	153	GLU
2	G	158	CYS
2	H	67	GLN
2	H	121	SER
2	H	153	GLU

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
2	H	158	CYS
2	J	67	GLN
2	J	139	CYS
2	J	153	GLU
2	J	158	CYS
2	K	67	GLN
2	L	67	GLN
2	L	101	MET
2	L	153	GLU
2	I	108	GLN
2	I	153	GLU
2	I	158	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	99	ASN
1	A	146	GLN
1	B	99	ASN
1	C	99	ASN
1	C	146	GLN
1	D	99	ASN
1	D	194	ASN
1	D	230	GLN
1	E	99	ASN
1	E	146	GLN
1	F	99	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	NAG	G	500	2	14,14,15	0.37	0	17,19,21	1.48	4 (23%)
3	NAG	H	501	2	14,14,15	0.76	1 (7%)	17,19,21	1.26	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	G	500	2	-	1/6/23/26	0/1/1/1
3	NAG	H	501	2	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	H	501	NAG	C1-C2	2.43	1.56	1.52

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	500	NAG	C1-O5-C5	3.29	116.65	112.19
3	H	501	NAG	C1-C2-N2	2.80	115.27	110.49
3	H	501	NAG	C1-O5-C5	2.73	115.89	112.19
3	G	500	NAG	C4-C3-C2	-2.45	107.42	111.02
3	G	500	NAG	O5-C1-C2	-2.34	107.59	111.29
3	G	500	NAG	O5-C5-C6	2.16	110.59	107.20
3	H	501	NAG	C2-N2-C7	2.14	125.95	122.90

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	G	500	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	152/165 (92%)	0.30	1 (0%) 87 81	88, 121, 179, 235	0
1	B	147/165 (89%)	0.35	4 (2%) 54 39	98, 144, 191, 224	0
1	C	147/165 (89%)	0.29	4 (2%) 54 39	109, 139, 179, 189	0
1	D	147/165 (89%)	0.26	2 (1%) 75 63	100, 132, 175, 211	0
1	E	144/165 (87%)	0.34	6 (4%) 36 23	107, 143, 180, 205	0
1	F	151/165 (91%)	0.43	8 (5%) 26 14	112, 156, 206, 244	0
2	G	131/138 (94%)	0.29	9 (6%) 16 9	104, 149, 199, 226	0
2	H	132/138 (95%)	0.50	18 (13%) 3 2	30, 156, 238, 249	0
2	I	131/138 (94%)	0.42	14 (10%) 6 3	120, 156, 268, 313	0
2	J	129/138 (93%)	0.45	13 (10%) 7 4	30, 198, 234, 249	0
2	K	132/138 (95%)	1.10	34 (25%) 0 0	183, 231, 275, 298	0
2	L	129/138 (93%)	0.47	11 (8%) 10 6	30, 167, 224, 235	0
All	All	1672/1818 (91%)	0.43	124 (7%) 14 8	30, 154, 237, 313	0

All (124) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	36	PHE	5.0
2	L	35	THR	4.6
2	K	119	ASP	4.6
2	K	137	THR	4.6
2	K	102	CYS	4.6
2	L	31	CYS	4.5
2	L	33	ALA	4.5
2	L	78	CYS	4.5
1	F	152	VAL	4.4
2	H	84	ALA	4.3
2	J	59	GLN	4.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	K	126	ASN	4.1
2	I	71	VAL	3.9
1	F	222	ALA	3.9
2	G	153	GLU	3.9
2	K	57	GLY	3.8
2	I	59	GLN	3.8
2	G	133	CYS	3.8
1	D	176	ALA	3.8
2	I	34	GLY	3.8
2	L	37	CYS	3.8
2	K	111	GLU	3.8
2	J	72	PHE	3.8
2	I	56	ALA	3.8
2	G	54	SER	3.7
2	G	120	CYS	3.7
2	L	36	PHE	3.7
2	H	28	CYS	3.6
2	H	38	ASP	3.6
2	K	155	ASP	3.5
2	K	133	CYS	3.4
2	I	108	GLN	3.4
2	K	59	GLN	3.4
2	J	102	CYS	3.4
2	I	35	THR	3.3
2	K	112	LEU	3.3
2	G	59	GLN	3.3
2	K	35	THR	3.2
2	H	85	GLU	3.2
2	H	39	ASN	3.2
2	K	139	CYS	3.1
1	F	200	GLN	3.1
2	H	37	CYS	3.1
2	K	148	VAL	3.0
2	J	58	GLY	3.0
2	H	59	GLN	3.0
2	K	127	ASP	3.0
1	E	176	ALA	3.0
2	K	28	CYS	3.0
2	K	120	CYS	3.0
2	J	36	PHE	3.0
2	K	117	CYS	3.0
2	H	95	LEU	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	J	125	PHE	3.0
2	H	125	PHE	2.9
2	G	136	TRP	2.9
2	K	110	GLN	2.9
1	D	115	LEU	2.8
2	L	102	CYS	2.8
2	H	108	GLN	2.8
2	I	55	SER	2.8
2	J	56	ALA	2.8
2	K	71	VAL	2.7
2	K	43	GLN	2.7
2	L	38	ASP	2.7
1	F	231	GLY	2.7
2	H	57	GLY	2.7
2	I	72	PHE	2.7
1	E	118	VAL	2.7
1	C	90	GLY	2.6
1	F	192	ALA	2.6
2	J	78	CYS	2.6
2	I	78	CYS	2.6
2	L	65	CYS	2.6
2	I	54	SER	2.5
2	K	39	ASN	2.5
2	G	146	VAL	2.5
2	K	48	CYS	2.4
2	J	65	CYS	2.4
2	G	55	SER	2.4
2	K	136	TRP	2.4
2	H	43	GLN	2.4
1	B	175	GLY	2.4
2	H	40	ASN	2.4
2	L	34	GLY	2.4
2	K	142	ASP	2.4
2	K	125	PHE	2.3
1	E	135	VAL	2.3
2	J	126	ASN	2.3
1	F	91	MET	2.3
2	K	37	CYS	2.3
2	K	157	VAL	2.3
2	K	56	ALA	2.3
2	H	71	VAL	2.3
2	K	54	SER	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	I	36	PHE	2.3
1	F	154	ALA	2.3
2	H	45	CYS	2.3
1	C	242	PRO	2.2
2	J	52	SER	2.2
1	E	233	THR	2.2
2	K	26	ASP	2.2
2	K	32	PRO	2.2
1	B	177	ALA	2.2
1	C	236	GLY	2.2
2	H	64	ILE	2.2
2	I	84	ALA	2.2
1	A	242	PRO	2.2
1	F	201	GLY	2.2
2	K	31	CYS	2.2
2	G	124	THR	2.2
2	K	109	GLY	2.1
2	I	33	ALA	2.1
1	C	198	GLY	2.1
2	I	86	CYS	2.1
1	B	197	PHE	2.1
2	K	33	ALA	2.1
2	H	33	ALA	2.1
1	E	201	GLY	2.1
1	B	231	GLY	2.1
2	L	58	GLY	2.1
2	J	37	CYS	2.1
2	J	92	PHE	2.1
1	E	146	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	G	500	14/15	0.79	0.25	190,198,201,201	0
3	NAG	H	501	14/15	0.80	0.20	186,195,202,204	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.