



# Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 04:46 am BST

PDB ID : 1CWO  
Title : HUMAN CYCLOPHILIN A COMPLEXED WITH THR2, LEU5, D-HIV8,  
LEU10 CYCLOSPORIN  
Authors : Mikol, V.; Kallen, J.; Taylor, P.; Walkinshaw, M.D.  
Deposited on : 1998-06-05  
Resolution : 1.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	165	
2	C	11	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2293 atoms, of which 723 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PEPTIDYL-PROLYL CIS-TRANS ISOMERASE A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	165	Total	C	H	N	O	S	0	0	0
			1551	802	285	218	237	9			

- Molecule 2 is a protein called CYCLOSPORIN C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	11	Total	C	H	N	O	0	0	0
			94	64	6	10	14			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	1	VAD	DAL	ENGINEERED MUTATION	NOR NOR00035
C	3	LEU	MLE	ENGINEERED MUTATION	NOR NOR00035
C	9	LEU	VAL	ENGINEERED MUTATION	NOR NOR00035

- Molecule 3 is water.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	210	Total	H	O	0	0
			630	420	210		
3	C	6	Total	H	O	0	0
			18	12	6		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: PEPTIDYL-PROLYL CIS-TRANS ISOMERASE A

Chain A: 



- Molecule 2: CYCLOSPORIN C

Chain C: 



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	36.41Å 57.44Å 71.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 1.86	Depositor
% Data completeness (in resolution range)	95.7 (8.00-1.86)	Depositor
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, $R_{free}$	(Not available) , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2293	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MVA, SAR, BMT, VAD, MLE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.49	0/1294	0.71	0/1733
2	C	0.34	0/24	0.70	0/27
All	All	0.49	0/1318	0.71	0/1760

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1266	285	1237	16	0
2	C	88	6	114	6	0
3	A	210	420	0	8	1
3	C	6	12	0	0	0
All	All	1570	723	1351	21	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:100:MET:CE	1:A:100:MET:SD	2.01	1.47
1:A:52:CYS:HB2	3:A:2098:HOH:O	1.87	0.73
1:A:52:CYS:SG	1:A:157:THR:HG22	2.38	0.63
1:A:129:PHE:HA	3:A:2165:HOH:O	1.99	0.60
2:C:10:MLE:CD2	2:C:11:ALA:H	2.16	0.59
1:A:70:HIS:HD2	3:A:2096:HOH:O	1.88	0.56
1:A:4:PRO:HG2	1:A:24:LEU:HB2	1.86	0.56
1:A:25:PHE:HB2	3:A:2165:HOH:O	2.05	0.56
1:A:90:LEU:HD12	3:A:2165:HOH:O	2.06	0.55
2:C:10:MLE:HD23	2:C:11:ALA:H	1.74	0.53
1:A:102:ASN:HD22	1:A:102:ASN:C	2.13	0.52
1:A:1:MET:HE1	3:A:2006:HOH:O	2.13	0.48
2:C:1:VAD:HA	2:C:2:MLE:HN1	1.51	0.48
1:A:102:ASN:ND2	1:A:104:GLY:H	2.12	0.47
1:A:113:PHE:CD1	2:C:4:MVA:HG11	2.49	0.47
1:A:98:LEU:HG	1:A:129:PHE:CZ	2.51	0.45
1:A:155:LYS:HD3	3:A:2197:HOH:O	2.17	0.44
2:C:9:LEU:HA	2:C:10:MLE:HN1	1.47	0.43
1:A:15:GLU:HA	1:A:16:PRO:HD3	1.92	0.41
1:A:70:HIS:CD2	3:A:2096:HOH:O	2.69	0.41
2:C:5:BMT:HD22	2:C:10:MLE:HN2	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:2075:HOH:H1	3:A:2196:HOH:H1[4_577]	1.26	0.34

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	163/165 (99%)	159 (98%)	4 (2%)	0	<b>100</b> <b>100</b>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	3/11 (27%)	3 (100%)	0	0	100	100
All	All	166/176 (94%)	162 (98%)	4 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	133/133 (100%)	128 (96%)	5 (4%)	33	16
2	C	3/3 (100%)	3 (100%)	0	100	100
All	All	136/136 (100%)	131 (96%)	5 (4%)	34	17

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	61	MET
1	A	76	LYS
1	A	81	GLU
1	A	102	ASN
1	A	151	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	HIS
1	A	102	ASN
1	A	149	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.



## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

7 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	MLE	C	2	2	7,8,9	0.39	0	6,9,11	0.89	0
2	MLE	C	10	2	7,8,9	0.45	0	6,9,11	1.52	1 (16%)
2	BMT	C	5	2	11,12,13	0.66	0	12,14,16	0.92	1 (8%)
2	SAR	C	7	2	4,4,5	0.75	0	1,3,5	1.69	0
2	VAD	C	1	2	5,6,7	1.55	1 (20%)	6,7,9	1.29	1 (16%)
2	MLE	C	8	2	7,8,9	0.49	0	6,9,11	1.32	1 (16%)
2	MVA	C	4	2	6,7,8	0.68	0	7,8,10	1.59	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MLE	C	2	2	-	1/5/8/10	-
2	MLE	C	10	2	-	3/5/8/10	-
2	BMT	C	5	2	-	1/13/16/18	-
2	SAR	C	7	2	-	1/1/2/3	-
2	VAD	C	1	2	-	2/5/6/8	-
2	MLE	C	8	2	-	1/5/8/10	-
2	MVA	C	4	2	-	2/6/8/10	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1	VAD	CB-CA	2.85	1.57	1.53

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	4	MVA	CB-CA-C	-3.16	109.07	113.04
2	C	1	VAD	OE-CA-CB	-3.00	103.18	109.64
2	C	10	MLE	CG-CB-CA	2.96	122.71	115.34
2	C	8	MLE	CG-CB-CA	2.50	121.57	115.34
2	C	5	BMT	CG2-CD2-CE	2.40	117.67	113.98

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	2	MLE	O-C-CA-CB
2	C	10	MLE	C-CA-CB-CG
2	C	10	MLE	O-C-CA-CB
2	C	1	VAD	C-CA-CB-CG2
2	C	10	MLE	N-CA-CB-CG
2	C	8	MLE	N-CA-CB-CG
2	C	5	BMT	CB-CA-N-CN
2	C	4	MVA	CB-CA-N-CN
2	C	1	VAD	C-CA-CB-CG1
2	C	4	MVA	O-C-CA-CB
2	C	7	SAR	C-CA-N-CN

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	2	MLE	1	0
2	C	10	MLE	4	0
2	C	5	BMT	1	0
2	C	1	VAD	1	0
2	C	4	MVA	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.