



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 9, 2020 – 09:39 PM BST

PDB ID : 1CX2
Title : CYCLOOXYGENASE-2 (PROSTAGLANDIN SYNTHASE-2) COM-
PLEXED WITH A SELECTIVE INHIBITOR, SC-558
Authors : Kurumbail, R.; Stallings, W.
Deposited on : 1996-12-17
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	587	
1	B	587	
1	C	587	
1	D	587	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	S58	A	701	-	-	X	-
4	S58	C	701	-	-	X	-
4	S58	D	701	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 22376 atoms, of which 4040 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYCLOOXYGENASE-2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	552	Total	C	H	N	O	S	0	0	0
			5439	2886	966	748	814	25			
1	B	552	Total	C	H	N	O	S	0	0	0
			5439	2886	966	748	814	25			
1	C	552	Total	C	H	N	O	S	0	0	0
			5439	2886	966	748	814	25			
1	D	552	Total	C	H	N	O	S	0	0	0
			5439	2886	966	748	814	25			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	310	GLN	ASN	conflict	UNP Q05769
A	333	LYS	ARG	conflict	UNP Q05769
B	310	GLN	ASN	conflict	UNP Q05769
B	333	LYS	ARG	conflict	UNP Q05769
C	310	GLN	ASN	conflict	UNP Q05769
C	333	LYS	ARG	conflict	UNP Q05769
D	310	GLN	ASN	conflict	UNP Q05769
D	333	LYS	ARG	conflict	UNP Q05769

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
2	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
2	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
2	B	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
2	B	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
2	B	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
2	C	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
2	C	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
2	C	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
2	D	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
2	D	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
2	D	1	Total	C	H	N	O	0	0
			28	8	14	1	5		

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




-
- The chemical structure of S58 is a benzimidazole derivative. It features a central benzimidazole core. The imidazole ring is substituted with a trifluoromethyl group (CF₃) at the 2-position. The benzimidazole core is further substituted with a sulfonamide group (SO₂NH₂) at the 4-position and a 4-bromophenyl group at the 5-position. The bromine atom is highlighted in red. The atoms are labeled with their respective element symbols and indices: C1-C16 for carbon, N1-N3 for nitrogen, O1-O2 for oxygen, F1-F3 for fluorine, and Br1 for bromine. The sulfonamide group is labeled S1, O1, O2, and N3.

Mol	Chain	Residues	Atoms								ZeroOcc	AltConf
4	A	1	Total	Br	C	F	H	N	O	S	0	0
			28	1	16	3	2	3	2	1		
4	B	1	Total	Br	C	F	H	N	O	S	0	0
			28	1	16	3	2	3	2	1		
4	C	1	Total	Br	C	F	H	N	O	S	0	0
			28	1	16	3	2	3	2	1		
4	D	1	Total	Br	C	F	H	N	O	S	0	0
			28	1	16	3	2	3	2	1		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: CYCLOOXYGENASE-2



• Molecule 1: CYCLOOXYGENASE-2








• Molecule 1: CYCLOOXYGENASE-2

Chain D: 31% 52% 11% 6%



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	180.34Å 133.92Å 121.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 3.00 20.00 – 3.00	Depositor EDS
% Data completeness (in resolution range)	54.0 (8.00-3.00) 61.9 (20.00-3.00)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.66 (at 2.98Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.216 , 0.218 0.231 , 0.311	Depositor DCC
R_{free} test set	3652 reflections (9.98%)	wwPDB-VP
Wilson B-factor (Å ²)	28.9	Xtriage
Anisotropy	0.960	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 62.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.38$, $\langle L^2 \rangle = 0.20$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.84	EDS
Total number of atoms	22376	wwPDB-VP
Average B, all atoms (Å ²)	9.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 39.14 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.3164e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, NAG, S58

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	0/4600	0.88	4/6237 (0.1%)
1	B	0.71	0/4600	0.88	4/6237 (0.1%)
1	C	0.69	0/4600	0.88	3/6237 (0.0%)
1	D	0.72	0/4600	0.89	3/6237 (0.0%)
All	All	0.71	0/18400	0.88	14/24948 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	2
1	C	0	2
1	D	0	4
All	All	0	11

There are no bond length outliers.

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	575	CYS	N-CA-C	-6.66	93.02	111.00
1	D	575	CYS	N-CA-C	-6.47	93.54	111.00
1	C	575	CYS	N-CA-C	-6.38	93.78	111.00
1	B	575	CYS	N-CA-C	-6.35	93.85	111.00
1	D	287	VAL	N-CA-C	5.94	127.03	111.00

There are no chirality outliers.

5 of 11 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	134	TYR	Sidechain
1	A	262	TYR	Sidechain
1	A	348	TYR	Sidechain
1	B	262	TYR	Sidechain
1	B	348	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4473	966	4375	399	0
1	B	4473	966	4375	409	0
1	C	4473	966	4375	389	0
1	D	4473	966	4375	392	0
2	A	42	42	39	1	0
2	B	42	42	39	6	0
2	C	42	42	39	1	0
2	D	42	42	39	7	0
3	A	43	0	30	3	0
3	B	43	0	30	6	0
3	C	43	0	30	7	0
3	D	43	0	30	5	0
4	A	26	2	11	9	0
4	B	26	2	11	5	0
4	C	26	2	11	11	0
4	D	26	2	11	7	0
All	All	18336	4040	17820	1552	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 43.

The worst 5 of 1552 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:52:PHE:N	1:D:322:GLU:HG2	1.80	0.96
1:C:322:GLU:HG2	1:D:52:PHE:N	1.83	0.93
1:A:275:TYR:CE2	1:A:284:GLN:HA	2.04	0.93

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:184:ARG:HA	1:C:438:ARG:O	1.71	0.89
1:C:275:TYR:CE2	1:C:284:GLN:HA	2.07	0.89

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	550/587 (94%)	439 (80%)	86 (16%)	25 (4%)	2	14
1	B	550/587 (94%)	438 (80%)	85 (16%)	27 (5%)	2	13
1	C	550/587 (94%)	439 (80%)	84 (15%)	27 (5%)	2	13
1	D	550/587 (94%)	444 (81%)	78 (14%)	28 (5%)	2	12
All	All	2200/2348 (94%)	1760 (80%)	333 (15%)	107 (5%)	2	13

5 of 107 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	67	GLU
1	A	138	SER
1	A	226	HIS
1	A	270	GLN
1	A	284	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	493/525 (94%)	422 (86%)	71 (14%)	3	15
1	B	493/525 (94%)	411 (83%)	82 (17%)	2	11
1	C	493/525 (94%)	423 (86%)	70 (14%)	3	16
1	D	493/525 (94%)	418 (85%)	75 (15%)	3	14
All	All	1972/2100 (94%)	1674 (85%)	298 (15%)	3	14

5 of 298 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	428	ARG
1	C	150	ARG
1	D	412	SER
1	B	469	ARG
1	B	577	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 71 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	351	HIS
1	C	87	ASN
1	D	318	GLN
1	B	382	ASN
1	B	464	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry

20 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	671	1	14,14,15	0.87	1 (7%)	17,19,21	1.18	1 (5%)
2	NAG	D	671	1	14,14,15	0.60	0	17,19,21	1.15	3 (17%)
2	NAG	A	671	1	14,14,15	0.55	0	17,19,21	1.13	3 (17%)
2	NAG	B	671	1	14,14,15	0.62	0	17,19,21	0.92	1 (5%)
2	NAG	D	661	1	14,14,15	0.64	0	17,19,21	0.76	0
2	NAG	D	681	1	14,14,15	0.55	0	17,19,21	0.92	1 (5%)
4	S58	C	701	-	27,28,28	2.65	6 (22%)	37,43,43	1.51	6 (16%)
2	NAG	B	681	1	14,14,15	0.70	0	17,19,21	0.81	0
3	HEM	B	682	1	27,50,50	1.93	6 (22%)	17,82,82	1.29	3 (17%)
2	NAG	B	661	1	14,14,15	0.63	0	17,19,21	0.84	1 (5%)
3	HEM	D	682	1	27,50,50	1.78	5 (18%)	17,82,82	1.14	2 (11%)
2	NAG	C	661	1	14,14,15	0.93	0	17,19,21	0.72	1 (5%)
2	NAG	C	681	1	14,14,15	0.88	1 (7%)	17,19,21	0.88	1 (5%)
4	S58	D	701	-	27,28,28	2.76	8 (29%)	37,43,43	1.62	8 (21%)
3	HEM	A	682	1	27,50,50	1.71	5 (18%)	17,82,82	1.20	1 (5%)
2	NAG	A	681	1	14,14,15	0.84	0	17,19,21	0.95	1 (5%)
3	HEM	C	682	1	27,50,50	1.75	6 (22%)	17,82,82	1.32	3 (17%)
4	S58	B	701	-	27,28,28	2.67	7 (25%)	37,43,43	1.65	7 (18%)
2	NAG	A	661	1	14,14,15	0.75	0	17,19,21	0.88	0
4	S58	A	701	-	27,28,28	2.54	6 (22%)	37,43,43	1.63	8 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	671	1	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	D	671	1	-	1/6/23/26	0/1/1/1
2	NAG	A	671	1	-	0/6/23/26	0/1/1/1
2	NAG	B	671	1	-	0/6/23/26	0/1/1/1
2	NAG	D	661	1	-	1/6/23/26	0/1/1/1
2	NAG	D	681	1	-	2/6/23/26	0/1/1/1
4	S58	C	701	-	-	8/20/20/20	0/3/3/3
2	NAG	B	681	1	-	3/6/23/26	0/1/1/1
3	HEM	B	682	1	-	1/6/54/54	-
2	NAG	B	661	1	-	2/6/23/26	0/1/1/1
3	HEM	D	682	1	-	1/6/54/54	-
2	NAG	C	661	1	-	1/6/23/26	0/1/1/1
2	NAG	C	681	1	-	3/6/23/26	0/1/1/1
4	S58	D	701	-	-	8/20/20/20	0/3/3/3
3	HEM	A	682	1	-	1/6/54/54	-
2	NAG	A	681	1	-	2/6/23/26	0/1/1/1
3	HEM	C	682	1	-	1/6/54/54	-
4	S58	B	701	-	-	8/20/20/20	0/3/3/3
2	NAG	A	661	1	-	2/6/23/26	0/1/1/1
4	S58	A	701	-	-	9/20/20/20	0/3/3/3

The worst 5 of 51 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	701	S58	S1-N3	8.09	1.76	1.60
4	C	701	S58	C1-C3	8.04	1.50	1.39
4	D	701	S58	C1-C3	7.98	1.50	1.39
4	B	701	S58	S1-N3	7.98	1.76	1.60
4	A	701	S58	S1-N3	7.89	1.76	1.60

The worst 5 of 51 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	701	S58	O2-S1-O1	-5.50	109.71	118.76
4	B	701	S58	O2-S1-O1	-5.32	110.02	118.76
4	D	701	S58	O2-S1-O1	-5.10	110.38	118.76
4	C	701	S58	O2-S1-O1	-4.39	111.55	118.76
4	B	701	S58	C4-C3-N2	3.69	124.06	119.72

There are no chirality outliers.

5 of 54 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	682	HEM	C2A-CAA-CBA-CGA
3	D	682	HEM	C2A-CAA-CBA-CGA
3	A	682	HEM	C2A-CAA-CBA-CGA
3	C	682	HEM	C2A-CAA-CBA-CGA
2	B	681	NAG	O5-C5-C6-O6


There are no ring outliers.


14 monomers are involved in 68 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	671	NAG	1	0
2	D	661	NAG	5	0
2	D	681	NAG	2	0
4	C	701	S58	11	0
2	B	681	NAG	4	0
3	B	682	HEM	6	0
2	B	661	NAG	2	0
3	D	682	HEM	5	0
2	C	661	NAG	1	0
4	D	701	S58	7	0
3	A	682	HEM	3	0
3	C	682	HEM	7	0
4	B	701	S58	5	0
4	A	701	S58	9	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	552/587 (94%)	-0.54	0 100 100	2, 6, 16, 30	0
1	B	552/587 (94%)	-0.57	1 (0%) 95 87	2, 6, 16, 27	0
1	C	552/587 (94%)	-0.55	0 100 100	2, 6, 16, 26	0
1	D	552/587 (94%)	-0.52	1 (0%) 95 87	2, 7, 17, 30	0
All	All	2208/2348 (94%)	-0.55	2 (0%) 95 89	2, 6, 16, 30	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	105	ASN	2.6
1	B	583	GLN	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	A	661	14/15	0.80	0.29	12,17,20,21	0
2	NAG	C	661	14/15	0.81	0.31	12,17,21,22	0
2	NAG	B	681	14/15	0.81	0.28	11,17,22,24	0
2	NAG	D	681	14/15	0.84	0.42	12,17,20,24	0
2	NAG	C	681	14/15	0.85	0.29	11,17,21,22	0
2	NAG	D	661	14/15	0.88	0.30	15,17,19,21	0
2	NAG	A	681	14/15	0.89	0.23	7,17,18,21	0
2	NAG	B	661	14/15	0.89	0.29	15,17,19,20	0
2	NAG	C	671	14/15	0.93	0.15	7,13,17,17	0
2	NAG	D	671	14/15	0.93	0.15	5,11,17,17	0
2	NAG	A	671	14/15	0.93	0.16	4,12,17,17	0
2	NAG	B	671	14/15	0.93	0.17	5,12,17,17	0
3	HEM	B	682	43/43	0.94	0.19	2,3,7,9	0
3	HEM	D	682	43/43	0.94	0.17	2,3,7,9	0
3	HEM	A	682	43/43	0.95	0.17	2,3,8,10	0
3	HEM	C	682	43/43	0.96	0.14	2,3,7,8	0
4	S58	C	701	26/26	0.97	0.13	2,8,15,17	0
4	S58	D	701	26/26	0.97	0.12	2,8,13,17	0
4	S58	A	701	26/26	0.98	0.13	2,9,15,17	0
4	S58	B	701	26/26	0.98	0.12	2,8,14,17	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around HEM B 682:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM D 682:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM A 682:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM C 682:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around S58 C 701:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around S58 D 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around S58 A 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around S58 B 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.