



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 10, 2021 – 03:36 PM EDT

PDB ID : 3CX4
Title : Crystal Structure of E.coli GS mutant E377A in complex with ADP and oligosaccharides
Authors : Sheng, F.; Geiger, J.H.
Deposited on : 2008-04-23
Resolution : 2.29 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	485	<div> <div>10%</div> <div>76%</div> <div>20%</div> <div>..</div> </div>
2	B	3	<div> <div>67%</div> <div>33%</div> </div>
3	C	2	<div> <div>50%</div> <div>50%</div> </div>
4	D	5	<div> <div>20%</div> <div>80%</div> </div>
5	E	6	<div> <div>67%</div> <div>33%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GLC	C	2	-	-	-	X
4	GLC	D	1	-	-	-	X
9	ETE	A	521	-	-	X	-
9	ETE	A	522	-	-	X	X
9	ETE	A	527	-	-	-	X
9	ETE	A	528	-	-	X	-

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 4285 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycogen synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	476	3751	2415	656	666	14	13	4	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	377	ALA	GLU	engineered mutation	UNP P0A6U8
A	478	LEU	-	expression tag	UNP P0A6U8
A	479	GLU	-	expression tag	UNP P0A6U8
A	480	HIS	-	expression tag	UNP P0A6U8
A	481	HIS	-	expression tag	UNP P0A6U8
A	482	HIS	-	expression tag	UNP P0A6U8
A	483	HIS	-	expression tag	UNP P0A6U8
A	484	HIS	-	expression tag	UNP P0A6U8
A	485	HIS	-	expression tag	UNP P0A6U8

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	B	3	34	18	16	0	0	0

- Molecule 3 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	C	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 4 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
4	D	5	Total	C	O	0	0	0
			56	30	26			

- Molecule 5 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
5	E	6	Total	C	O	0	0	0
			67	36	31			

- Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	Na	0	0
			2	2		


- Molecule 7 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂).



- 250
-
- ORTEP diagram of the chemical structure of 1,4-bis(2-hydroxyethyl)piperazine-2-sulfonic acid. The structure shows a central piperazine ring with two nitrogen atoms, N1 and N2. N1 is bonded to a 2-hydroxyethyl group (C3-C2-C1) and a sulfonic acid group (S1-O1-O2-O3). N2 is bonded to another 2-hydroxyethyl group (C8-C9-C10). The hydroxyl groups are shown in red, and the sulfonic acid group is shown in yellow. The piperazine ring is shown in blue. The structure is labeled with atom names and numbers: N1, N2, C1, C2, C3, C4, C5, C6, C7, C8, C9, C10, O1, O2, O3, O4, O5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	A	1	Total	C	N	O	S	0	0
			17	9	2	5	1		

- Molecule 9 is 2-{2-[2-2-(METHOXY-ETHOXY)-ETHOXY]-ETHOXY}-ETHANOL (three-letter code: ETE) (formula: C₉H₂₀O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	C	O	0	0
			14	9	5		
9	A	1	Total	C	O	0	0
			11	7	4		
9	A	1	Total	C	O	0	0
			14	9	5		
9	A	1	Total	C	O	0	0
			12	8	4		
9	A	1	Total	C	O	0	0
			6	4	2		
9	A	1	Total	C	O	0	0
			4	2	2		
9	A	1	Total	C	O	0	0
			5	3	2		
9	A	1	Total	C	O	0	0
			11	8	3		
9	A	1	Total	C	O	0	0
			8	5	3		
9	A	1	Total	C	O	0	0
			6	4	2		
9	A	1	Total	C	O	0	0
			10	7	3		
9	A	1	Total	C	O	0	0
			6	4	2		

- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	201	Total 201	O 201	0	0



- Molecule 5: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranose

Chain E: 67% 33%



4 Data and refinement statistics

Property	Value	Source
Space group	I 41	Depositor
Cell constants a, b, c, α , β , γ	126.32Å 126.32Å 152.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	9.31 – 2.29 32.61 – 2.29	Depositor EDS
% Data completeness (in resolution range)	99.8 (9.31-2.29) 99.9 (32.61-2.29)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.92 (at 2.29Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.186 , 0.221 0.190 , 0.223	Depositor DCC
R_{free} test set	2728 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	49.0	Xtriage
Anisotropy	0.271	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 67.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.023 for -h,k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4285	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP, 250, ETE, BGC, GLC, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	0/3865	0.68	3/5258 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	396	ARG	NE-CZ-NH2	-6.71	116.94	120.30
1	A	267	ARG	NE-CZ-NH2	-6.41	117.09	120.30
1	A	48	ARG	NE-CZ-NH2	-5.90	117.35	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3751	0	3698	128	0
2	B	34	0	30	4	0
3	C	23	0	21	0	0
4	D	56	0	48	1	0
5	E	67	0	57	3	0
6	A	2	0	0	0	0
7	A	27	0	12	0	0
8	A	17	0	20	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	A	107	0	123	56	0
10	A	201	0	0	12	0
All	All	4285	0	4009	139	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 139 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:A:517:250:H42	9:A:527:ETE:H132	1.23	1.16
1:A:252:ILE:CD1	9:A:528:ETE:H141	1.79	1.12
1:A:380:GLY:H	9:A:521:ETE:H242	1.16	1.10
1:A:252:ILE:HD11	9:A:528:ETE:H141	1.22	1.09
1:A:21:ASP:OD1	9:A:528:ETE:H252	1.58	1.01

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	478/485 (99%)	457 (96%)	21 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	384/389 (99%)	357 (93%)	27 (7%)	15	19

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	228	MET
1	A	259	LEU
1	A	430	SER
1	A	252	ILE
1	A	261	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	233	GLN
1	A	246	ASN
1	A	454	GLN
1	A	383	GLN
1	A	412	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

16 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	B	1	2	12,12,12	0.61	0	17,17,17	1.02	1 (5%)
2	GLC	B	2	2	11,11,12	0.61	0	15,15,17	1.76	3 (20%)
2	GLC	B	3	2	11,11,12	0.62	0	15,15,17	1.73	4 (26%)
3	GLC	C	1	3	12,12,12	0.64	0	17,17,17	1.33	3 (17%)
3	GLC	C	2	3	11,11,12	0.59	0	15,15,17	0.80	0
4	GLC	D	1	4	12,12,12	0.56	0	17,17,17	0.70	0
4	GLC	D	2	4	11,11,12	0.67	0	15,15,17	0.73	0
4	GLC	D	3	4	11,11,12	0.78	0	15,15,17	1.18	2 (13%)
4	GLC	D	4	4	11,11,12	0.65	0	15,15,17	1.03	1 (6%)
4	GLC	D	5	4	11,11,12	0.52	0	15,15,17	0.93	1 (6%)
5	BGC	E	1	5	12,12,12	0.59	0	17,17,17	1.56	4 (23%)
5	GLC	E	2	5	11,11,12	0.60	0	15,15,17	1.14	1 (6%)
5	GLC	E	3	5	11,11,12	0.69	0	15,15,17	1.04	1 (6%)
5	GLC	E	4	5	11,11,12	0.82	0	15,15,17	1.28	2 (13%)
5	GLC	E	5	5	11,11,12	0.48	0	15,15,17	1.24	2 (13%)
5	GLC	E	6	5	11,11,12	0.76	0	15,15,17	1.20	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2	-	0/2/22/22	0/1/1/1
2	GLC	B	2	2	-	2/2/19/22	0/1/1/1
2	GLC	B	3	2	-	0/2/19/22	0/1/1/1
3	GLC	C	1	3	-	0/2/22/22	0/1/1/1
3	GLC	C	2	3	-	2/2/19/22	0/1/1/1
4	GLC	D	1	4	-	0/2/22/22	0/1/1/1
4	GLC	D	2	4	-	0/2/19/22	0/1/1/1
4	GLC	D	3	4	-	0/2/19/22	0/1/1/1
4	GLC	D	4	4	-	2/2/19/22	0/1/1/1
4	GLC	D	5	4	-	2/2/19/22	0/1/1/1
5	BGC	E	1	5	-	1/2/22/22	0/1/1/1
5	GLC	E	2	5	-	0/2/19/22	0/1/1/1
5	GLC	E	3	5	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GLC	E	4	5	-	0/2/19/22	0/1/1/1
5	GLC	E	5	5	-	0/2/19/22	0/1/1/1
5	GLC	E	6	5	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	GLC	C1-O5-C5	4.60	118.42	112.19
5	E	2	GLC	C1-O5-C5	3.37	116.75	112.19
5	E	1	BGC	O5-C5-C4	3.34	115.76	109.69
3	C	1	GLC	O5-C5-C4	3.22	115.54	109.69
5	E	1	BGC	C3-C4-C5	3.18	115.92	110.24

There are no chirality outliers.

5 of 9 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	2	GLC	O5-C5-C6-O6
4	D	5	GLC	O5-C5-C6-O6
2	B	2	GLC	C4-C5-C6-O6
4	D	5	GLC	C4-C5-C6-O6
4	D	4	GLC	C4-C5-C6-O6

There are no ring outliers.


4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	3	GLC	4	0
5	E	1	BGC	3	0
5	E	2	GLC	3	0
4	D	2	GLC	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.




Oligosaccharide Chain C



Oligosaccharide Chain D





5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 2 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
9	ETE	A	531	-	5,5,13	0.52	0	4,4,12	0.09	0
9	ETE	A	525	-	3,3,13	0.43	0	2,2,12	0.32	0
9	ETE	A	527	-	10,10,13	0.52	0	9,9,12	0.24	0
9	ETE	A	528	-	7,7,13	0.51	0	6,6,12	0.64	0
7	ADP	A	500	6	24,29,29	0.83	0	29,45,45	1.23	2 (6%)
9	ETE	A	530	-	9,9,13	0.57	0	8,8,12	0.15	0
8	250	A	517	-	17,17,17	0.87	1 (5%)	21,23,23	2.57	11 (52%)
9	ETE	A	524	-	5,5,13	0.49	0	4,4,12	0.23	0
9	ETE	A	522	-	13,13,13	0.59	0	12,12,12	0.20	0
9	ETE	A	523	-	11,11,13	0.55	0	10,10,12	0.21	0
9	ETE	A	529	-	5,5,13	0.51	0	4,4,12	0.32	0
9	ETE	A	521	-	10,10,13	0.52	0	9,9,12	0.33	0
9	ETE	A	520	-	13,13,13	0.46	0	12,12,12	0.33	0
9	ETE	A	526	-	4,4,13	0.49	0	3,3,12	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	ETE	A	531	-	-	3/3/3/11	-
9	ETE	A	525	-	-	1/1/1/11	-
9	ETE	A	527	-	-	5/8/8/11	-
9	ETE	A	528	-	-	5/5/5/11	-
7	ADP	A	500	6	-	4/12/32/32	0/3/3/3
9	ETE	A	530	-	-	4/7/7/11	-
8	250	A	517	-	-	4/12/22/22	0/1/1/1
9	ETE	A	524	-	-	1/3/3/11	-
9	ETE	A	522	-	-	8/11/11/11	-
9	ETE	A	523	-	-	6/9/9/11	-
9	ETE	A	529	-	-	2/3/3/11	-
9	ETE	A	521	-	-	5/8/8/11	-
9	ETE	A	520	-	-	5/11/11/11	-
9	ETE	A	526	-	-	1/2/2/11	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	A	517	250	C1-S1	-2.66	1.67	1.77

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	517	250	C5-N2-C6	4.94	119.95	108.83
7	A	500	ADP	N3-C2-N1	-4.70	121.34	128.68
8	A	517	250	C4-N1-C7	4.13	118.12	108.83
8	A	517	250	C8-N2-C6	3.77	120.88	111.23
8	A	517	250	C7-N1-C3	3.65	121.81	111.20

There are no chirality outliers.

5 of 54 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
9	A	520	ETE	C14-C24-OH4-C13
9	A	522	ETE	OH4-C13-C23-OH3
9	A	520	ETE	OH4-C13-C23-OH3
9	A	520	ETE	OH6-C15-C25-OH5
9	A	530	ETE	OH5-C14-C24-OH4


There are no ring outliers.

11 monomers are involved in 56 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	A	531	ETE	1	0
9	A	527	ETE	5	0
9	A	528	ETE	21	0
9	A	530	ETE	1	0
8	A	517	250	3	0
9	A	524	ETE	1	0
9	A	522	ETE	8	0
9	A	523	ETE	1	0
9	A	521	ETE	14	0
9	A	520	ETE	3	0
9	A	526	ETE	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	476/485 (98%)	0.34	49 (10%) 6 9	35, 47, 68, 80	34 (7%)

The worst 5 of 49 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	98	THR	6.4
1	A	99	ASN	6.2
1	A	6	VAL	4.7
1	A	113	LEU	4.4
1	A	135	ALA	4.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GLC	C	1	12/12	0.77	0.30	90,91,91,92	1
3	GLC	C	2	11/12	0.78	0.47	92,93,95,95	0
4	GLC	D	1	12/12	0.79	0.50	66,67,68,68	12
2	GLC	B	1	12/12	0.82	0.36	50,51,51,52	12
2	GLC	B	2	11/12	0.84	0.21	50,52,55,57	0
5	BGC	E	1	12/12	0.88	0.28	72,75,75,76	12
4	GLC	D	5	11/12	0.91	0.34	79,81,82,83	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	GLC	D	4	11/12	0.91	0.22	67,69,74,76	0
4	GLC	D	2	11/12	0.92	0.21	60,62,63,64	0
5	GLC	E	3	11/12	0.93	0.26	66,68,68,68	0
2	GLC	B	3	11/12	0.94	0.18	54,55,56,57	0
5	GLC	E	4	11/12	0.94	0.26	62,63,64,65	0
4	GLC	D	3	11/12	0.95	0.14	58,60,64,65	0
5	GLC	E	5	11/12	0.95	0.15	56,58,59,60	0
5	GLC	E	2	11/12	0.96	0.22	69,70,71,71	11
5	GLC	E	6	11/12	0.96	0.15	55,56,56,57	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
9	ETE	A	522	14/14	0.60	0.42	52,59,68,68	10
9	ETE	A	529	6/14	0.64	0.23	82,83,84,84	1
9	ETE	A	525	4/14	0.66	0.33	96,96,96,96	0
9	ETE	A	523	12/14	0.67	0.32	78,79,80,80	8
9	ETE	A	527	11/14	0.75	0.73	89,89,90,90	11

Continued on next page...




Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
9	ETE	A	531	6/14	0.77	0.23	93,94,95,95	0
9	ETE	A	530	10/14	0.80	0.23	100,101,103,103	0
8	250	A	517	17/17	0.80	0.24	99,101,103,103	5
9	ETE	A	526	5/14	0.82	0.37	86,87,87,88	0
9	ETE	A	521	11/14	0.85	0.55	55,58,61,61	5
9	ETE	A	520	14/14	0.86	0.18	69,70,75,75	9
9	ETE	A	528	8/14	0.88	0.39	8,10,12,12	8
9	ETE	A	524	6/14	0.92	0.34	94,94,94,94	0
7	ADP	A	500	27/27	0.96	0.13	43,47,48,49	0
6	NA	A	518	1/1	0.96	0.42	58,58,58,58	0
6	NA	A	519	1/1	0.97	0.19	58,58,58,58	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 250 A 517:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.