



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 6, 2020 – 02:20 PM BST

PDB ID : 4CYG
Title : The structure of vanin-1: defining the link between metabolic disease, oxidative stress and inflammation
Authors : Boersma, Y.L.; Newman, J.; Adams, T.E.; Sparrow, L.; Cowieson, N.; Lucent, D.; Krippner, G.; Bozaoglu, K.; Peat, T.S.
Deposited on : 2014-04-11
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	506	
1	B	506	
2	C	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	B	1527	-	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7820 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PANTETHEINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	463	Total	C	N	O	S	0	4	0
			3670	2340	606	704	20			
1	B	462	Total	C	N	O	S	0	5	0
			3675	2343	609	703	20			

There are 30 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	8	ALA	-	expression tag	UNP O95497
A	9	SER	-	expression tag	UNP O95497
A	10	ILE	-	expression tag	UNP O95497
A	11	SER	-	expression tag	UNP O95497
A	12	ALA	-	expression tag	UNP O95497
A	13	ARG	-	expression tag	UNP O95497
A	14	ASP	-	expression tag	UNP O95497
A	15	TYR	-	expression tag	UNP O95497
A	16	LYS	-	expression tag	UNP O95497
A	17	ASP	-	expression tag	UNP O95497
A	18	ASP	-	expression tag	UNP O95497
A	19	ASP	-	expression tag	UNP O95497
A	20	ASP	-	expression tag	UNP O95497
A	21	LYS	-	expression tag	UNP O95497
A	26	ILE	THR	variant	UNP O95497
B	8	ALA	-	expression tag	UNP O95497
B	9	SER	-	expression tag	UNP O95497
B	10	ILE	-	expression tag	UNP O95497
B	11	SER	-	expression tag	UNP O95497
B	12	ALA	-	expression tag	UNP O95497
B	13	ARG	-	expression tag	UNP O95497
B	14	ASP	-	expression tag	UNP O95497
B	15	TYR	-	expression tag	UNP O95497
B	16	LYS	-	expression tag	UNP O95497
B	17	ASP	-	expression tag	UNP O95497

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	18	ASP	-	expression tag	UNP O95497
B	19	ASP	-	expression tag	UNP O95497
B	20	ASP	-	expression tag	UNP O95497
B	21	LYS	-	expression tag	UNP O95497
B	26	ILE	THR	variant	UNP O95497

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is (2R)-2,4-dihydroxy-N-[(3S)-3-hydroxy-4-phenylbutyl]-3,3-dimethylbutanamide (three-letter code: RRV) (formula: C₁₆H₂₅NO₄).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			21	16	1	4		
3	B	1	Total	C	N	O	0	0
			21	16	1	4		

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			7	4	3		
5	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	181	Total	O	0	0
			181	181		
6	B	126	Total	O	0	0
			126	126		

- Molecule 1: PANTETHEINASE

ILE	VAL	ILE	ALA	PRO	ILE	VAL	CYS	LEU	SER	SER	TRP	ALA	SER	ILE	SER	ALA	ARG	LYS	ASP	ASP	ASP	ASP	K21	L52	I62	Y63	S132	P150	Q151	R157	N186	R200	C211	R259	G276	A312	N424	D483	T49	ALA	SER	ASN	ALA	SER	SER	GLY	LEU	THR	ALA	GLN	ALA	ARG	ILE	ILE	ILE	MET	LEU
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

- Molecule 1: PANTETHEINASE

TRP	ALA
SER	SER
ASN	ILE
ALA	ALA
SER	ARG
SER	ASP
GLY	TVR
LEU	LYS
THR	ASP
ALA	ASP
GLN	ASP
ARG	LYS
ILE	Q22
MET	A39
LEU	T40
ILE	L41
VAL	T42
ILE	Y83
ALA	Y83
PRO	N105
ILE	N105
VAL	S132
CYS	S132
SER	D146
LEU	T147
SER	P150
TRP	P150
	R157
	K171
	C211
	D218
	K258
	R259
	G276
	V313
	K316
	E324
	V350
	K372
	V436
	N445
	D462

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

NAG1
NAG2

4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	120.09 Å 120.09 Å 221.90 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	105.62 – 2.30 46.62 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.4 (105.62-2.30) 99.5 (46.62-2.30)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.40 (at 2.29 Å)	Xtriage
Refinement program	REFMAC 5.8.0069	Depositor
R, R_{free}	0.186 , 0.214 0.193 , 0.217	Depositor DCC
R_{free} test set	3638 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	36.0	Xtriage
Anisotropy	0.024	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 27.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7820	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: RRV, PEG, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/3766	0.57	0/5128
1	B	0.35	0/3771	0.57	0/5134
All	All	0.35	0/7537	0.57	0/10262

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3670	0	3512	5	0
1	B	3675	0	3522	8	0
2	C	28	0	25	2	0
3	A	21	0	25	1	0
3	B	21	0	25	1	0
4	A	42	0	39	1	0
4	B	42	0	39	0	0
5	B	14	0	20	0	0
6	A	181	0	0	2	0
6	B	126	0	0	1	0
All	All	7820	0	7207	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:211:CYS:SG	3:A:1502:RRV:H12	1.94	1.06
1:B:211:CYS:SG	3:B:1501:RRV:H12	1.98	1.03
1:B:350:VAL:HG12	1:B:372:ASN:HB2	1.89	0.54
1:A:52:LEU:HD13	4:A:1524:NAG:C8	2.43	0.49
1:B:350:VAL:CG1	1:B:372:ASN:HB2	2.43	0.48
2:C:2:NAG:C8	2:C:2:NAG:O3	2.66	0.43
1:A:200[A]:ASN:ND2	6:A:2102:HOH:O	2.20	0.43
1:A:82[B]:ILE:HG22	6:A:2049:HOH:O	2.19	0.43
1:B:105:ASN:OD1	1:B:171:LYS:HE2	2.19	0.43
1:B:259[A]:ARG:HH21	1:B:313:VAL:HG22	1.83	0.42
1:B:258:MET:O	1:B:259[A]:ARG:HG2	2.20	0.42
1:A:259:ARG:NH2	1:A:312:ALA:O	2.52	0.41
2:C:2:NAG:H83	2:C:2:NAG:O3	2.20	0.41
1:B:218:ASP:OD1	6:B:2042:HOH:O	2.22	0.40
1:B:316:TRP:CE2	1:B:436:VAL:HB	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	465/506 (92%)	448 (96%)	15 (3%)	2 (0%)	34	42
1	B	465/506 (92%)	450 (97%)	13 (3%)	2 (0%)	34	42
All	All	930/1012 (92%)	898 (97%)	28 (3%)	4 (0%)	34	42

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	132	SER
1	B	132	SER
1	A	276	GLY
1	B	276	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	406/438 (93%)	401 (99%)	5 (1%)	71	84
1	B	407/438 (93%)	402 (99%)	5 (1%)	71	84
All	All	813/876 (93%)	803 (99%)	10 (1%)	71	84

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	83	TYR
1	A	132	SER
1	A	157	ARG
1	A	186	ASN
1	A	424	MET
1	B	83	TYR
1	B	132	SER
1	B	157	ARG
1	B	324	GLU
1	B	445	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	117	GLN
1	A	186	ASN
1	A	434	GLN
1	A	445	ASN
1	B	445	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	1	1,2	14,14,15	0.47	0	17,19,21	1.63	4 (23%)
2	NAG	C	2	2	14,14,15	0.97	1 (7%)	17,19,21	3.06	8 (47%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	C	2	2	-	5/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	2	NAG	C1-C2	2.71	1.56	1.52

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	NAG	C2-N2-C7	8.32	134.74	122.90
2	C	2	NAG	C8-C7-N2	6.13	126.47	116.10
2	C	1	NAG	C2-N2-C7	3.48	127.86	122.90
2	C	2	NAG	O7-C7-C8	-3.46	115.63	122.06
2	C	1	NAG	C8-C7-N2	3.39	121.84	116.10
2	C	2	NAG	O5-C1-C2	-2.59	107.20	111.29
2	C	2	NAG	C4-C3-C2	2.45	114.61	111.02
2	C	2	NAG	O3-C3-C4	-2.45	104.69	110.35
2	C	2	NAG	C1-O5-C5	2.23	115.21	112.19
2	C	2	NAG	O7-C7-N2	-2.21	117.89	121.95
2	C	1	NAG	C1-O5-C5	2.20	115.17	112.19
2	C	1	NAG	O7-C7-C8	-2.18	118.00	122.06

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	2	NAG	C3-C2-N2-C7
2	C	2	NAG	O5-C5-C6-O6
2	C	2	NAG	C4-C5-C6-O6
2	C	1	NAG	C8-C7-N2-C2
2	C	1	NAG	O7-C7-N2-C2
2	C	2	NAG	C8-C7-N2-C2
2	C	2	NAG	O7-C7-N2-C2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	2	NAG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	RRV	A	1502	1	19,21,21	0.71	0	24,28,28	0.93	1 (4%)
4	NAG	B	1527	1	14,14,15	0.61	0	17,19,21	1.20	1 (5%)
4	NAG	A	1522	1	14,14,15	0.49	0	17,19,21	1.10	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	A	1512	1	14,14,15	0.73	0	17,19,21	2.01	4 (23%)
5	PEG	B	1484	-	6,6,6	0.57	0	5,5,5	0.19	0
4	NAG	B	1525	1	14,14,15	0.43	0	17,19,21	1.79	3 (17%)
4	NAG	B	1522	1	14,14,15	0.43	0	17,19,21	1.36	1 (5%)
5	PEG	B	1485	-	6,6,6	0.52	0	5,5,5	0.24	0
3	RRV	B	1501	1	19,21,21	0.75	1 (5%)	24,28,28	0.91	2 (8%)
4	NAG	A	1524	1	14,14,15	0.56	0	17,19,21	1.11	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	RRV	A	1502	1	-	5/23/23/23	0/1/1/1
4	NAG	B	1527	1	-	1/6/23/26	0/1/1/1
4	NAG	A	1522	1	-	2/6/23/26	0/1/1/1
4	NAG	A	1512	1	-	1/6/23/26	0/1/1/1
5	PEG	B	1484	-	-	2/4/4/4	-
4	NAG	B	1525	1	-	2/6/23/26	0/1/1/1
4	NAG	B	1522	1	-	0/6/23/26	0/1/1/1
5	PEG	B	1485	-	-	2/4/4/4	-
3	RRV	B	1501	1	-	2/23/23/23	0/1/1/1
4	NAG	A	1524	1	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1501	RRV	C7-N9	2.18	1.38	1.33

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1512	NAG	O5-C1-C2	-5.34	102.85	111.29
4	B	1525	NAG	C1-O5-C5	4.74	118.62	112.19
4	B	1522	NAG	C1-O5-C5	4.28	117.99	112.19
4	A	1512	NAG	C3-C4-C5	3.83	117.07	110.24
4	B	1527	NAG	C4-C3-C2	3.45	116.07	111.02
4	B	1525	NAG	C8-C7-N2	3.40	121.86	116.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1512	NAG	C4-C3-C2	3.39	115.99	111.02
3	A	1502	RRV	C6-C7-N9	-2.78	111.03	116.58
4	A	1524	NAG	O5-C1-C2	-2.48	107.37	111.29
4	A	1522	NAG	C1-O5-C5	2.45	115.50	112.19
3	B	1501	RRV	C6-C7-N9	-2.39	111.82	116.58
4	A	1512	NAG	C1-C2-N2	2.35	114.50	110.49
4	A	1524	NAG	C4-C3-C2	-2.25	107.72	111.02
4	B	1525	NAG	C4-C3-C2	-2.18	107.82	111.02
3	B	1501	RRV	C1-C2-C6	2.17	112.59	108.82
4	A	1522	NAG	O5-C5-C6	2.01	110.36	107.20

There are no chirality outliers.

All (17) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1502	RRV	C11-C12-C15-C16
3	A	1502	RRV	O13-C12-C15-C16
3	B	1501	RRV	C11-C12-C15-C16
3	B	1501	RRV	O13-C12-C15-C16
4	B	1525	NAG	C8-C7-N2-C2
4	B	1525	NAG	O7-C7-N2-C2
4	A	1522	NAG	O5-C5-C6-O6
5	B	1485	PEG	O2-C3-C4-O4
4	A	1522	NAG	C4-C5-C6-O6
5	B	1485	PEG	O1-C1-C2-O2
5	B	1484	PEG	O1-C1-C2-O2
4	B	1527	NAG	O5-C5-C6-O6
5	B	1484	PEG	O2-C3-C4-O4
4	A	1512	NAG	O5-C5-C6-O6
3	A	1502	RRV	O8-C7-N9-C10
3	A	1502	RRV	O22-C6-C7-O8
3	A	1502	RRV	C6-C7-N9-C10

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1502	RRV	1	0
3	B	1501	RRV	1	0
4	A	1524	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	463/506 (91%)	-0.42	2 (0%) 92 95	21, 32, 56, 86	0
1	B	462/506 (91%)	-0.30	6 (1%) 77 81	21, 34, 61, 101	0
All	All	925/1012 (91%)	-0.36	8 (0%) 84 88	21, 33, 60, 101	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	41	LEU	5.5
1	B	39	ALA	3.6
1	B	150	PRO	2.8
1	A	151	GLN	2.7
1	B	147	THR	2.7
1	B	42	THR	2.5
1	A	150	PRO	2.5
1	B	146	ASP	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	C	2	14/15	0.75	0.38	78,89,93,98	0
2	NAG	C	1	14/15	0.95	0.13	39,46,55,56	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	B	1527	14/15	0.71	0.44	88,104,110,110	0
5	PEG	B	1485	7/7	0.73	0.17	62,63,67,68	0
5	PEG	B	1484	7/7	0.74	0.26	68,70,73,75	0
4	NAG	A	1512	14/15	0.85	0.29	53,76,89,93	0
4	NAG	B	1525	14/15	0.89	0.15	63,69,73,73	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	RRV	A	1502	21/21	0.90	0.15	38,47,50,52	0
4	NAG	A	1522	14/15	0.90	0.14	54,59,62,62	0
3	RRV	B	1501	21/21	0.90	0.13	45,53,56,57	0
4	NAG	B	1522	14/15	0.91	0.16	57,59,62,64	0
4	NAG	A	1524	14/15	0.96	0.10	35,39,41,43	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.