



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 23, 2020 – 04:56 PM JST

PDB ID : 7CY7  
Title : Crystal Structure of CMD1 in complex with DNA  
Authors : Li, W.; Zhang, T.; Sun, M.; Ding, J.  
Deposited on : 2020-09-03  
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.16  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.16


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	917	<div> <div>4%</div> <div>87%</div> <div>8%</div> <div>5%</div> </div>
2	C	14	<div> <div>36%</div> <div>64%</div> <div>36%</div> </div>
2	D	14	<div> <div>21%</div> <div>50%</div> <div>50%</div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7408 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltodextrin-binding protein,5-methylcytosine-modifying enzyme 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	872	Total	C	N	O	S	0	3	0
			6642	4222	1143	1250	27			


There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-373	MET	-	initiating methionine	UNP A0A376KDN7
A	-291	ALA	ASP	engineered mutation	UNP A0A376KDN7
A	-290	ALA	LYS	engineered mutation	UNP A0A376KDN7
A	-201	ALA	GLU	engineered mutation	UNP A0A376KDN7
A	-200	ALA	ASN	engineered mutation	UNP A0A376KDN7
A	-134	ALA	LYS	engineered mutation	UNP A0A376KDN7
A	-11	ALA	LYS	engineered mutation	UNP A0A376KDN7
A	-10	ALA	ASP	engineered mutation	UNP A0A376KDN7
A	-6	ASN	-	linker	UNP A0A376KDN7
A	-5	ALA	-	linker	UNP A0A376KDN7
A	-4	ALA	-	linker	UNP A0A376KDN7
A	-3	ARG	-	linker	UNP A0A376KDN7
A	-2	ALA	-	linker	UNP A0A376KDN7
A	-1	ALA	-	linker	UNP A0A376KDN7
A	0	ALA	-	linker	UNP A0A376KDN7
A	533	ALA	-	expression tag	UNP A0A2K3D5Z7
A	534	ALA	-	expression tag	UNP A0A2K3D5Z7
A	535	ALA	-	expression tag	UNP A0A2K3D5Z7
A	536	LEU	-	expression tag	UNP A0A2K3D5Z7
A	537	GLU	-	expression tag	UNP A0A2K3D5Z7
A	538	HIS	-	expression tag	UNP A0A2K3D5Z7
A	539	HIS	-	expression tag	UNP A0A2K3D5Z7
A	540	HIS	-	expression tag	UNP A0A2K3D5Z7
A	541	HIS	-	expression tag	UNP A0A2K3D5Z7
A	542	HIS	-	expression tag	UNP A0A2K3D5Z7
A	543	HIS	-	expression tag	UNP A0A2K3D5Z7

- Molecule 2 is a DNA chain called DNA (5'-D(P\*CP\*GP\*CP\*GP\*CP\*GP\*GP\*GP\*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	9	Total	C	N	O	P	0	0	0
			188	87	39	53	9			
2	D	7	Total	C	N	O	P	0	0	0
			139	65	25	42	7			


- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is ISOPROPYL ALCOHOL (three-letter code: IPA) (formula: C<sub>3</sub>H<sub>8</sub>O) (labeled

as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	3	1		

- Molecule 5 is FE (II) ION (three-letter code: FE2) (formula: Fe) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Fe	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	395	Total	O	0	0
			395	395		
6	C	3	Total	O	0	0
			3	3		
6	D	4	Total	O	0	0
			4	4		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Maltodextrin-binding protein, 5-methylcytosine-modifying enzyme 1



- Molecule 2: DNA (5'-D(P\*CP\*GP\*CP\*GP\*CP\*GP\*GP\*GP\*A)-3')



- Molecule 2: DNA (5'-D(P\*CP\*GP\*CP\*GP\*CP\*GP\*GP\*GP\*A)-3')



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	153.61Å 125.76Å 64.03Å 90.00° 102.82° 90.00°	Depositor
Resolution (Å)	50.01 – 2.15 49.28 – 2.15	Depositor EDS
% Data completeness (in resolution range)	94.0 (50.01-2.15) 94.0 (49.28-2.15)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.12 (at 2.14Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
R, $R_{free}$	0.188 , 0.221 0.190 , 0.220	Depositor DCC
$R_{free}$ test set	3035 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.1	Xtriage
Anisotropy	0.035	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 41.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7408	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IPA, FE2, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.34	0/6796	0.54	0/9238
2	C	0.51	0/211	0.69	0/324
2	D	0.37	0/154	0.77	0/234
All	All	0.35	0/7161	0.55	0/9796

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6642	0	6584	35	0
2	C	188	0	100	0	0
2	D	139	0	78	0	0
3	A	32	0	48	0	0
4	A	4	0	8	0	0
5	A	1	0	0	0	0
6	A	395	0	0	0	0
6	C	3	0	0	0	0
6	D	4	0	0	0	0
All	All	7408	0	6818	35	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:394:THR:HG21	1:A:497:SER:HB2	1.81	0.63
1:A:26:ARG:HG3	1:A:105:ASP:OD1	1.98	0.62
1:A:-311:TRP:HB3	1:A:-306:PHE:HE1	1.68	0.57
1:A:325:MET:HE2	1:A:332:ASN:HB3	1.87	0.57
1:A:41:THR:O	1:A:45:GLU:HG2	2.05	0.57
1:A:-86:ASP:OD1	1:A:-67:SER:OG	2.24	0.55
1:A:-305:GLY:HA3	1:A:-41:ASN:O	2.07	0.54
1:A:239:PHE:HB3	1:A:252:ASN:HB3	1.89	0.54
1:A:-122:LYS:O	1:A:-120[B]:GLN:NE2	2.42	0.52
1:A:187:PHE:HB3	1:A:501:TRP:CZ2	2.44	0.52
1:A:-329:GLU:HB3	1:A:-311:TRP:CZ2	2.44	0.52
1:A:-365:VAL:HG13	1:A:-316:PRO:HA	1.91	0.52
1:A:-275:ARG:HG3	1:A:-270:LEU:HD23	1.91	0.52
1:A:12:VAL:HG21	1:A:487:VAL:HG21	1.93	0.51
1:A:127:ARG:NH1	1:A:184:GLY:O	2.44	0.49
1:A:27:LYS:O	1:A:31:ILE:HG12	2.12	0.49
1:A:-294:ILE:HG22	1:A:-292:PRO:HD3	1.94	0.48
1:A:-243:GLU:O	1:A:-240:PRO:HD2	2.15	0.47
1:A:-214:PRO:HG3	1:A:-116:PRO:HA	1.97	0.46
1:A:419:PHE:HB2	1:A:486:VAL:HG21	1.97	0.46
1:A:350:ASP:HA	1:A:417:LEU:O	2.15	0.46
1:A:-123:PHE:CE2	1:A:-122:LYS:HD2	2.51	0.45
1:A:458:GLY:O	1:A:462:ARG:HG3	2.17	0.45
1:A:325:MET:CE	1:A:332:ASN:HB3	2.47	0.45
1:A:91:PRO:HG3	1:A:377:GLY:HA3	1.98	0.45
1:A:369:GLY:HA2	1:A:400:ARG:O	2.17	0.45
1:A:-352:ALA:O	1:A:-348:LYS:HG2	2.18	0.43
1:A:-311:TRP:HB3	1:A:-306:PHE:CE1	2.51	0.42
1:A:62:LYS:HG3	1:A:160:CYS:HB2	2.01	0.41
1:A:496:PRO:HG2	1:A:499:LEU:HD21	2.03	0.41
1:A:-363:TRP:CD1	1:A:-316:PRO:HB3	2.56	0.41
1:A:-296:ALA:HB2	1:A:-100:LYS:HE3	2.02	0.41
1:A:419:PHE:HA	1:A:422:VAL:HG12	2.02	0.41
1:A:339:ASN:HB3	1:A:402:LEU:HD11	2.03	0.40
1:A:-102:PRO:HG2	1:A:-101:ASN:ND2	2.37	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	871/917 (95%)	854 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	682/711 (96%)	664 (97%)	18 (3%)	46	47

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-367	LYS
1	A	-342	THR
1	A	-327	LYS
1	A	-140	SER
1	A	-120[A]	GLN
1	A	-120[B]	GLN
1	A	-95	GLU
1	A	-71	VAL
1	A	27	LYS
1	A	28	SER
1	A	219	ILE
1	A	244	ARG
1	A	251	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	261	ASN
1	A	266	ILE
1	A	312	TYR
1	A	484	ARG
1	A	487	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 1 is monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	EDO	A	602	-	3,3,3	0.45	0	2,2,2	0.22	0
3	EDO	A	606	-	3,3,3	0.45	0	2,2,2	0.40	0
3	EDO	A	608	-	3,3,3	0.57	0	2,2,2	0.09	0
3	EDO	A	603	-	3,3,3	0.44	0	2,2,2	0.22	0
3	EDO	A	607	-	3,3,3	0.48	0	2,2,2	0.30	0
4	IPA	A	609	-	3,3,3	0.67	0	3,3,3	0.25	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	EDO	A	601	-	3,3,3	0.47	0	2,2,2	0.28	0
3	EDO	A	604	-	3,3,3	0.52	0	2,2,2	0.03	0
3	EDO	A	605	-	3,3,3	0.44	0	2,2,2	0.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	602	-	-	0/1/1/1	-
3	EDO	A	606	-	-	1/1/1/1	-
3	EDO	A	608	-	-	1/1/1/1	-
3	EDO	A	603	-	-	0/1/1/1	-
3	EDO	A	607	-	-	0/1/1/1	-
3	EDO	A	601	-	-	0/1/1/1	-
3	EDO	A	604	-	-	0/1/1/1	-
3	EDO	A	605	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	606	EDO	O1-C1-C2-O2
3	A	608	EDO	O1-C1-C2-O2
3	A	605	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	872/917 (95%)	0.30	37 (4%) 36 45	17, 36, 68, 89	0
2	C	9/14 (64%)	2.36	5 (55%) 0 0	91, 104, 158, 160	0
2	D	7/14 (50%)	1.94	3 (42%) 0 0	60, 68, 132, 147	0
All	All	888/945 (93%)	0.34	45 (5%) 28 36	17, 36, 69, 160	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	-321	ALA	4.9
1	A	-232	ALA	4.7
2	C	6	DG	4.7
2	C	4	DG	4.2
2	D	1	DC	4.0
1	A	-317	GLY	3.9
1	A	457	VAL	3.9
1	A	184	GLY	3.5
1	A	-371	ILE	3.5
2	C	3	DC	3.4
1	A	453	SER	3.2
2	D	7	DC	3.2
1	A	-171	LYS	3.2
1	A	456	GLY	3.2
1	A	-272	GLY	2.8
1	A	-201	ALA	2.8
1	A	-293	THR	2.8
1	A	-338	VAL	2.7
1	A	-289	ALA	2.7
1	A	-238	LEU	2.7
1	A	-173	LYS	2.7
1	A	-236	LYS	2.6
1	A	-97	ALA	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	-90	TYR	2.5
1	A	-340	ILE	2.5
1	A	455	ASP	2.4
2	C	5	DC	2.4
1	A	-347	LYS	2.4
1	A	-346	PHE	2.4
1	A	-337	THR	2.4
1	A	-367	LYS	2.3
1	A	491	CYS	2.3
1	A	449	ASP	2.3
1	A	-177	VAL	2.3
1	A	462	ARG	2.2
1	A	-235	GLU	2.2
1	A	458	GLY	2.2
1	A	-249	ASN	2.2
1	A	459	ALA	2.2
2	C	11	DA	2.1
1	A	-342	THR	2.1
1	A	176	ALA	2.1
2	D	2	DC	2.1
1	A	452	ASN	2.1
1	A	-166	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	EDO	A	601	4/4	0.40	0.35	71,73,75,75	0

*Continued on next page...*






*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	EDO	A	606	4/4	0.76	0.27	62,66,66,68	0
3	EDO	A	604	4/4	0.76	0.30	50,50,51,55	0
4	IPA	A	609	4/4	0.83	0.29	53,55,55,58	0
3	EDO	A	607	4/4	0.87	0.27	52,52,53,55	0
3	EDO	A	608	4/4	0.91	0.21	49,52,53,55	0
3	EDO	A	605	4/4	0.92	0.21	39,39,39,41	0
3	EDO	A	602	4/4	0.94	0.15	43,46,49,50	0
3	EDO	A	603	4/4	0.95	0.14	41,43,43,44	0
5	FE2	A	610	1/1	0.95	0.10	44,44,44,44	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around IPA A 609:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around FE2 A 610:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.