



wwPDB X-ray Structure Validation Summary Report ⓘ

May 28, 2020 – 09:27 pm BST

PDB ID : 2D00
Title : Subunit F of V-type ATPase/synthase
Authors : Makyio, H.; Iino, R.; Ikeda, C.; Imamura, H.; Tamakoshi, M.; Iwata, M.;
Stock, D.; Bernal, R.A.; Carpenter, E.P.; Yoshida, M.; Yokoyama, K.; Iwata,
S.
Deposited on : 2005-07-21
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	109	67% 28% 6%
1	B	109	56% 39% 6%
1	C	109	54% 39% 5% .
1	D	109	63% 32% . .
1	E	109	61% 33% . .
1	F	109	60% 33% 5% .

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5289 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called V-type ATP synthase subunit F.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	109	Total	C	N	O	S	0	0	0
			833	527	144	158	4			
1	B	109	Total	C	N	O	S	0	0	0
			833	527	144	158	4			
1	C	109	Total	C	N	O	S	0	0	0
			833	527	144	158	4			
1	D	109	Total	C	N	O	S	0	0	0
			833	527	144	158	4			
1	E	109	Total	C	N	O	S	0	0	0
			833	527	144	158	4			
1	F	109	Total	C	N	O	S	0	0	0
			833	527	144	158	4			

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	CLONING ARTIFACT	UNP P74903
A	2	VAL	-	CLONING ARTIFACT	UNP P74903
A	3	PRO	-	CLONING ARTIFACT	UNP P74903
A	4	VAL	-	CLONING ARTIFACT	UNP P74903
A	5	ARG	-	CLONING ARTIFACT	UNP P74903
B	1	MET	-	CLONING ARTIFACT	UNP P74903
B	2	VAL	-	CLONING ARTIFACT	UNP P74903
B	3	PRO	-	CLONING ARTIFACT	UNP P74903
B	4	VAL	-	CLONING ARTIFACT	UNP P74903
B	5	ARG	-	CLONING ARTIFACT	UNP P74903
C	1	MET	-	CLONING ARTIFACT	UNP P74903
C	2	VAL	-	CLONING ARTIFACT	UNP P74903
C	3	PRO	-	CLONING ARTIFACT	UNP P74903
C	4	VAL	-	CLONING ARTIFACT	UNP P74903
C	5	ARG	-	CLONING ARTIFACT	UNP P74903
D	1	MET	-	CLONING ARTIFACT	UNP P74903
D	2	VAL	-	CLONING ARTIFACT	UNP P74903

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	3	PRO	-	CLONING ARTIFACT	UNP P74903
D	4	VAL	-	CLONING ARTIFACT	UNP P74903
D	5	ARG	-	CLONING ARTIFACT	UNP P74903
E	1	MET	-	CLONING ARTIFACT	UNP P74903
E	2	VAL	-	CLONING ARTIFACT	UNP P74903
E	3	PRO	-	CLONING ARTIFACT	UNP P74903
E	4	VAL	-	CLONING ARTIFACT	UNP P74903
E	5	ARG	-	CLONING ARTIFACT	UNP P74903
F	1	MET	-	CLONING ARTIFACT	UNP P74903
F	2	VAL	-	CLONING ARTIFACT	UNP P74903
F	3	PRO	-	CLONING ARTIFACT	UNP P74903
F	4	VAL	-	CLONING ARTIFACT	UNP P74903
F	5	ARG	-	CLONING ARTIFACT	UNP P74903

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	2	Total Ca 2 2	0	0
2	D	1	Total Ca 1 1	0	0
2	C	1	Total Ca 1 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	54	Total O 54 54	0	0
3	B	45	Total O 45 45	0	0
3	C	51	Total O 51 51	0	0
3	D	42	Total O 42 42	0	0
3	E	49	Total O 49 49	0	0
3	F	46	Total O 46 46	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: V-type ATP synthase subunit F

Chain A: 




- Molecule 1: V-type ATP synthase subunit F

Chain B: 




- Molecule 1: V-type ATP synthase subunit F

Chain C: 




- Molecule 1: V-type ATP synthase subunit F


Chain D: 



- Molecule 1: V-type ATP synthase subunit F



● Molecule 1: V-type ATP synthase subunit F



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	81.70 Å 138.29 Å 66.10 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 2.20 24.77 – 2.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (25.00-2.20) 99.4 (24.77-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.71 (at 2.19 Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.214 , 0.262 0.223 , 0.268	Depositor DCC
R_{free} test set	1206 reflections (3.13%)	wwPDB-VP
Wilson B-factor (Å ²)	49.2	Xtriage
Anisotropy	0.025	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 76.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5289	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.60	9/845 (1.1%)	1.29	7/1140 (0.6%)
1	B	1.32	2/845 (0.2%)	1.18	2/1140 (0.2%)
1	C	2.10	23/845 (2.7%)	1.38	6/1140 (0.5%)
1	D	1.51	7/845 (0.8%)	1.31	10/1140 (0.9%)
1	E	1.55	8/845 (0.9%)	1.32	7/1140 (0.6%)
1	F	1.72	12/845 (1.4%)	1.27	6/1140 (0.5%)
All	All	1.65	61/5070 (1.2%)	1.29	38/6840 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	1
1	D	0	1
1	E	0	1
1	F	0	2
All	All	0	6

The worst 5 of 61 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	43	GLU	CD-OE1	26.75	1.55	1.25
1	F	97	GLU	CD-OE2	17.57	1.45	1.25
1	C	39	GLU	CD-OE2	15.09	1.42	1.25
1	C	39	GLU	CD-OE1	14.83	1.42	1.25
1	A	43	GLU	CG-CD	13.04	1.71	1.51

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	72	ASP	CB-CG-OD2	9.64	126.97	118.30
1	C	72	ASP	CB-CG-OD2	8.88	126.29	118.30
1	C	72	ASP	CB-CG-OD1	-8.58	110.58	118.30
1	C	68	MET	CG-SD-CE	-8.41	86.74	100.20
1	F	5	ARG	CG-CD-NE	-8.01	94.98	111.80

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	44	ARG	Peptide
1	C	44	ARG	Peptide
1	D	72	ASP	Peptide
1	E	71	ARG	Peptide
1	F	68	MET	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	833	0	850	35	1
1	B	833	0	850	48	1
1	C	833	0	850	46	0
1	D	833	0	850	30	0
1	E	833	0	850	47	0
1	F	833	0	850	38	0
2	B	2	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	54	0	0	8	0
3	B	45	0	0	18	0
3	C	51	0	0	7	0
3	D	42	0	0	8	0
3	E	49	0	0	17	1
3	F	46	0	0	8	1
All	All	5289	0	5100	211	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 211 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:68:MET:SD	1:C:68:MET:CG	2.07	1.41
1:A:37:LEU:HA	3:A:162:HOH:O	1.40	1.19
1:B:43:GLU:HG3	3:B:1043:HOH:O	1.48	1.14
1:F:32:GLU:HG3	3:F:117:HOH:O	1.58	1.02
1:E:74:PRO:HD3	3:E:153:HOH:O	1.59	1.02

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:92:GLU:OE2	3:E:157:HOH:O[2_665]	2.06	0.14
1:B:89:HIS:ND1	3:F:126:HOH:O[1_554]	2.08	0.12

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	107/109 (98%)	100 (94%)	6 (6%)	1 (1%)	17	16
1	B	107/109 (98%)	98 (92%)	6 (6%)	3 (3%)	5	2
1	C	107/109 (98%)	99 (92%)	7 (6%)	1 (1%)	17	16
1	D	107/109 (98%)	95 (89%)	11 (10%)	1 (1%)	17	16
1	E	107/109 (98%)	101 (94%)	4 (4%)	2 (2%)	8	5
1	F	107/109 (98%)	99 (92%)	6 (6%)	2 (2%)	8	5
All	All	642/654 (98%)	592 (92%)	40 (6%)	10 (2%)	9	7

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	89	HIS
1	B	45	GLY
1	F	105	PHE
1	B	40	THR
1	C	40	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/85 (100%)	81 (95%)	4 (5%)	26	33
1	B	85/85 (100%)	79 (93%)	6 (7%)	14	16
1	C	85/85 (100%)	81 (95%)	4 (5%)	26	33
1	D	85/85 (100%)	83 (98%)	2 (2%)	49	62
1	E	85/85 (100%)	82 (96%)	3 (4%)	36	46
1	F	85/85 (100%)	76 (89%)	9 (11%)	6	6
All	All	510/510 (100%)	482 (94%)	28 (6%)	21	26

5 of 28 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	49	LEU
1	D	102	THR
1	F	100	ARG
1	C	107	ILE
1	D	62	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	87	GLN
1	D	87	GLN
1	D	89	HIS
1	F	89	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.