



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 08:23 pm BST

PDB ID : 3D0T
Title : Structure of the BNB domain of the Hsp70 cochaperone Bag2
Authors : Xu, Z.; Nix, J.C.; Devlin, K.; Misra, S.
Deposited on : 2008-05-02
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1332 (2.56-2.52)
Ramachandran outliers	138981	1315 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)
RSRZ outliers	127900	1272 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	88	<div> <div>3%</div> <div>60% 27% 13%</div> </div>
1	B	88	<div> <div>6%</div> <div>63% 24% 13%</div> </div>
1	C	88	<div> <div>5%</div> <div>57% 27% 14%</div> </div>
1	D	88	<div> <div>6%</div> <div>56% 32% 11%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2483 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BAG family molecular chaperone regulator 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	77	Total	C	N	O	S	0	0	0
			609	378	108	120	3			
1	B	77	Total	C	N	O	S	0	0	0
			606	375	108	120	3			
1	C	76	Total	C	N	O	S	0	0	0
			602	373	107	119	3			
1	D	78	Total	C	N	O	S	0	0	0
			613	380	109	121	3			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	102	GLY	-	EXPRESSION TAG	UNP Q91YN9
A	103	ALA	-	EXPRESSION TAG	UNP Q91YN9
A	104	MET	-	EXPRESSION TAG	UNP Q91YN9
A	105	GLY	-	EXPRESSION TAG	UNP Q91YN9
A	106	SER	-	EXPRESSION TAG	UNP Q91YN9
B	102	GLY	-	EXPRESSION TAG	UNP Q91YN9
B	103	ALA	-	EXPRESSION TAG	UNP Q91YN9
B	104	MET	-	EXPRESSION TAG	UNP Q91YN9
B	105	GLY	-	EXPRESSION TAG	UNP Q91YN9
B	106	SER	-	EXPRESSION TAG	UNP Q91YN9
C	102	GLY	-	EXPRESSION TAG	UNP Q91YN9
C	103	ALA	-	EXPRESSION TAG	UNP Q91YN9
C	104	MET	-	EXPRESSION TAG	UNP Q91YN9
C	105	GLY	-	EXPRESSION TAG	UNP Q91YN9
C	106	SER	-	EXPRESSION TAG	UNP Q91YN9
D	102	GLY	-	EXPRESSION TAG	UNP Q91YN9
D	103	ALA	-	EXPRESSION TAG	UNP Q91YN9
D	104	MET	-	EXPRESSION TAG	UNP Q91YN9
D	105	GLY	-	EXPRESSION TAG	UNP Q91YN9
D	106	SER	-	EXPRESSION TAG	UNP Q91YN9


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	22	Total 22	O 22	0	0
2	B	8	Total 8	O 8	0	0
2	C	8	Total 8	O 8	0	0
2	D	15	Total 15	O 15	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: BAG family molecular chaperone regulator 2




- Molecule 1: BAG family molecular chaperone regulator 2



- Molecule 1: BAG family molecular chaperone regulator 2



- Molecule 1: BAG family molecular chaperone regulator 2



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	104.68 Å 104.68 Å 164.30 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.69 – 2.55 52.34 – 2.55	Depositor EDS
% Data completeness (in resolution range)	97.8 (39.69-2.55) 97.9 (52.34-2.55)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.42 (at 2.55 Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.253 , 0.284 0.251 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	49.7	Xtriage
Anisotropy	0.338	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 51.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.37$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2483	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/612	0.62	0/814
1	B	0.40	0/609	0.59	0/809
1	C	0.38	0/605	0.58	0/804
1	D	0.41	0/616	0.57	0/819
All	All	0.40	0/2442	0.59	0/3246

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	609	0	629	20	0
1	B	606	0	623	19	0
1	C	602	0	620	22	2
1	D	613	0	632	27	0
2	A	22	0	0	3	1
2	B	8	0	0	4	0
2	C	8	0	0	0	0
2	D	15	0	0	3	0
All	All	2483	0	2504	79	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:142:CYS:HB3	1:C:171:LYS:HE2	1.61	0.81
1:D:164:LEU:HD11	1:D:168:LYS:HE2	1.63	0.78
1:B:141:ALA:HB1	1:B:156:GLN:HB3	1.68	0.75
1:D:164:LEU:HD13	1:D:164:LEU:C	2.08	0.73
1:A:162:CYS:HA	2:B:41:HOH:O	1.90	0.71
1:D:151:VAL:HG22	1:D:152:ASP:N	2.08	0.69
1:D:165:GLU:HG2	2:D:40:HOH:O	1.95	0.66
1:D:164:LEU:HD13	1:D:164:LEU:O	1.96	0.65
1:C:166:ASP:O	1:C:170:ILE:HG12	1.97	0.64
1:C:115:ARG:O	1:C:119:GLU:HG3	2.01	0.61
1:B:106:SER:O	2:B:41:HOH:O	2.15	0.60
1:B:128:LEU:HG	1:B:181:ILE:HD11	1.83	0.60
1:A:189:LYS:H	1:B:180:ASN:ND2	1.99	0.60
1:A:189:LYS:H	1:B:180:ASN:HD22	1.51	0.59
1:D:141:ALA:HA	1:D:151:VAL:HG21	1.82	0.59
1:C:155:PHE:O	1:C:159:VAL:HG23	2.02	0.59
1:A:142:CYS:O	1:A:143:SER:C	2.41	0.59
1:A:123:LYS:HD2	2:A:20:HOH:O	2.03	0.59
1:D:171:LYS:O	1:D:175:GLU:HG3	2.03	0.58
1:C:128:LEU:HD23	1:C:128:LEU:O	2.03	0.58
1:C:138:LEU:HD13	1:D:117:ILE:HD13	1.86	0.58
1:D:164:LEU:CD1	1:D:168:LYS:HE2	2.34	0.57
1:C:164:LEU:HD22	1:C:164:LEU:N	2.21	0.55
1:A:126:ASP:OD1	2:A:42:HOH:O	2.18	0.54
1:C:132:LYS:O	1:C:136:MET:HG3	2.08	0.54
1:B:123:LYS:NZ	1:B:127:ASP:OD1	2.42	0.53
1:D:151:VAL:CG2	1:D:152:ASP:N	2.71	0.53
1:D:128:LEU:HG	1:D:181:ILE:CD1	2.38	0.53
1:D:186:LYS:HD3	2:D:7:HOH:O	2.08	0.53
1:A:173:ARG:NH2	1:B:118:ASP:OD1	2.35	0.53
1:A:128:LEU:O	1:A:128:LEU:HD23	2.08	0.52
1:D:164:LEU:C	1:D:164:LEU:CD1	2.76	0.52
1:B:128:LEU:HG	1:B:181:ILE:CD1	2.40	0.51
1:B:172:ARG:HA	1:B:175:GLU:OE1	2.10	0.51
1:A:156:GLN:O	1:A:160:ILE:HG12	2.12	0.50
1:C:153:GLN:HA	1:C:153:GLN:NE2	2.27	0.50
1:A:162:CYS:CA	2:B:41:HOH:O	2.51	0.50
1:B:156:GLN:O	1:B:160:ILE:HG12	2.13	0.49
1:B:171:LYS:O	1:B:175:GLU:HG3	2.12	0.49
1:D:151:VAL:HG22	1:D:152:ASP:H	1.76	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:174:LEU:HD11	1:D:117:ILE:HD12	1.95	0.49
1:C:134:HIS:O	1:C:138:LEU:HG	2.13	0.49
1:D:154:LYS:O	1:D:158:ILE:HG12	2.13	0.49
1:B:106:SER:HB3	2:B:41:HOH:O	2.12	0.48
1:B:159:VAL:O	1:B:162:CYS:HB2	2.13	0.48
1:C:154:LYS:O	1:C:158:ILE:HG13	2.13	0.47
1:C:164:LEU:CD2	1:C:164:LEU:H	2.27	0.47
1:C:164:LEU:HD22	1:C:164:LEU:H	1.79	0.47
1:B:132:LYS:HG2	1:B:136:MET:CE	2.44	0.47
1:A:184:SER:HA	1:B:184:SER:HA	1.97	0.47
1:C:153:GLN:HE21	1:C:153:GLN:HA	1.80	0.46
1:D:151:VAL:CG2	1:D:152:ASP:H	2.29	0.46
1:A:164:LEU:HD21	1:A:168:LYS:HE3	1.97	0.45
1:D:138:LEU:O	1:D:141:ALA:HB3	2.17	0.45
1:D:132:LYS:NZ	2:D:50:HOH:O	2.49	0.45
1:C:128:LEU:HD23	1:C:128:LEU:C	2.37	0.45
1:A:180:ASN:ND2	1:B:189:LYS:H	2.15	0.45
1:A:171:LYS:O	1:A:175:GLU:HG3	2.18	0.43
1:C:138:LEU:O	1:C:141:ALA:HB3	2.18	0.43
1:A:173:ARG:NH2	2:A:4:HOH:O	2.34	0.43
1:A:110:LEU:HD23	1:B:166:ASP:HB3	2.01	0.43
1:A:183:ASN:OD1	1:A:186:LYS:NZ	2.40	0.43
1:C:142:CYS:HB3	1:C:171:LYS:CE	2.40	0.43
1:A:110:LEU:HD22	1:B:170:ILE:HD11	2.01	0.43
1:D:141:ALA:HA	1:D:151:VAL:CG2	2.49	0.42
1:D:165:GLU:OE1	1:D:165:GLU:HA	2.19	0.42
1:C:164:LEU:CD2	1:C:164:LEU:N	2.81	0.42
1:D:115:ARG:O	1:D:119:GLU:HG3	2.19	0.42
1:A:134:HIS:O	1:A:137:SER:HB3	2.19	0.42
1:C:113:ALA:O	1:C:117:ILE:HG13	2.19	0.42
1:D:138:LEU:HD22	1:D:155:PHE:CG	2.55	0.41
1:A:128:LEU:C	1:A:128:LEU:HD23	2.41	0.41
1:C:107:GLU:HA	1:C:107:GLU:OE1	2.20	0.41
1:B:110:LEU:HD13	1:B:110:LEU:C	2.41	0.41
1:D:126:ASP:O	1:D:130:ASN:ND2	2.55	0.41
1:C:179:ARG:HH11	1:C:179:ARG:HG2	1.85	0.40
1:D:177:LEU:HD23	1:D:177:LEU:HA	1.88	0.40
1:D:118:ASP:O	1:D:122:SER:HB2	2.22	0.40
1:D:155:PHE:O	1:D:159:VAL:HG23	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:168:LYS:NZ	2:A:42:HOH:O[2_665]	1.73	0.47
1:C:108:GLU:OE2	1:C:108:GLU:OE2[12_556]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	73/88 (83%)	71 (97%)	2 (3%)	0	100	100
1	B	73/88 (83%)	73 (100%)	0	0	100	100
1	C	72/88 (82%)	71 (99%)	1 (1%)	0	100	100
1	D	74/88 (84%)	74 (100%)	0	0	100	100
All	All	292/352 (83%)	289 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	70/77 (91%)	67 (96%)	3 (4%)	29	39
1	B	69/77 (90%)	68 (99%)	1 (1%)	67	79
1	C	69/77 (90%)	66 (96%)	3 (4%)	29	39
1	D	70/77 (91%)	65 (93%)	5 (7%)	14	19
All	All	278/308 (90%)	266 (96%)	12 (4%)	29	39

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	130	ASN
1	A	136	MET
1	A	174	LEU
1	B	181	ILE
1	C	110	LEU
1	C	115	ARG
1	C	174	LEU
1	D	110	LEU
1	D	133	SER
1	D	136	MET
1	D	174	LEU
1	D	181	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	180	ASN
1	B	167	GLN
1	B	180	ASN
1	C	153	GLN
1	C	167	GLN
1	C	180	ASN
1	D	130	ASN
1	D	134	HIS
1	D	153	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	77/88 (87%)	0.36	3 (3%)	39 45	35, 50, 76, 90	0
1	B	77/88 (87%)	0.59	5 (6%)	18 22	38, 52, 77, 83	0
1	C	76/88 (86%)	0.39	4 (5%)	26 31	35, 54, 79, 90	0
1	D	78/88 (88%)	0.36	5 (6%)	19 22	36, 50, 78, 85	0
All	All	308/352 (87%)	0.42	17 (5%)	25 30	35, 52, 78, 90	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	152	ASP	4.6
1	B	164	LEU	3.9
1	D	112	HIS	3.7
1	B	143	SER	3.5
1	C	152	ASP	3.4
1	C	106	SER	2.9
1	B	142	CYS	2.8
1	A	153	GLN	2.7
1	D	151	VAL	2.6
1	A	119	GLU	2.6
1	D	105	GLY	2.4
1	C	143	SER	2.2
1	D	165	GLU	2.2
1	B	153	GLN	2.2
1	C	159	VAL	2.2
1	D	115	ARG	2.2
1	A	160	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.