



# Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 04:05 am BST

PDB ID : 5D15  
Title : Crystal structure of an adenylyl cyclase Ma1120 from Mycobacterium avium  
in complex with ATP and calcium ion  
Authors : Bharambe, N.G.; Barathy, D.V.; Suguna, K.  
Deposited on : 2015-08-03  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	177	<div> <div>5%</div> <div> <div></div> <div>86%</div> <div>5%</div> <div>8%</div> </div> </div>
1	B	177	<div> <div>3%</div> <div> <div></div> <div>88%</div> <div>5%</div> <div>7%</div> </div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5391 atoms, of which 2472 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	162	Total	C	H	N	O	S	0	19	0
			2494	789	1229	229	240	7			
1	B	164	Total	C	H	N	O	S	0	8	0
			2427	767	1201	224	229	6			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	99	PHE	-	expression tag	UNP V7LAR8
A	100	GLN	-	expression tag	UNP V7LAR8
A	101	GLY	-	expression tag	UNP V7LAR8
A	102	ALA	-	expression tag	UNP V7LAR8
A	103	MET	-	expression tag	UNP V7LAR8
A	104	GLY	-	expression tag	UNP V7LAR8
A	105	SER	-	expression tag	UNP V7LAR8
B	99	PHE	-	expression tag	UNP V7LAR8
B	100	GLN	-	expression tag	UNP V7LAR8
B	101	GLY	-	expression tag	UNP V7LAR8
B	102	ALA	-	expression tag	UNP V7LAR8
B	103	MET	-	expression tag	UNP V7LAR8
B	104	GLY	-	expression tag	UNP V7LAR8
B	105	SER	-	expression tag	UNP V7LAR8

- Molecule 2 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>13</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	P	0	1
			43	10	12	5	13	3		
2	B	1	Total	C	H	N	O	P	0	1
			86	20	24	10	26	6		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Ca	0	0
			1	1		
3	A	1	Total	Ca	0	0
			1	1		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			10	2	6	2		



- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	187	Total	O	0	0
			187	187		
5	B	142	Total	O	0	0
			142	142		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Cyclase



#### • Molecule 1: Cyclase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.53 Å   55.64 Å   55.07 Å 90.00°   109.79°   90.00°	Depositor
Resolution (Å)	37.92 – 1.50 37.92 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.7 (37.92-1.50) 99.7 (37.92-1.50)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.56 (at 1.50 Å)	Xtriage
Refinement program	PHENIX 1.10pre_2084	Depositor
R, $R_{free}$	0.195 , 0.212 0.196 , 0.211	Depositor DCC
$R_{free}$ test set	2339 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.4	Xtriage
Anisotropy	0.369	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 48.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.023 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5391	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.43% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, EDO, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.64	1/1347 (0.1%)	0.56	0/1809
1	B	0.36	0/1268	0.49	0/1704
All	All	0.52	1/2615 (0.0%)	0.53	0/3513

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	263	TYR	CE2-CZ	-5.56	1.31	1.38

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1265	1229	1159	4	0
1	B	1226	1201	1165	5	0
2	A	31	12	12	2	0
2	B	62	24	24	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	4	6	6	0	0
5	A	187	0	0	1	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	142	0	0	1	0
All	All	2919	2472	2366	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:301[A]:ATP:N6	1:B:210:LEU:O	2.09	0.86
1:A:234[C]:GLN:OE1	5:A:401:HOH:O	2.00	0.77
1:B:113:ASP:OD2	1:B:195[A]:ARG:NH1	2.36	0.58
1:A:234[B]:GLN:HB3	1:A:235:PRO:HD3	1.95	0.48
1:B:217[A]:MET:SD	1:B:256:LEU:CD1	3.04	0.46
1:B:203[B]:SER:OG	5:B:401:HOH:O	2.20	0.44
1:B:217[A]:MET:SD	1:B:256:LEU:HD13	2.60	0.42
2:A:301[A]:ATP:H2'	2:A:301[A]:ATP:N3	2.33	0.41
1:A:152:VAL:HG11	1:A:160[A]:MET:HE2	2.01	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	177/177 (100%)	176 (99%)	1 (1%)	0	100	100
1	B	168/177 (95%)	165 (98%)	3 (2%)	0	100	100
All	All	345/354 (98%)	341 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	132/141 (94%)	129 (98%)	3 (2%)	50	20
1	B	122/141 (86%)	119 (98%)	3 (2%)	47	18
All	All	254/282 (90%)	248 (98%)	6 (2%)	46	19

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	121	ASN
1	A	192	ILE
1	A	213	ARG
1	B	121	ASN
1	B	153	LYS
1	B	213	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	ATP	B	301[A]	1,3	26,33,33	0.90	1 (3%)	31,52,52	1.43	6 (19%)
2	ATP	A	301[A]	3	26,33,33	1.97	10 (38%)	31,52,52	1.85	10 (32%)
4	EDO	A	303	-	3,3,3	0.49	0	2,2,2	0.23	0
2	ATP	B	301[B]	3	26,33,33	0.92	1 (3%)	31,52,52	1.43	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ATP	B	301[A]	1,3	-	8/18/38/38	0/3/3/3
2	ATP	A	301[A]	3	-	9/18/38/38	0/3/3/3
4	EDO	A	303	-	-	1/1/1/1	-
2	ATP	B	301[B]	3	-	5/18/38/38	0/3/3/3

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301[A]	ATP	PG-O3G	-3.66	1.40	1.54
2	A	301[A]	ATP	C4-N3	-3.39	1.31	1.35
2	A	301[A]	ATP	PG-O2G	-3.15	1.42	1.54
2	A	301[A]	ATP	PA-O5'	-2.63	1.48	1.59
2	A	301[A]	ATP	C2-N3	-2.50	1.28	1.32
2	A	301[A]	ATP	PA-O1A	-2.48	1.42	1.50
2	A	301[A]	ATP	PB-O2B	-2.47	1.43	1.55
2	B	301[B]	ATP	C5-C4	2.36	1.47	1.40
2	B	301[A]	ATP	C5-C4	2.32	1.47	1.40
2	A	301[A]	ATP	O5'-C5'	-2.20	1.36	1.44
2	A	301[A]	ATP	PA-O2A	-2.19	1.45	1.55
2	A	301[A]	ATP	PB-O1B	-2.12	1.43	1.50

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301[A]	ATP	N3-C2-N1	-3.69	122.91	128.68
2	B	301[B]	ATP	N3-C2-N1	-3.64	122.99	128.68
2	A	301[A]	ATP	O5'-PA-O1A	-3.55	95.19	109.07
2	A	301[A]	ATP	C4-C5-N7	-3.36	105.90	109.40
2	A	301[A]	ATP	C2'-C3'-C4'	-3.09	96.64	102.64
2	B	301[B]	ATP	C3'-C2'-C1'	2.94	105.40	100.98
2	A	301[A]	ATP	C2-N1-C6	2.75	123.46	118.75
2	A	301[A]	ATP	C5'-C4'-C3'	-2.72	105.00	115.18
2	B	301[A]	ATP	C4-C5-N7	-2.71	106.58	109.40
2	A	301[A]	ATP	O4'-C4'-C5'	2.69	118.21	109.37
2	A	301[A]	ATP	O5'-C5'-C4'	-2.68	99.75	108.99
2	A	301[A]	ATP	C1'-N9-C4	-2.67	121.95	126.64
2	B	301[B]	ATP	C4-C5-N7	-2.61	106.67	109.40
2	A	301[A]	ATP	N3-C2-N1	-2.55	124.69	128.68
2	B	301[A]	ATP	C3'-C2'-C1'	2.47	104.70	100.98
2	B	301[B]	ATP	PA-O3A-PB	-2.25	125.11	132.83
2	B	301[A]	ATP	PB-O3B-PG	-2.24	125.13	132.83
2	B	301[A]	ATP	PA-O3A-PB	-2.18	125.35	132.83
2	B	301[A]	ATP	C2-N1-C6	2.07	122.29	118.75
2	B	301[B]	ATP	PB-O3B-PG	-2.06	125.74	132.83
2	A	301[A]	ATP	O2A-PA-O1A	2.03	122.30	112.24

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301[A]	ATP	PB-O3B-PG-O2G
2	B	301[A]	ATP	C5'-O5'-PA-O1A
2	B	301[A]	ATP	C5'-O5'-PA-O2A
2	B	301[B]	ATP	PB-O3B-PG-O3G
2	A	301[A]	ATP	O4'-C4'-C5'-O5'
2	A	301[A]	ATP	C3'-C4'-C5'-O5'
2	B	301[A]	ATP	O4'-C4'-C5'-O5'
2	B	301[A]	ATP	C3'-C4'-C5'-O5'
2	B	301[B]	ATP	O4'-C4'-C5'-O5'
2	B	301[B]	ATP	C3'-C4'-C5'-O5'
4	A	303	EDO	O1-C1-C2-O2
2	A	301[A]	ATP	PB-O3A-PA-O1A
2	B	301[B]	ATP	C4'-C5'-O5'-PA
2	A	301[A]	ATP	PB-O3A-PA-O2A
2	B	301[A]	ATP	C4'-C5'-O5'-PA
2	A	301[A]	ATP	PA-O3A-PB-O1B
2	B	301[A]	ATP	PA-O3A-PB-O1B

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
2	A	301[A]	ATP	PB-O3B-PG-O3G
2	B	301[B]	ATP	PB-O3B-PG-O2G
2	B	301[A]	ATP	C5'-O5'-PA-O3A
2	A	301[A]	ATP	PA-O3A-PB-O2B
2	B	301[A]	ATP	PA-O3A-PB-O2B
2	A	301[A]	ATP	PB-O3B-PG-O1G


There are no ring outliers.


1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301[A]	ATP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	162/177 (91%)	0.19	9 (5%) 24 26	11, 22, 45, 64	0
1	B	164/177 (92%)	0.23	6 (3%) 41 46	15, 26, 52, 69	0
All	All	326/354 (92%)	0.21	15 (4%) 32 35	11, 24, 51, 69	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	243	SER	5.6
1	A	270	ALA	3.3
1	A	181	ARG	3.3
1	B	270	ALA	3.2
1	A	242	ARG	3.2
1	B	208	ASP	3.0
1	B	244	ASP	2.9
1	A	192	ILE	2.8
1	B	184	ALA	2.7
1	A	271	SER	2.6
1	A	102	ALA	2.5
1	B	103	MET	2.3
1	A	269	LEU	2.2
1	B	185	ASN	2.1
1	A	156	GLY	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	EDO	A	303	4/4	0.66	0.22	52,63,66,66	0
2	ATP	B	301[A]	31/31	0.95	0.15	16,47,67,69	43
2	ATP	B	301[B]	31/31	0.95	0.15	16,49,67,69	43
2	ATP	A	301[A]	31/31	0.97	0.10	18,63,85,94	0
3	CA	B	302	1/1	0.99	0.07	18,18,18,18	0
3	CA	A	302	1/1	0.99	0.08	19,19,19,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




### Electron density around ATP B 301 (A):

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
 mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
 and green (positive)






**Electron density around ATP B 301 (B):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around ATP A 301 (A):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers

There are no such residues in this entry.