



# Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 12:23 pm BST

PDB ID : 6D2W  
Title : Crystal structure of Prevotella bryantii endo-beta-mannanase/endo-beta-glucanase PbGH26A-GH5A  
Authors : Stogios, P.J.; Skarina, T.; McGregor, N.; Nocek, B.; Di Leo, R.; Brumer, H.; Savchenko, A.  
Deposited on : 2018-04-14  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	776	<div> <div>2%</div> <div> <div></div> <div>83%</div> <div>6%</div> <div>11%</div> </div> </div>
1	B	776	<div> <div>2%</div> <div> <div></div> <div>82%</div> <div>6%</div> <div>12%</div> </div> </div>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 12730 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aryl-phospho-beta-D-glucosidase BglC, GH1 family.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	687	Total	C	N	O	S	0	3	0
			5454	3464	907	1056	27			
1	B	685	Total	C	N	O	S	0	1	0
			5433	3450	904	1052	27			

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: C<sub>8</sub>H<sub>18</sub>N<sub>2</sub>O<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
3	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
3	B	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	936	Total	O	0	28
			964	964		
4	B	801	Total	O	0	15
			816	816		

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A:
- 
- 83% 2% 6% 11%
- MET LYS LYS PHE ILE PHE SER LEU LEU THR ALA LEU THR ILE PHE SER VAL PHE ALA GLN ALA SER ASP THR TYR VAL VAL ASN LYS ALA ASP GLN SER TYR LYS MET GLU PRO ASN ILE LYS PHE ASN GLY ASP GLY THR PHE GLY TYR MET GLY

- Chain B:
- 
- 82% 6% 12%
- PRE K230 F231 E234 E237 K245 K255 F258 A271 N272 N273 D283 Y288 L343 C372 N376 P398 W399 S403 I425 F443 F447 D454 W490 P495 K517 N533 A539 K547 V552 Q597 K637 T686
- PRE LYS LYS PHE GLY GLN ASN VAL TRP ASN ILE SER ASP VAL LYS SER VAL THR PHE ALA GLN SER ASP THR TYR VAL ASN LYS ASP GLY THR SER GLN SER TYR LYS MET MET GLU PHE PRO ASN ILE LYS PHE ASN GLY ASP GLY THR PHE GLY TYR MET THR GLY

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	54.48 Å 96.75 Å 138.09 Å 90.00° 99.27° 90.00°	Depositor
Resolution (Å)	19.81 – 2.10 19.81 – 2.10	Depositor EDS
% Data completeness (in resolution range)	93.7 (19.81-2.10) 93.7 (19.81-2.10)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.16 (at 2.09 Å)	Xtriage
Refinement program	PHENIX (dev_3026: ???)	Depositor
R, $R_{free}$	0.146 , 0.193 0.146 , 0.193	Depositor DCC
$R_{free}$ test set	2013 reflections (2.60%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.0	Xtriage
Anisotropy	0.026	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 62.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.022 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	12730	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.92 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.8316e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, EPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.30	0/5618	0.47	0/7661
1	B	0.29	0/5591	0.46	0/7623
All	All	0.30	0/11209	0.47	0/15284

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5454	0	5089	27	0
1	B	5433	0	5062	26	0
2	A	12	0	16	0	0
2	B	6	0	8	0	0
3	A	30	0	34	1	0
3	B	15	0	17	0	0
4	A	964	0	0	8	0
4	B	816	0	0	2	0
All	All	12730	0	10226	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:642:LEU:O	1:A:693:LYS:NZ	2.29	0.59
1:B:547:LYS:HB2	1:B:552:VAL:HG11	1.85	0.59
1:B:227:TRP:HA	1:B:230:LYS:HE3	1.86	0.58
1:A:672:LYS:HD2	1:A:679:CYS:SG	2.45	0.56
1:A:290:GLN:NE2	4:A:916:HOH:O	2.39	0.55
1:A:133:GLU:O	1:A:137:ILE:HG12	2.06	0.55
1:B:258:PHE:HB3	1:B:288:TYR:HE1	1.70	0.55
1:B:754:SER:O	1:B:757:ILE:HG22	2.07	0.55
1:B:343:LEU:O	1:B:372:CYS:HA	2.08	0.53
1:A:343:LEU:O	1:A:372:CYS:HA	2.10	0.52
1:A:380:THR:HG22	4:A:1505:HOH:O	2.10	0.51
1:A:605:LEU:C	1:A:605:LEU:HD23	2.32	0.50
1:B:197:ASN:OD1	4:B:901:HOH:O	2.20	0.50
1:A:710:SER:HB3	1:A:712:ILE:HD12	1.93	0.49
1:A:197:ASN:ND2	4:A:926:HOH:O	2.45	0.49
1:B:196:GLU:HG3	1:B:227:TRP:CG	2.47	0.49
1:B:234:GLU:HA	1:B:237:GLU:HB3	1.94	0.49
1:B:245:LYS:NZ	4:B:922:HOH:O	2.47	0.48
1:B:153:ILE:HB	1:B:189:HIS:HB2	1.97	0.47
1:A:194:GLN:NE2	4:A:939:HOH:O	2.48	0.46
1:B:193:PRO:HD3	1:B:231:PHE:CZ	2.51	0.46
1:A:707:HIS:HB3	1:A:711:ASP:OD2	2.16	0.46
1:B:219:SER:OG	1:B:283:ASP:OD2	2.32	0.46
1:A:499:TYR:CE2	1:A:500:GLU:HG3	2.49	0.46
3:A:804:EPE:H82	3:A:804:EPE:H31	1.55	0.46
1:B:186:LEU:HB2	1:B:255:TRP:HE3	1.81	0.46
1:A:453:LEU:HD11	1:A:519:ILE:HD12	1.98	0.46
1:A:659:GLN:NE2	1:A:702:GLY:HA3	2.31	0.45
1:A:733:MET:HG3	1:A:744:THR:HG21	1.97	0.45
1:A:398:PRO:HG2	1:A:410:TRP:CZ2	2.52	0.44
1:B:637:LYS:HE3	1:B:686:ILE:HD13	2.00	0.44
1:A:217:LYS:HB3	1:A:217:LYS:HE2	1.68	0.44
1:A:147:ILE:HG12	1:A:183:ILE:HB	1.99	0.44
1:A:509:ASP:OD2	4:A:901:HOH:O	2.21	0.44
1:B:517:LYS:HE2	1:B:517:LYS:HB3	1.82	0.43
1:B:227:TRP:HA	1:B:230:LYS:HG2	2.00	0.43
1:B:146:ALA:HB1	1:B:425:ILE:HD11	2.01	0.43
1:B:495:PRO:HA	1:B:533:ASN:OD1	2.19	0.43
1:B:761:TRP:CD2	1:B:767:VAL:HG21	2.53	0.43
1:A:258:PHE:HB3	1:A:288:TYR:HE1	1.83	0.43

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:447:PHE:CZ	1:B:751:PHE:HB3	2.54	0.43
1:A:99:SER:OG	4:A:902:HOH:O	2.22	0.42
1:B:372:CYS:O	1:B:398:PRO:HA	2.18	0.42
1:B:399:TRP:HB3	1:B:403:SER:HB2	2.01	0.42
1:A:90:VAL:N	4:A:948:HOH:O	2.52	0.42
1:B:704:TYR:CE1	1:B:744:THR:HB	2.54	0.42
1:B:736:LEU:O	1:B:740:HIS:HD2	2.03	0.42
1:A:495:PRO:HA	1:A:533:ASN:OD1	2.20	0.41
1:A:90:VAL:HG11	4:A:1724:HOH:O	2.20	0.41
1:A:547:LYS:HB2	1:A:552:VAL:HG11	2.01	0.41
1:B:454:ASP:HB3	1:B:495:PRO:HB2	2.03	0.41
1:B:443:PHE:O	1:B:490:ASN:HB2	2.20	0.41
1:A:412:LYS:HE2	1:A:412:LYS:HB3	1.85	0.40
1:A:447:PHE:CZ	1:A:751:PHE:HB3	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	688/776 (89%)	665 (97%)	22 (3%)	1 (0%)	51	54
1	B	684/776 (88%)	657 (96%)	25 (4%)	2 (0%)	41	41
All	All	1372/1552 (88%)	1322 (96%)	47 (3%)	3 (0%)	47	49

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	539	ALA
1	A	539	ALA
1	B	127	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	568/643 (88%)	567 (100%)	1 (0%)	93	96
1	B	564/643 (88%)	562 (100%)	2 (0%)	91	94
All	All	1132/1286 (88%)	1129 (100%)	3 (0%)	92	95

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	190	PHE
1	B	597	GLN
1	B	692	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	659	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	GOL	A	801	-	5,5,5	0.96	0	5,5,5	1.03	0
3	EPE	A	804	-	15,15,15	0.77	1 (6%)	18,20,20	1.77	4 (22%)
3	EPE	B	802	-	15,15,15	0.77	1 (6%)	18,20,20	1.84	4 (22%)
2	GOL	B	801	-	5,5,5	0.86	0	5,5,5	0.99	0
3	EPE	A	803	-	15,15,15	0.78	1 (6%)	18,20,20	1.82	5 (27%)
2	GOL	A	802	-	5,5,5	0.87	0	5,5,5	1.01	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	801	-	-	2/4/4/4	-
3	EPE	A	804	-	-	2/9/19/19	0/1/1/1
3	EPE	B	802	-	-	2/9/19/19	0/1/1/1
2	GOL	B	801	-	-	4/4/4/4	-
3	EPE	A	803	-	-	6/9/19/19	0/1/1/1
2	GOL	A	802	-	-	0/4/4/4	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	803	EPE	C10-S	2.48	1.81	1.77
3	A	804	EPE	C10-S	2.46	1.81	1.77
3	B	802	EPE	C10-S	2.42	1.80	1.77

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	804	EPE	C5-N4-C3	4.38	118.68	108.83
3	B	802	EPE	C5-N4-C3	3.92	117.66	108.83
3	A	803	EPE	C5-N4-C3	3.70	117.17	108.83
3	B	802	EPE	C7-N4-C3	3.46	120.09	111.23
3	B	802	EPE	C7-N4-C5	3.41	119.97	111.23
3	A	803	EPE	C7-N4-C3	3.29	119.65	111.23
3	A	804	EPE	C7-N4-C5	3.29	119.64	111.23
3	A	803	EPE	C7-N4-C5	3.19	119.39	111.23
3	B	802	EPE	O1S-S-C10	2.89	110.39	106.92
3	A	804	EPE	O2S-S-C10	2.50	109.92	106.92
3	A	803	EPE	O1S-S-C10	2.49	109.91	106.92
3	A	803	EPE	C6-N1-C2	2.43	114.29	108.83
3	A	804	EPE	C7-N4-C3	2.16	116.75	111.23

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	801	GOL	C1-C2-C3-O3
3	A	804	EPE	C8-C7-N4-C3
3	B	802	EPE	C8-C7-N4-C3
2	B	801	GOL	O1-C1-C2-O2
2	B	801	GOL	O1-C1-C2-C3
2	B	801	GOL	C1-C2-C3-O3
2	B	801	GOL	O2-C2-C3-O3
3	A	803	EPE	C8-C7-N4-C3
3	A	803	EPE	C9-C10-S-O1S
3	A	803	EPE	C9-C10-S-O2S
2	A	801	GOL	O2-C2-C3-O3
3	A	803	EPE	C9-C10-S-O3S
3	A	803	EPE	N4-C7-C8-O8
3	B	802	EPE	N4-C7-C8-O8
3	A	804	EPE	N4-C7-C8-O8
3	A	803	EPE	C10-C9-N1-C6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	804	EPE	1	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	687/776 (88%)	-0.56	13 (1%) 66 71	14, 26, 47, 91	0
1	B	685/776 (88%)	-0.37	13 (1%) 66 71	16, 30, 58, 103	0
All	All	1372/1552 (88%)	-0.46	26 (1%) 66 71	14, 27, 56, 103	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	94	PHE	6.9
1	B	92	GLY	6.6
1	B	93	VAL	6.4
1	B	272	ASN	5.0
1	A	91	SER	4.9
1	A	376	ASN	3.9
1	B	271	ALA	3.6
1	A	90	VAL	3.5
1	B	197	ASN	3.4
1	A	711	ASP	3.1
1	A	377	SER	2.9
1	A	162	SER	2.9
1	A	94	PHE	2.7
1	A	696	THR	2.4
1	B	222	LEU	2.4
1	B	162	SER	2.3
1	A	272	ASN	2.3
1	B	273	TRP	2.3
1	B	376	ASN	2.2
1	A	161	GLY	2.2
1	A	776	ASN	2.2
1	B	161	GLY	2.1
1	B	224	SER	2.1
1	B	711	ASP	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	92	GLY	2.1
1	A	93	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	EPE	A	804	15/15	0.64	0.28	85,93,117,118	0
2	GOL	B	801	6/6	0.72	0.17	70,73,73,75	0
2	GOL	A	801	6/6	0.77	0.26	83,84,86,87	0
2	GOL	A	802	6/6	0.78	0.15	68,70,70,71	0
3	EPE	B	802	15/15	0.84	0.19	79,82,108,108	0
3	EPE	A	803	15/15	0.90	0.15	73,75,90,91	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.