



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 18, 2021 – 06:09 PM JST

PDB ID : 7D2T  
Title : Crystal structure of Rsu1/PINCH1\_LIM45C complex  
Authors : Yang, H.; Wei, Z.; Yu, C.  
Deposited on : 2020-09-17  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.16  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.16


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	282	<div> <div>5%</div> <div>85%</div> <div>12%</div> </div>
1	C	282	<div> <div>5%</div> <div>85%</div> <div>12%</div> </div>
2	B	142	<div> <div>18%</div> <div>87%</div> <div>5% 8%</div> </div>
2	D	142	<div> <div>6%</div> <div>89%</div> <div>9%</div> </div>

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6487 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ras suppressor protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	247	Total	C	N	O	S	2	1	0
			1992	1273	337	377	5			
1	C	247	Total	C	N	O	S	0	2	0
			1979	1265	335	374	5			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	expression tag	UNP Q15404
A	-3	ALA	-	expression tag	UNP Q15404
A	-2	MET	-	expression tag	UNP Q15404
A	-1	GLY	-	expression tag	UNP Q15404
A	0	SER	-	expression tag	UNP Q15404
C	-4	GLY	-	expression tag	UNP Q15404
C	-3	ALA	-	expression tag	UNP Q15404
C	-2	MET	-	expression tag	UNP Q15404
C	-1	GLY	-	expression tag	UNP Q15404
C	0	SER	-	expression tag	UNP Q15404

- Molecule 2 is a protein called LIM and senescent cell antigen-like-containing domain protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	130	Total	C	N	O	S	0	0	0
			1048	670	190	173	15			
2	D	129	Total	C	N	O	S	1	1	0
			1036	661	187	173	15			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	184	GLY	-	expression tag	UNP P48059

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	185	PRO	-	expression tag	UNP P48059
B	186	GLY	-	expression tag	UNP P48059
B	187	SER	-	expression tag	UNP P48059
D	184	GLY	-	expression tag	UNP P48059
D	185	PRO	-	expression tag	UNP P48059
D	186	GLY	-	expression tag	UNP P48059
D	187	SER	-	expression tag	UNP P48059

- Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula:  $C_4H_{12}NO_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			8	4	1	3		
3	C	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	4	Total	Zn	0	0
			4	4		
5	D	4	Total	Zn	0	0
			4	4		

- Molecule 6 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			10	6	4		
6	D	1	Total	C	O	0	0
			10	6	4		



- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	146	Total	O	0	0
			146	146		
7	B	42	Total	O	0	0
			42	42		
7	C	133	Total	O	0	0
			133	133		
7	D	37	Total	O	0	0
			37	37		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Ras suppressor protein 1




- Molecule 1: Ras suppressor protein 1



- Molecule 2: LIM and senescent cell antigen-like-containing domain protein 1



- Molecule 2: LIM and senescent cell antigen-like-containing domain protein 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	114.62Å 51.29Å 119.63Å 90.00° 101.55° 90.00°	Depositor
Resolution (Å)	50.00 – 2.20 48.15 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.7 (50.00-2.20) 93.7 (48.15-2.20)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.38 (at 2.20Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.170 , 0.198 0.175 , 0.199	Depositor DCC
$R_{free}$ test set	2009 reflections (2.86%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	37.3	Xtriage
Anisotropy	0.220	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 42.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.115 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6487	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.85% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, PGE, TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.23	0/2033	0.41	0/2758
1	C	0.24	0/2023	0.40	0/2748
2	B	0.23	0/1074	0.38	0/1441
2	D	0.23	0/1065	0.38	0/1432
All	All	0.23	0/6195	0.40	0/8379

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1992	0	2023	4	0
1	C	1979	0	1998	3	0
2	B	1048	0	1045	5	0
2	D	1036	0	1019	1	0
3	A	8	0	12	0	0
3	C	8	0	12	0	0
4	A	18	0	24	1	0
4	B	6	0	8	0	0
4	C	6	0	8	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	4	0	0	0	0
5	D	4	0	0	0	0
6	B	10	0	14	1	0
6	D	10	0	14	1	0
7	A	146	0	0	0	0
7	B	42	0	0	0	0
7	C	133	0	0	0	0
7	D	37	0	0	0	0
All	All	6487	0	6177	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:316:LYS:HA	6:B:406:PGE:H3	1.82	0.62
1:A:120[A]:ASN:ND2	4:A:302:GOL:O2	2.26	0.58
2:B:222:CYS:HB3	2:B:243:HIS:HE1	1.74	0.52
1:C:165:ARG:HG3	1:C:188:GLN:HG2	1.98	0.46
2:D:316:LYS:HA	6:D:405:PGE:H42	1.99	0.43
2:B:281:CYS:HB3	2:B:303:CYS:SG	2.59	0.43
1:A:6:LYS:HG2	1:A:34:ASN:HB3	2.00	0.42
1:A:117:THR:HG23	1:A:142:SER:HB2	2.00	0.42
2:B:222:CYS:HB3	2:B:243:HIS:CE1	2.53	0.42
1:A:221:PRO:O	1:A:225:GLN:HG2	2.20	0.42
1:C:142:SER:OG	1:C:163:SER:OG	2.26	0.41
2:B:191:PRO:HB3	2:B:211:GLN:HG2	2.02	0.41
1:C:93:LEU:HB2	1:C:116:LEU:HD23	2.01	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	246/282 (87%)	235 (96%)	11 (4%)	0	100	100
1	C	247/282 (88%)	238 (96%)	9 (4%)	0	100	100
2	B	128/142 (90%)	126 (98%)	2 (2%)	0	100	100
2	D	128/142 (90%)	125 (98%)	3 (2%)	0	100	100
All	All	749/848 (88%)	724 (97%)	25 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/258 (89%)	229 (100%)	1 (0%)	91	96
1	C	227/258 (88%)	226 (100%)	1 (0%)	91	96
2	B	115/123 (94%)	115 (100%)	0	100	100
2	D	113/123 (92%)	112 (99%)	1 (1%)	78	88
All	All	685/762 (90%)	682 (100%)	3 (0%)	91	96

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	61	GLU
1	C	61	GLU
2	D	201	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 17 ligands modelled in this entry, 8 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	GOL	A	304	-	5,5,5	0.38	0	5,5,5	0.39	0
4	GOL	A	302	-	5,5,5	0.38	0	5,5,5	0.25	0
4	GOL	C	302	-	5,5,5	0.36	0	5,5,5	0.28	0
6	PGE	D	405	-	9,9,9	0.30	0	8,8,8	0.36	0
4	GOL	A	303	-	5,5,5	0.37	0	5,5,5	0.27	0
3	TRS	A	301	-	7,7,7	0.32	0	9,9,9	0.31	0
6	PGE	B	406	-	9,9,9	0.30	0	8,8,8	0.31	0
3	TRS	C	301	-	7,7,7	0.31	0	9,9,9	0.35	0
4	GOL	B	405	-	5,5,5	0.37	0	5,5,5	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	304	-	-	2/4/4/4	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	302	-	-	4/4/4/4	-
4	GOL	C	302	-	-	2/4/4/4	-
6	PGE	D	405	-	-	3/7/7/7	-
4	GOL	A	303	-	-	2/4/4/4	-
3	TRS	A	301	-	-	3/9/9/9	-
6	PGE	B	406	-	-	3/7/7/7	-
3	TRS	C	301	-	-	3/9/9/9	-
4	GOL	B	405	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	304	GOL	O1-C1-C2-C3
4	C	302	GOL	O1-C1-C2-O2
3	C	301	TRS	C2-C-C1-O1
3	C	301	TRS	C3-C-C1-O1
4	A	302	GOL	O1-C1-C2-C3
4	C	302	GOL	O1-C1-C2-C3
4	A	303	GOL	O1-C1-C2-C3
4	B	405	GOL	O1-C1-C2-C3
6	D	405	PGE	O3-C5-C6-O4
6	B	406	PGE	O1-C1-C2-O2
4	A	304	GOL	O1-C1-C2-O2
4	A	303	GOL	O1-C1-C2-O2
4	A	302	GOL	O2-C2-C3-O3
4	B	405	GOL	O1-C1-C2-O2
6	B	406	PGE	C4-C3-O2-C2
3	A	301	TRS	C2-C-C3-O3
3	A	301	TRS	N-C-C3-O3
3	C	301	TRS	N-C-C1-O1
4	A	302	GOL	O1-C1-C2-O2
3	A	301	TRS	C1-C-C3-O3
6	B	406	PGE	O2-C3-C4-O3
6	D	405	PGE	O2-C3-C4-O3
4	A	302	GOL	C1-C2-C3-O3
6	D	405	PGE	C3-C4-O3-C5

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	302	GOL	1	0
6	D	405	PGE	1	0
6	B	406	PGE	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	247/282 (87%)	0.25	14 (5%) 23 22	25, 39, 76, 137	6 (2%)
1	C	247/282 (87%)	0.28	13 (5%) 26 25	26, 42, 86, 147	3 (1%)
2	B	130/142 (91%)	0.77	26 (20%) 1 1	29, 62, 106, 124	1 (0%)
2	D	129/142 (90%)	0.49	9 (6%) 16 15	32, 64, 90, 98	2 (1%)
All	All	753/848 (88%)	0.39	62 (8%) 11 10	25, 46, 101, 147	12 (1%)

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	13	ARG	7.4
1	C	8	LEU	7.1
1	C	10	GLU	6.7
1	C	9	VAL	6.4
1	A	8	LEU	6.3
1	A	6	LYS	6.3
1	A	13	ARG	6.3
1	C	7	LYS	5.9
1	A	7	LYS	5.4
1	C	5	LEU	5.3
1	A	11	GLU	5.1
1	A	14	GLU	4.9
1	A	10	GLU	4.9
1	C	6	LYS	4.7
1	A	5	LEU	4.6
1	A	9	VAL	4.2
2	B	198	ARG	4.2
2	B	199	PRO	4.0
2	B	192	ILE	4.0
2	B	191	PRO	3.9
1	C	11	GLU	3.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	D	189	GLY	3.9
2	B	201	GLU	3.8
1	A	15	LYS	3.7
2	D	197	ARG	3.6
2	B	189	GLY	3.5
1	C	14	GLU	3.4
2	B	190	VAL	3.4
2	B	211	GLN	3.3
2	B	197	ARG	3.3
2	B	205	VAL	3.2
2	B	202	GLY	3.2
1	A	12	SER	3.1
2	D	242	THR	3.1
1	C	12	SER	3.0
1	A	16	ASN	3.0
2	B	203	ARG	3.0
2	B	318	LEU	2.9
2	D	200	ILE	2.8
1	C	15	LYS	2.8
1	A	251	GLN	2.7
2	B	214	VAL	2.7
2	B	208	MET	2.5
2	B	200	ILE	2.5
1	C	16	ASN	2.4
2	B	207	ALA	2.4
2	B	236	GLY	2.4
2	B	195	ALA	2.4
2	D	201	GLU	2.4
2	B	209	GLY	2.3
2	B	217	PHE	2.3
2	D	202	GLY	2.3
2	D	190	VAL	2.3
2	B	215	GLU	2.2
2	B	213	HIS	2.1
2	D	209	GLY	2.1
1	A	249	HIS	2.1
2	B	196	CYS	2.1
2	D	192	ILE	2.0
2	B	212	TRP	2.0
1	C	244	TYR	2.0
2	B	228	GLY	2.0



## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	PGE	D	405	10/10	0.76	0.21	78,84,90,91	0
4	GOL	A	304	6/6	0.77	0.37	105,105,107,108	0
6	PGE	B	406	10/10	0.80	0.22	75,85,97,99	0
4	GOL	A	303	6/6	0.81	0.34	103,105,106,107	0
4	GOL	C	302	6/6	0.82	0.23	57,64,69,73	0
4	GOL	B	405	6/6	0.86	0.40	88,91,92,93	0
3	TRS	C	301	8/8	0.88	0.16	47,52,58,60	0
4	GOL	A	302	6/6	0.89	0.31	77,83,84,85	0
3	TRS	A	301	8/8	0.90	0.14	44,51,55,56	0
5	ZN	B	404	1/1	0.92	0.06	95,95,95,95	0
5	ZN	D	404	1/1	0.93	0.05	75,75,75,75	0
5	ZN	B	403	1/1	0.96	0.09	68,68,68,68	0
5	ZN	D	402	1/1	0.98	0.09	64,64,64,64	0
5	ZN	B	402	1/1	0.99	0.14	49,49,49,49	0
5	ZN	D	401	1/1	1.00	0.12	39,39,39,39	0
5	ZN	D	403	1/1	1.00	0.10	51,51,51,51	0
5	ZN	B	401	1/1	1.00	0.14	41,41,41,41	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.