



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 12:29 pm BST

PDB ID : 5D6R
Title : Acetolactate Synthase from *Klebsiella pneumoniae* in Complex with Mechanism-Based Inhibitor
Authors : Latta, A.J.; Andrews, F.H.; McLeish, M.J.
Deposited on : 2015-08-12
Resolution : 2.28 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.28 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6980 (2.30-2.26)
Clashscore	141614	7711 (2.30-2.26)
Ramachandran outliers	138981	7597 (2.30-2.26)
Sidechain outliers	138945	7598 (2.30-2.26)
RSRZ outliers	127900	6849 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	579	<div> <div>3%</div> <div>92%</div> <div>5%</div> </div>
1	M	579	<div> <div>3%</div> <div>91%</div> <div>5%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PO4	M	603	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8538 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetolactate synthase, catabolic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	548	Total	C	N	O	S	0	0	0
			4090	2584	721	767	18			
1	M	553	Total	C	N	O	S	0	0	0
			4098	2594	714	772	18			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-19	MET	-	initiating methionine	UNP P27696
B	-18	GLY	-	expression tag	UNP P27696
B	-17	SER	-	expression tag	UNP P27696
B	-16	SER	-	expression tag	UNP P27696
B	-15	HIS	-	expression tag	UNP P27696
B	-14	HIS	-	expression tag	UNP P27696
B	-13	HIS	-	expression tag	UNP P27696
B	-12	HIS	-	expression tag	UNP P27696
B	-11	HIS	-	expression tag	UNP P27696
B	-10	HIS	-	expression tag	UNP P27696
B	-9	SER	-	expression tag	UNP P27696
B	-8	SER	-	expression tag	UNP P27696
B	-7	GLY	-	expression tag	UNP P27696
B	-6	LEU	-	expression tag	UNP P27696
B	-5	VAL	-	expression tag	UNP P27696
B	-4	PRO	-	expression tag	UNP P27696
B	-3	ARG	-	expression tag	UNP P27696
B	-2	GLY	-	expression tag	UNP P27696
B	-1	SER	-	expression tag	UNP P27696
B	0	HIS	-	expression tag	UNP P27696
B	407	SER	THR	engineered mutation	UNP P27696
M	-19	MET	-	initiating methionine	UNP P27696
M	-18	GLY	-	expression tag	UNP P27696
M	-17	SER	-	expression tag	UNP P27696
M	-16	SER	-	expression tag	UNP P27696

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
M	-15	HIS	-	expression tag	UNP P27696
M	-14	HIS	-	expression tag	UNP P27696
M	-13	HIS	-	expression tag	UNP P27696
M	-12	HIS	-	expression tag	UNP P27696
M	-11	HIS	-	expression tag	UNP P27696
M	-10	HIS	-	expression tag	UNP P27696
M	-9	SER	-	expression tag	UNP P27696
M	-8	SER	-	expression tag	UNP P27696
M	-7	GLY	-	expression tag	UNP P27696
M	-6	LEU	-	expression tag	UNP P27696
M	-5	VAL	-	expression tag	UNP P27696
M	-4	PRO	-	expression tag	UNP P27696
M	-3	ARG	-	expression tag	UNP P27696
M	-2	GLY	-	expression tag	UNP P27696
M	-1	SER	-	expression tag	UNP P27696
M	0	HIS	-	expression tag	UNP P27696
M	407	SER	THR	engineered mutation	UNP P27696

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	3	Total Mg 3 3	0	0
2	M	2	Total Mg 2 2	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total O P 5 4 1	0	0
3	M	1	Total O P 5 4 1	0	0

- Molecule 4 is 3-[(4-amino-2-methylpyrimidin-5-yl)methyl]-2-[(Z)-2-fluoro-1-hydroxy-2-phosphonoethenyl]-5-(2-[(S)-hydroxy(phosphonooxy)phosphoryl]oxy)ethyl)-4-methyl-1,3-thiazol-3-ium (three-letter code: EN0) (formula: C₁₄H₂₁FN₄O₁₁P₃S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C F N O P S 34 14 1 4 11 3 1	0	0
4	M	1	Total C F N O P S 34 14 1 4 11 3 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	121	Total O 121 121	0	0
5	M	146	Total O 146 146	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Acetolactate synthase, catabolic



- Molecule 1: Acetolactate synthase, catabolic



4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	85.12Å 133.35Å 110.62Å 90.00° 95.40° 90.00°	Depositor
Resolution (Å)	55.07 – 2.28 55.07 – 2.28	Depositor EDS
% Data completeness (in resolution range)	99.8 (55.07-2.28) 93.8 (55.07-2.28)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.07 (at 2.27Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.201 , 0.236 0.208 , 0.237	Depositor DCC
R_{free} test set	2000 reflections (3.56%)	wwPDB-VP
Wilson B-factor (Å ²)	33.4	Xtriage
Anisotropy	0.471	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 45.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8538	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EN0, PO4, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.22	0/4169	0.40	0/5676
1	M	0.22	0/4181	0.39	0/5701
All	All	0.22	0/8350	0.39	0/11377

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	4090	0	4032	8	0
1	M	4098	0	4012	14	0
2	B	3	0	0	0	0
2	M	2	0	0	0	0
3	B	5	0	0	0	0
3	M	5	0	0	2	0
4	B	34	0	0	1	0
4	M	34	0	0	1	0
5	B	121	0	0	1	0
5	M	146	0	0	4	0
All	All	8538	0	8044	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:366:LEU:O	1:B:375:ARG:NH1	1.94	1.01
1:B:555:LEU:O	1:B:556:SER:OG	1.94	0.84
1:B:53:PRO:O	1:B:462:ARG:NH2	2.28	0.66
1:M:-3:ARG:NH1	1:M:182:PRO:O	2.30	0.64
1:M:228:ARG:NH2	5:M:703:HOH:O	2.31	0.63
1:B:365:GLN:NE2	5:B:704:HOH:O	2.32	0.62
1:M:365:GLN:OE1	1:M:365:GLN:N	2.39	0.56
1:B:153:ARG:NH1	1:B:184:SER:OG	2.38	0.55
1:M:114:LYS:NZ	5:M:709:HOH:O	2.40	0.54
1:M:440:LYS:NZ	1:M:466:ASN:OD1	2.41	0.54
1:B:458:GLU:OE1	1:B:462:ARG:NH1	2.45	0.49
1:M:27:ARG:NH1	5:M:718:HOH:O	2.48	0.47
1:M:366:LEU:O	1:M:375:ARG:NH1	2.48	0.46
1:M:264:ASN:ND2	1:M:553:LEU:O	2.48	0.46
1:B:420:GLN:O	4:B:605:EN0:N33	2.48	0.46
1:M:426:LEU:HB3	1:M:427:PRO:HD3	1.98	0.45
1:M:8:ARG:NH1	5:M:719:HOH:O	2.48	0.44
1:B:426:LEU:HB3	1:B:427:PRO:HD3	1.99	0.43
1:M:266:GLN:NE2	3:M:603:PO4:O1	2.52	0.42
1:M:352:ARG:NH1	3:M:603:PO4:O1	2.45	0.42
1:M:522:GLU:HB2	1:M:523:PRO:HD3	2.01	0.42
1:M:82:GLY:HA3	1:M:120:GLN:HG3	2.02	0.41
4:M:604:EN0:C08	4:M:604:EN0:N33	2.83	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	544/579 (94%)	527 (97%)	17 (3%)	0	100	100
1	M	549/579 (95%)	535 (97%)	14 (3%)	0	100	100
All	All	1093/1158 (94%)	1062 (97%)	31 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	418/462 (90%)	414 (99%)	4 (1%)	76	86
1	M	417/462 (90%)	413 (99%)	4 (1%)	76	86
All	All	835/924 (90%)	827 (99%)	8 (1%)	76	86

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	312	ARG
1	B	366	LEU
1	B	367	ASN
1	B	455	MET
1	M	118	VAL
1	M	235	HIS
1	M	358	LEU
1	M	455	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 5 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EN0	M	604	2	27,35,35	1.43	4 (14%)	32,54,54	3.13	9 (28%)
4	EN0	B	605	2	27,35,35	1.41	4 (14%)	32,54,54	3.00	9 (28%)
3	PO4	B	604	-	4,4,4	0.94	0	6,6,6	0.41	0
3	PO4	M	603	-	4,4,4	0.93	0	6,6,6	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EN0	M	604	2	-	10/16/31/31	0/2/2/2
4	EN0	B	605	2	-	10/16/31/31	0/2/2/2

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	605	EN0	C32-N33	2.87	1.41	1.34
4	M	604	EN0	P13-O15	-2.83	1.47	1.54
4	M	604	EN0	C32-N33	2.81	1.41	1.34
4	B	605	EN0	P13-O15	-2.69	1.47	1.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	M	604	EN0	C06-C05	2.67	1.56	1.51
4	B	605	EN0	C06-C05	2.43	1.56	1.51
4	B	605	EN0	C19-C18	2.12	1.51	1.50
4	M	604	EN0	C19-C18	2.01	1.51	1.50

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	M	604	EN0	C30-N07-C08	15.28	117.14	108.64
4	B	605	EN0	C30-N07-C08	14.49	116.70	108.64
4	M	604	EN0	C05-C06-N07	-4.62	105.62	113.26
4	B	605	EN0	C05-C06-N07	-3.66	107.21	113.26
4	B	605	EN0	C18-C30-N07	3.16	114.26	107.66
4	M	604	EN0	C18-C30-N07	3.02	113.97	107.66
4	B	605	EN0	C04-N03-C02	2.81	120.74	115.96
4	M	604	EN0	C04-N03-C02	2.75	120.65	115.96
4	M	604	EN0	C19-C18-C30	2.51	129.45	127.43
4	M	604	EN0	P22-O24-P25	-2.50	124.25	132.83
4	B	605	EN0	N03-C02-N34	-2.48	121.28	125.54
4	B	605	EN0	P22-O24-P25	-2.45	124.42	132.83
4	M	604	EN0	N03-C02-N34	-2.42	121.38	125.54
4	B	605	EN0	C19-C18-C30	2.36	129.33	127.43
4	B	605	EN0	C02-N34-C32	2.31	121.68	118.08
4	M	604	EN0	C02-N34-C32	2.25	121.60	118.08
4	M	604	EN0	C01-C02-N03	2.20	119.56	117.14
4	B	605	EN0	C31-C30-C18	-2.16	122.88	127.60

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	605	EN0	C32-C05-C06-N07
4	B	605	EN0	C18-C19-C20-O21
4	B	605	EN0	C20-O21-P22-O24
4	B	605	EN0	P22-O24-P25-O28
4	M	604	EN0	C32-C05-C06-N07
4	M	604	EN0	C18-C19-C20-O21
4	M	604	EN0	C20-O21-P22-O24
4	B	605	EN0	C19-C20-O21-P22
4	B	605	EN0	C04-C05-C06-N07
4	B	605	EN0	C20-O21-P22-O23
4	B	605	EN0	C20-O21-P22-O29

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	M	604	EN0	C04-C05-C06-N07
4	M	604	EN0	C20-O21-P22-O23
4	M	604	EN0	C20-O21-P22-O29
4	M	604	EN0	C30-C18-C19-C20
4	B	605	EN0	P22-O24-P25-O26
4	B	605	EN0	P25-O24-P22-O29
4	M	604	EN0	P22-O24-P25-O26
4	M	604	EN0	P22-O24-P25-O27
4	M	604	EN0	C19-C20-O21-P22


There are no ring outliers.

3 monomers are involved in 4 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	M	604	EN0	1	0
4	B	605	EN0	1	0
3	M	603	PO4	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand EN0 M 604



Ligand EN0 B 605



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	B	548/579 (94%)	0.09	16 (2%)	51 57	27, 53, 85, 118	0
1	M	553/579 (95%)	-0.01	16 (2%)	51 57	28, 49, 86, 139	0
All	All	1101/1158 (95%)	0.04	32 (2%)	51 57	27, 51, 86, 139	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	M	200	VAL	5.1
1	M	358	LEU	3.8
1	M	249	ASN	3.7
1	B	118	VAL	3.3
1	B	45	LEU	3.2
1	M	250	GLN	3.1
1	B	516	GLU	3.1
1	M	3	LYS	2.9
1	M	196	ALA	2.9
1	M	354	HIS	2.9
1	B	178	GLY	2.8
1	B	365	GLN	2.7
1	M	203	LEU	2.7
1	B	370	ALA	2.7
1	M	5	TYR	2.6
1	M	526	ARG	2.5
1	B	517	SER	2.4
1	B	364	ALA	2.4
1	M	195	ASP	2.4
1	M	334	ASN	2.3
1	B	119	HIS	2.3
1	B	367	ASN	2.3
1	B	265	ASN	2.2
1	B	48	SER	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	120	GLN	2.2
1	M	1	MET	2.1
1	B	126	ALA	2.1
1	B	518	ALA	2.1
1	M	337	HIS	2.1
1	B	509	GLY	2.0
1	M	367	ASN	2.0
1	M	244	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MG	M	602	1/1	0.84	0.14	45,45,45,45	0
2	MG	M	601	1/1	0.92	0.10	34,34,34,34	0
2	MG	B	602	1/1	0.94	0.27	47,47,47,47	0
2	MG	B	603	1/1	0.94	0.09	55,55,55,55	0
4	EN0	B	605	34/34	0.95	0.15	34,45,104,105	2
2	MG	B	601	1/1	0.95	0.11	39,39,39,39	0
4	EN0	M	604	34/34	0.96	0.14	29,36,122,122	3
3	PO4	M	603	5/5	0.96	0.11	51,51,56,57	0
3	PO4	B	604	5/5	0.99	0.09	33,35,38,42	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around EN0 B 605:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around EN0 M 604:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.