



wwPDB X-ray Structure Validation Summary Report ⓘ

Jul 27, 2021 – 02:08 PM JST

PDB ID : 7D6C
Title : Crystal structure of CcmM N-terminal domain in complex with CcmN
Authors : Sun, H.; Cui, N.; Han, S.J.; Chen, Z.P.; Xia, L.Y.; Chen, Y.; Jiang, Y.L.;
Zhou, C.Z.
Deposited on : 2020-09-30
Resolution : 2.89 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.22
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.22


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	1	218	<div> <div>3%</div> <div> <div></div> <div>49%</div> <div>28%</div> <div>23%</div> </div> </div>
1	2	218	<div> <div>4%</div> <div> <div></div> <div>62%</div> <div>15%</div> <div>22%</div> </div> </div>
2	3	122	<div> <div>2%</div> <div> <div></div> <div>75%</div> <div>11%</div> <div>13%</div> </div> </div>
2	4	122	<div> <div>2%</div> <div> <div></div> <div>71%</div> <div>13%</div> <div>15%</div> </div> </div>
2	5	122	<div> <div>2%</div> <div> <div></div> <div>79%</div> <div>20%</div> <div>..</div> </div> </div>
2	F	122	<div> <div>3%</div> <div> <div></div> <div>74%</div> <div>25%</div> <div>..</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5696 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Carbon dioxide concentrating mechanism protein CcmM.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	1	167	Total	C	N	O	S	0	0	0
			1258	793	228	234	3			
1	2	169	Total	C	N	O	S	0	0	0
			1272	803	230	236	3			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
1	-8	MET	-	initiating methionine	UNP Q03513
1	-7	ASP	-	expression tag	UNP Q03513
1	-6	TYR	-	expression tag	UNP Q03513
1	-5	LYS	-	expression tag	UNP Q03513
1	-4	ASP	-	expression tag	UNP Q03513
1	-3	ASP	-	expression tag	UNP Q03513
1	-2	ASP	-	expression tag	UNP Q03513
1	-1	ASP	-	expression tag	UNP Q03513
1	0	LYS	-	expression tag	UNP Q03513
2	-8	MET	-	initiating methionine	UNP Q03513
2	-7	ASP	-	expression tag	UNP Q03513
2	-6	TYR	-	expression tag	UNP Q03513
2	-5	LYS	-	expression tag	UNP Q03513
2	-4	ASP	-	expression tag	UNP Q03513
2	-3	ASP	-	expression tag	UNP Q03513
2	-2	ASP	-	expression tag	UNP Q03513
2	-1	ASP	-	expression tag	UNP Q03513
2	0	LYS	-	expression tag	UNP Q03513

- Molecule 2 is a protein called Carboxysome assembly protein CcmN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	3	106	Total	C	N	O	S	0	0	0
			720	451	124	143	2			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	4	104	Total	C	N	O	S	0	0	0
			704	442	122	138	2			
2	5	121	Total	C	N	O	S	0	0	0
			836	527	143	163	3			
2	F	121	Total	C	N	O	S	0	0	0
			840	531	143	163	3			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
3	-3	PHE	-	linker	UNP P46204
3	-2	GLN	-	linker	UNP P46204
3	-1	SER	-	linker	UNP P46204
3	0	ASN	-	linker	UNP P46204
4	-3	PHE	-	linker	UNP P46204
4	-2	GLN	-	linker	UNP P46204
4	-1	SER	-	linker	UNP P46204
4	0	ASN	-	linker	UNP P46204
5	-3	PHE	-	linker	UNP P46204
5	-2	GLN	-	linker	UNP P46204
5	-1	SER	-	linker	UNP P46204
5	0	ASN	-	linker	UNP P46204
F	-3	PHE	-	linker	UNP P46204
F	-2	GLN	-	linker	UNP P46204
F	-1	SER	-	linker	UNP P46204
F	0	ASN	-	linker	UNP P46204


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	1	14	Total	O	0	0
			14	14		
3	2	9	Total	O	0	0
			9	9		
3	3	9	Total	O	0	0
			9	9		
3	4	10	Total	O	0	0
			10	10		
3	5	6	Total	O	0	0
			6	6		
3	F	18	Total	O	0	0
			18	18		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Carbon dioxide concentrating mechanism protein CcmM



- Molecule 1: Carbon dioxide concentrating mechanism protein CcmM



- Molecule 2: Carboxysome assembly protein CcmN




- Molecule 2: Carboxysome assembly protein CcmN






● Molecule 2: Carboxysome assembly protein CcmN



● Molecule 2: Carboxysome assembly protein CcmN



PRO

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	84.18Å 92.33Å 125.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.16 – 2.89 44.16 – 2.89	Depositor EDS
% Data completeness (in resolution range)	99.8 (44.16-2.89) 99.9 (44.16-2.89)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.25 (at 2.90Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.209 , 0.272 0.209 , 0.269	Depositor DCC
R_{free} test set	1105 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	63.6	Xtriage
Anisotropy	0.499	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 40.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5696	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	1	0.42	0/1284	0.62	0/1746
1	2	0.51	0/1300	0.92	4/1770 (0.2%)
2	3	0.49	0/724	0.75	1/986 (0.1%)
2	4	0.46	0/708	0.66	1/964 (0.1%)
2	5	0.66	2/847 (0.2%)	1.22	7/1159 (0.6%)
2	F	0.56	1/851 (0.1%)	0.77	1/1163 (0.1%)
All	All	0.52	3/5714 (0.1%)	0.85	14/7788 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	2	0	1
2	5	0	1
All	All	0	2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	5	13	ARG	CG-CD	10.04	1.77	1.51
2	5	13	ARG	CZ-NH1	5.63	1.40	1.33
2	F	88	CYS	CB-SG	-5.00	1.73	1.81

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	5	13	ARG	NE-CZ-NH2	21.68	131.14	120.30
1	2	164	ARG	NE-CZ-NH1	-19.36	110.62	120.30
2	5	13	ARG	NE-CZ-NH1	-18.82	110.89	120.30
2	5	13	ARG	CD-NE-CZ	12.44	141.01	123.60
1	2	164	ARG	NE-CZ-NH2	10.84	125.72	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	2	164	ARG	Sidechain
2	5	13	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1	1258	0	1243	41	0
1	2	1272	0	1257	21	1
2	3	720	0	751	9	0
2	4	704	0	738	13	0
2	5	836	0	870	14	2
2	F	840	0	872	22	2
3	1	14	0	0	1	0
3	2	9	0	0	0	0
3	3	9	0	0	0	0
3	4	10	0	0	1	0
3	5	6	0	0	0	0
3	F	18	0	0	2	0
All	All	5696	0	5731	110	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 110 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:5:13:ARG:CD	2:5:13:ARG:CG	1.77	1.60
1:1:135:MET:HE3	2:F:111:LEU:HD11	1.62	0.80
1:1:75:LEU:H	1:1:106:GLY:HA2	1.47	0.79
2:F:80:GLN:NE2	3:F:201:HOH:O	2.20	0.74
2:4:25:ASP:OD1	2:4:46:SER:HB2	1.92	0.70

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:2:17:GLU:OE1	1:2:164:ARG:NH2[4_554]	1.99	0.21
2:5:46:SER:OG	2:F:12:ASP:OD2[3_554]	2.16	0.04
2:5:13:ARG:NE	2:F:21:THR:O[3_554]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	1	165/218 (76%)	156 (94%)	9 (6%)	0	100	100
1	2	167/218 (77%)	154 (92%)	13 (8%)	0	100	100
2	3	104/122 (85%)	99 (95%)	5 (5%)	0	100	100
2	4	102/122 (84%)	98 (96%)	4 (4%)	0	100	100
2	5	119/122 (98%)	118 (99%)	1 (1%)	0	100	100
2	F	119/122 (98%)	113 (95%)	6 (5%)	0	100	100
All	All	776/924 (84%)	738 (95%)	38 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	1	127/172 (74%)	126 (99%)	1 (1%)	81	94

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	2	129/172 (75%)	129 (100%)	0	100	100
2	3	73/89 (82%)	73 (100%)	0	100	100
2	4	71/89 (80%)	71 (100%)	0	100	100
2	5	88/89 (99%)	88 (100%)	0	100	100
2	F	88/89 (99%)	88 (100%)	0	100	100
All	All	576/700 (82%)	575 (100%)	1 (0%)	93	98

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	1	79	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	1	94	GLN
1	2	94	GLN
2	F	0	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	1	167/218 (76%)	0.21	7 (4%) 36 32	54, 71, 98, 105	0
1	2	169/218 (77%)	0.31	9 (5%) 26 22	46, 68, 97, 129	0
2	3	106/122 (86%)	0.05	2 (1%) 66 65	41, 55, 76, 105	0
2	4	104/122 (85%)	0.13	2 (1%) 66 65	44, 55, 67, 85	0
2	5	121/122 (99%)	0.03	2 (1%) 70 69	41, 51, 78, 101	0
2	F	121/122 (99%)	0.17	4 (3%) 46 41	44, 55, 72, 110	0
All	All	788/924 (85%)	0.17	26 (3%) 46 41	41, 58, 92, 129	0

The worst 5 of 26 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	3	115	GLU	4.3
2	5	116	THR	4.0
1	2	184	PRO	3.9
1	2	55	GLY	3.5
2	3	116	THR	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.