



# Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 05:21 am BST

PDB ID : 6D91  
Title : Crystal structure of the *Deinococcus radiodurans* Nramp/MntH divalent transition metal transporter in the outward-open, apo conformation  
Authors : Bozzi, A.T.; Zimanyi, C.M.; Nicoludis, J.M.; Gaudet, R.  
Deposited on : 2018-04-27  
Resolution : 2.36 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.36 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	412	<div> <div>16%</div> <div>77%</div> <div>18%</div> <div>..</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6256 atoms, of which 3141 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Divalent metal cation transporter MntH.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	398	Total	C	H	N	O	S	0	0	0
			6153	1985	3141	505	505	17			

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	25	MET	-	expression tag	UNP Q9RTP8
A	26	HIS	-	expression tag	UNP Q9RTP8
A	27	HIS	-	expression tag	UNP Q9RTP8
A	28	HIS	-	expression tag	UNP Q9RTP8
A	29	HIS	-	expression tag	UNP Q9RTP8
A	30	HIS	-	expression tag	UNP Q9RTP8
A	31	HIS	-	expression tag	UNP Q9RTP8
A	32	HIS	-	expression tag	UNP Q9RTP8
A	33	HIS	-	expression tag	UNP Q9RTP8
A	34	MET	-	expression tag	UNP Q9RTP8
A	223	TRP	GLY	engineered mutation	UNP Q9RTP8

- Molecule 2 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLC) (formula: C<sub>21</sub>H<sub>40</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 21 17 4	0	0
2	A	1	Total C O 15 11 4	0	0
2	A	1	Total C O 15 11 4	0	0
2	A	1	Total C O 13 9 4	0	0
2	A	1	Total C 8 8	0	0
2	A	1	Total C 14 14	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	17	Total O 17 17	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Divalent metal cation transporter MntH



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	101.92Å 75.51Å 53.15Å 90.00° 98.07° 90.00°	Depositor
Resolution (Å)	38.14 – 2.36 38.14 – 2.36	Depositor EDS
% Data completeness (in resolution range)	64.0 (38.14-2.36) 64.0 (38.14-2.36)	Depositor EDS
$R_{merge}$	0.18	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.60 (at 2.37Å)	Xtriage
Refinement program	PHENIX (1.14_3211: ???)	Depositor
R, $R_{free}$	0.244 , 0.299 0.244 , 0.300	Depositor DCC
$R_{free}$ test set	1061 reflections (10.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	37.6	Xtriage
Anisotropy	0.095	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 49.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	6256	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.07% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: OLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.25	0/3080	0.40	0/4209

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3012	3141	3167	66	0
2	A	86	0	114	23	0
3	A	17	0	0	1	0
All	All	3115	3141	3281	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:261:ARG:NH2	2:A:501:OLC:H2	1.85	0.90

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:261:ARG:HD3	2:A:501:OLC:H4A	1.60	0.83
1:A:88:ILE:HD12	1:A:389:LEU:HD22	1.63	0.79
1:A:411:PHE:CZ	2:A:501:OLC:H5A	2.24	0.72
1:A:87:VAL:HG23	2:A:501:OLC:H11	1.74	0.68
1:A:339:MET:SD	1:A:352:ARG:NH2	2.66	0.68
2:A:502:OLC:O25	2:A:502:OLC:O20	2.09	0.66
1:A:192:VAL:HG11	1:A:315:LEU:HD13	1.79	0.65
1:A:94:ASN:ND2	2:A:501:OLC:H2A	2.10	0.65
1:A:375:ILE:HG23	1:A:433:LEU:HD22	1.77	0.64
1:A:287:HIS:NE2	3:A:602:HOH:O	2.30	0.64
1:A:261:ARG:CD	2:A:501:OLC:H4A	2.28	0.63
1:A:411:PHE:CE1	2:A:501:OLC:H3	2.34	0.63
2:A:502:OLC:H2A	2:A:504:OLC:O23	2.00	0.61
1:A:64:ILE:HD11	1:A:316:PHE:HE2	1.67	0.59
1:A:197:ALA:HA	1:A:308:LEU:HD21	1.85	0.58
1:A:197:ALA:HB1	1:A:284:ALA:HB3	1.86	0.56
1:A:411:PHE:CE1	2:A:501:OLC:H5A	2.40	0.56
1:A:44:LEU:O	1:A:48:VAL:HG23	2.07	0.55
1:A:87:VAL:HG13	1:A:411:PHE:CZ	2.42	0.55
1:A:103:LEU:HD11	1:A:391:PRO:HB2	1.89	0.55
1:A:156:VAL:O	1:A:160:VAL:HG12	2.07	0.54
1:A:48:VAL:HG21	2:A:503:OLC:H2	1.89	0.54
1:A:212:LEU:HD13	1:A:218:ALA:HA	1.89	0.54
1:A:113:ARG:N	1:A:114:PRO:HD2	2.23	0.54
2:A:502:OLC:H7	2:A:504:OLC:H2A	1.90	0.52
1:A:377:SER:OG	1:A:378:GLN:NE2	2.43	0.51
1:A:296:ASP:OD2	1:A:298:THR:HG22	2.11	0.51
1:A:225:ILE:HG21	1:A:425:LEU:HD12	1.93	0.50
1:A:64:ILE:HD11	1:A:316:PHE:CE2	2.47	0.50
1:A:200:ASP:O	1:A:204:VAL:HG23	2.12	0.50
1:A:88:ILE:HD12	1:A:389:LEU:CD2	2.36	0.50
1:A:90:ASN:ND2	2:A:501:OLC:H5	2.27	0.49
1:A:366:LEU:HD22	1:A:368:MET:HG2	1.94	0.48
1:A:52:ILE:HD12	1:A:55:MET:HG3	1.95	0.48
1:A:411:PHE:CE1	2:A:501:OLC:C3	2.97	0.48
1:A:265:ILE:HD11	2:A:501:OLC:H7	1.95	0.48
1:A:271:ALA:O	1:A:275:ASN:ND2	2.47	0.47
1:A:370:PRO:O	1:A:374:LEU:N	2.40	0.47
1:A:416:TRP:O	1:A:420:VAL:HG23	2.15	0.47
2:A:506:OLC:H14	2:A:506:OLC:H11	1.54	0.46
1:A:87:VAL:CG2	2:A:501:OLC:H11	2.45	0.46

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:54:TYR:OH	1:A:89:GLN:OE1	2.27	0.45
1:A:94:ASN:HD22	2:A:501:OLC:H2A	1.81	0.45
1:A:236:LEU:HD21	1:A:333:MET:HE2	1.99	0.44
1:A:248:ASP:OD1	1:A:248:ASP:N	2.51	0.44
1:A:192:VAL:CG1	1:A:315:LEU:HD13	2.46	0.44
1:A:379:VAL:HG22	1:A:433:LEU:HD13	2.00	0.44
1:A:297:LEU:HD23	1:A:316:PHE:HZ	1.84	0.43
1:A:160:VAL:HG13	1:A:161:THR:N	2.34	0.43
1:A:87:VAL:HG13	1:A:411:PHE:HZ	1.83	0.43
1:A:113:ARG:HG3	1:A:114:PRO:CD	2.49	0.43
1:A:213:GLN:OE1	1:A:213:GLN:N	2.52	0.43
1:A:102:ASN:HB3	1:A:239:ALA:HA	2.01	0.43
1:A:374:LEU:O	1:A:378:GLN:NE2	2.42	0.43
1:A:257:VAL:HG11	2:A:501:OLC:H21A	2.00	0.42
1:A:113:ARG:HG3	1:A:114:PRO:HD3	2.01	0.42
1:A:394:LEU:CD1	2:A:505:OLC:H6	2.50	0.42
1:A:143:GLN:O	1:A:147:GLY:N	2.52	0.41
1:A:184:LEU:O	1:A:188:VAL:HG22	2.21	0.41
1:A:315:LEU:HA	1:A:318:VAL:HG22	2.02	0.41
1:A:351:LEU:HD22	1:A:355:ILE:HD11	2.03	0.41
1:A:91:LEU:CD2	1:A:393:LEU:HD23	2.50	0.41
1:A:261:ARG:HD2	2:A:501:OLC:H6A	2.02	0.41
1:A:213:GLN:HG2	1:A:213:GLN:O	2.21	0.41
1:A:225:ILE:CG2	1:A:425:LEU:HD12	2.50	0.41
1:A:223:TRP:CD1	1:A:429:LEU:HD22	2.56	0.41
2:A:503:OLC:H6	2:A:503:OLC:H3A	1.88	0.41
1:A:411:PHE:CD1	2:A:501:OLC:H3A	2.56	0.41
1:A:161:THR:O	1:A:165:LEU:HD12	2.20	0.40
1:A:427:GLY:O	1:A:431:TRP:N	2.52	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	396/412 (96%)	379 (96%)	17 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	310/323 (96%)	302 (97%)	8 (3%)	46	56

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	102	ASN
1	A	168	GLN
1	A	177	LEU
1	A	200	ASP
1	A	223	TRP
1	A	270	LEU
1	A	383	PHE
1	A	433	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	90	ASN
1	A	102	ASN
1	A	426	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	OLC	A	505	-	7,7,24	0.45	0	6,6,25	0.79	0
2	OLC	A	503	-	14,14,24	1.02	2 (14%)	15,15,25	1.06	1 (6%)
2	OLC	A	504	-	12,12,24	1.09	2 (16%)	13,13,25	1.39	1 (7%)
2	OLC	A	502	-	14,14,24	1.00	2 (14%)	15,15,25	1.21	1 (6%)
2	OLC	A	506	-	13,13,24	0.37	0	12,12,25	0.55	0
2	OLC	A	501	-	20,20,24	0.88	2 (10%)	21,21,25	0.89	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OLC	A	505	-	-	3/5/5/24	-
2	OLC	A	503	-	-	7/14/14/24	-
2	OLC	A	504	-	-	3/12/12/24	-
2	OLC	A	502	-	-	3/14/14/24	-
2	OLC	A	506	-	-	4/11/11/24	-
2	OLC	A	501	-	-	11/20/20/24	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	504	OLC	O20-C1	2.46	1.40	1.33
2	A	503	OLC	O20-C1	2.42	1.40	1.33
2	A	501	OLC	O20-C1	2.41	1.40	1.33
2	A	502	OLC	O20-C1	2.38	1.40	1.33
2	A	503	OLC	O20-C21	-2.10	1.40	1.45
2	A	501	OLC	O20-C21	-2.10	1.40	1.45
2	A	502	OLC	O20-C21	-2.06	1.40	1.45
2	A	504	OLC	O20-C21	-2.00	1.40	1.45

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	504	OLC	O20-C1-C2	3.31	122.30	111.91
2	A	502	OLC	O20-C1-C2	2.93	121.11	111.91
2	A	503	OLC	O20-C1-C2	2.73	120.48	111.91
2	A	501	OLC	O20-C1-C2	2.54	119.88	111.91

There are no chirality outliers.

All (31) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	503	OLC	C21-C22-C24-O25
2	A	503	OLC	O23-C22-C24-O25
2	A	504	OLC	O20-C21-C22-C24
2	A	504	OLC	O20-C21-C22-O23
2	A	501	OLC	O20-C21-C22-O23
2	A	501	OLC	O19-C1-O20-C21
2	A	501	OLC	C2-C1-O20-C21
2	A	503	OLC	C2-C1-O20-C21
2	A	506	OLC	C11-C12-C13-C14
2	A	503	OLC	O19-C1-O20-C21
2	A	502	OLC	C1-C2-C3-C4
2	A	501	OLC	C1-C2-C3-C4
2	A	503	OLC	C1-C2-C3-C4
2	A	503	OLC	C2-C3-C4-C5
2	A	501	OLC	C5-C6-C7-C8
2	A	501	OLC	C21-C22-C24-O25
2	A	502	OLC	C2-C1-O20-C21
2	A	506	OLC	C3-C4-C5-C6
2	A	503	OLC	C3-C4-C5-C6
2	A	506	OLC	C10-C11-C12-C13

*Continued on next page...*

*Continued from previous page...*




Mol	Chain	Res	Type	Atoms
2	A	501	OLC	O20-C21-C22-C24
2	A	501	OLC	C4-C5-C6-C7
2	A	502	OLC	O19-C1-O20-C21
2	A	501	OLC	C2-C3-C4-C5
2	A	506	OLC	C4-C5-C6-C7
2	A	501	OLC	O23-C22-C24-O25
2	A	504	OLC	C3-C4-C5-C6
2	A	505	OLC	C6-C7-C8-C9
2	A	505	OLC	C7-C8-C9-C10
2	A	505	OLC	C4-C5-C6-C7
2	A	501	OLC	C7-C8-C9-C10


There are no ring outliers.

6 monomers are involved in 23 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	505	OLC	1	0
2	A	503	OLC	2	0
2	A	504	OLC	2	0
2	A	502	OLC	3	0
2	A	506	OLC	1	0
2	A	501	OLC	16	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	398/412 (96%)	1.14	66 (16%) ⓘ ⓘ	24, 41, 92, 287	0

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	371	SER	17.4
1	A	39	ARG	11.2
1	A	372	SER	8.2
1	A	370	PRO	7.8
1	A	368	MET	7.1
1	A	373	VAL	6.7
1	A	40	ILE	5.9
1	A	162	PHE	4.9
1	A	376	LEU	4.8
1	A	216	GLY	4.8
1	A	169	LYS	4.6
1	A	214	GLY	4.6
1	A	375	ILE	4.6
1	A	436	GLY	4.5
1	A	165	LEU	4.5
1	A	171	GLY	4.3
1	A	298	THR	4.3
1	A	246	GLN	4.2
1	A	367	GLY	4.2
1	A	248	ASP	4.2
1	A	366	LEU	4.1
1	A	369	ASP	3.9
1	A	149	PRO	3.8
1	A	163	TRP	3.8
1	A	435	GLY	3.8
1	A	346	HIS	3.7
1	A	167	LEU	3.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	201	LEU	3.7
1	A	251	GLU	3.6
1	A	151	PHE	3.6
1	A	377	SER	3.5
1	A	245	ILE	3.5
1	A	173	ARG	3.4
1	A	292	GLU	3.4
1	A	168	GLN	3.3
1	A	291	VAL	3.2
1	A	431	TRP	3.2
1	A	374	LEU	2.9
1	A	345	PHE	2.8
1	A	290	ASN	2.8
1	A	172	THR	2.8
1	A	223	TRP	2.7
1	A	166	ASN	2.7
1	A	164	LEU	2.6
1	A	365	LEU	2.6
1	A	170	ARG	2.5
1	A	41	LEU	2.5
1	A	192	VAL	2.5
1	A	148	LEU	2.5
1	A	384	GLY	2.4
1	A	296	ASP	2.4
1	A	154	ALA	2.4
1	A	350	TRP	2.3
1	A	227	ALA	2.3
1	A	383	PHE	2.3
1	A	293	ASN	2.2
1	A	430	LEU	2.2
1	A	399	ARG	2.1
1	A	401	VAL	2.1
1	A	152	TRP	2.1
1	A	52	ILE	2.1
1	A	379	VAL	2.1
1	A	132	LEU	2.0
1	A	364	ILE	2.0
1	A	297	LEU	2.0
1	A	299	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	OLC	A	501	21/25	0.76	0.24	39,51,70,73	0
2	OLC	A	506	14/25	0.78	0.32	44,47,48,48	0
2	OLC	A	503	15/25	0.86	0.17	50,56,59,60	0
2	OLC	A	504	13/25	0.88	0.17	42,47,49,50	0
2	OLC	A	502	15/25	0.89	0.25	47,48,50,51	0
2	OLC	A	505	8/25	0.91	0.19	43,44,46,46	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around OLC A 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around OLC A 506:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around OLC A 503:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around OLC A 504:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around OLC A 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around OLC A 505:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers

There are no such residues in this entry.