



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 25, 2020 – 03:17 pm BST

PDB ID : 3DBN  
Title : Crystal structure of the Streptococcus suis serotype2 D-mannonate dehydratase in complex with its substrate  
Authors : Peng, H.; Zhang, Q.; Gao, F.; Gao, G.F.  
Deposited on : 2008-06-02  
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	386	<div> <div>2%</div> <div> <div></div> <div>54%</div> <div>30%</div> <div>6%</div> <div>10%</div> </div> </div>
1	B	386	<div> <div>2%</div> <div> <div></div> <div>53%</div> <div>33%</div> <div>5%</div> <div>10%</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5484 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mannonate dehydratase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	349	Total	C	N	O	S	0	0	0
			2737	1745	465	513	14			
1	B	349	Total	C	N	O	S	0	0	0
			2732	1742	465	512	13			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	EXPRESSION TAG	UNP A4VVI4
A	2	GLY	-	EXPRESSION TAG	UNP A4VVI4
A	3	SER	-	EXPRESSION TAG	UNP A4VVI4
A	4	SER	-	EXPRESSION TAG	UNP A4VVI4
A	5	HIS	-	EXPRESSION TAG	UNP A4VVI4
A	6	HIS	-	EXPRESSION TAG	UNP A4VVI4
A	7	HIS	-	EXPRESSION TAG	UNP A4VVI4
A	8	HIS	-	EXPRESSION TAG	UNP A4VVI4
A	9	HIS	-	EXPRESSION TAG	UNP A4VVI4
A	10	HIS	-	EXPRESSION TAG	UNP A4VVI4
A	11	SER	-	EXPRESSION TAG	UNP A4VVI4
A	12	SER	-	EXPRESSION TAG	UNP A4VVI4
A	13	GLY	-	EXPRESSION TAG	UNP A4VVI4
A	14	LEU	-	EXPRESSION TAG	UNP A4VVI4
A	15	VAL	-	EXPRESSION TAG	UNP A4VVI4
A	16	PRO	-	EXPRESSION TAG	UNP A4VVI4
A	17	ARG	-	EXPRESSION TAG	UNP A4VVI4
A	18	GLY	-	EXPRESSION TAG	UNP A4VVI4
A	19	SER	-	EXPRESSION TAG	UNP A4VVI4
A	20	HIS	-	EXPRESSION TAG	UNP A4VVI4
B	1	MET	-	EXPRESSION TAG	UNP A4VVI4
B	2	GLY	-	EXPRESSION TAG	UNP A4VVI4
B	3	SER	-	EXPRESSION TAG	UNP A4VVI4
B	4	SER	-	EXPRESSION TAG	UNP A4VVI4
B	5	HIS	-	EXPRESSION TAG	UNP A4VVI4

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	6	HIS	-	EXPRESSION TAG	UNP A4VVI4
B	7	HIS	-	EXPRESSION TAG	UNP A4VVI4
B	8	HIS	-	EXPRESSION TAG	UNP A4VVI4
B	9	HIS	-	EXPRESSION TAG	UNP A4VVI4
B	10	HIS	-	EXPRESSION TAG	UNP A4VVI4
B	11	SER	-	EXPRESSION TAG	UNP A4VVI4
B	12	SER	-	EXPRESSION TAG	UNP A4VVI4
B	13	GLY	-	EXPRESSION TAG	UNP A4VVI4
B	14	LEU	-	EXPRESSION TAG	UNP A4VVI4
B	15	VAL	-	EXPRESSION TAG	UNP A4VVI4
B	16	PRO	-	EXPRESSION TAG	UNP A4VVI4
B	17	ARG	-	EXPRESSION TAG	UNP A4VVI4
B	18	GLY	-	EXPRESSION TAG	UNP A4VVI4
B	19	SER	-	EXPRESSION TAG	UNP A4VVI4
B	20	HIS	-	EXPRESSION TAG	UNP A4VVI4

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Mn 1 1	0	0
2	A	1	Total Mn 1 1	0	0

- Molecule 3 is D-MANNONIC ACID (three-letter code: CS2) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	6	7		



T342	P343	G344	Y348	D349	L352	T355	Y356	F357	L360	Y361	E362	A363	N364	M365	K370	T371	P372	D373	F374	K384	E385	GLY
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	105.52Å 105.52Å 160.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.90 25.87 – 2.90	Depositor EDS
% Data completeness (in resolution range)	95.8 (50.00-2.90) 96.0 (25.87-2.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	8.03 (at 2.89Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.231 , 0.280 0.230 , 0.281	Depositor DCC
$R_{free}$ test set	981 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.4	Xtriage
Anisotropy	0.019	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 39.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.31$ , $\langle L^2 \rangle = 0.14$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.82	EDS
Total number of atoms	5484	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.74% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN, CS2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.65	1/2804 (0.0%)	0.76	0/3807
1	B	0.64	0/2799	0.76	0/3801
All	All	0.65	1/5603 (0.0%)	0.76	0/7608

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	238	GLU	CB-CG	5.90	1.63	1.52

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	85	SER	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2737	0	2676	115	0
1	B	2732	0	2678	106	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	13	0	10	2	0
All	All	5484	0	5364	212	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 212 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:182:ILE:HG22	1:B:186:ARG:HD2	1.31	1.06
1:B:287:THR:HG23	1:B:287:THR:O	1.57	1.01
1:A:226:GLY:HA2	1:A:232:ARG:HD3	1.48	0.94
1:A:250:GLU:HG2	1:A:280:ASN:HD22	1.31	0.92
1:A:57:VAL:HG13	1:A:58:GLY:H	1.30	0.92

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	345/386 (89%)	290 (84%)	43 (12%)	12 (4%)	3	14
1	B	345/386 (89%)	301 (87%)	33 (10%)	11 (3%)	4	16
All	All	690/772 (89%)	591 (86%)	76 (11%)	23 (3%)	4	15

5 of 23 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	57	VAL
1	A	284	PHE
1	A	331	HIS
1	A	343	PRO
1	B	343	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/321 (90%)	263 (91%)	25 (9%)	10	30
1	B	288/321 (90%)	261 (91%)	27 (9%)	8	26
All	All	576/642 (90%)	524 (91%)	52 (9%)	9	29

5 of 52 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	341	LYS
1	B	79	GLU
1	B	341	LYS
1	A	381	VAL
1	B	25	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	105	ASN
1	B	219	HIS
1	B	324	GLN
1	B	137	HIS
1	B	237	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	CS2	A	388	2	9,12,12	0.74	0	12,16,16	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CS2	A	388	2	-	7/14/18/18	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	388	CS2	C1-C2-C3-O3
3	A	388	CS2	O2-C2-C3-O3
3	A	388	CS2	O2-C2-C3-C4
3	A	388	CS2	C4-C5-C6-O6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	A	388	CS2	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	388	CS2	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	349/386 (90%)	-0.15	6 (1%) 70 69	3, 22, 42, 49	0
1	B	349/386 (90%)	-0.22	8 (2%) 60 58	2, 20, 42, 63	0
All	All	698/772 (90%)	-0.19	14 (2%) 65 63	2, 21, 42, 63	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	180	ALA	4.4
1	A	56	PRO	3.6
1	B	148	ALA	2.9
1	B	340	THR	2.8
1	B	293	GLY	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CS2	A	388	13/13	0.93	0.17	41,44,46,47	0
2	MN	B	387	1/1	0.95	0.13	52,52,52,52	0
2	MN	A	387	1/1	0.95	0.06	31,31,31,31	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.