



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 04:34 pm BST

PDB ID : 5DBF
Title : Crystal Structure of Iridoid Synthase from *Cantharanthus roseus* in complex with NADPH
Authors : Hu, Y.M.; Liu, W.D.; Zheng, Y.Y.; Xu, Z.X.; Ko, T.P.; Chen, C.C.; Guo, R.T.
Deposited on : 2015-08-21
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	374	<div> <div>6%</div> <div> <div></div> <div>84%</div> <div>11%</div> <div>..</div> </div> </div>
1	B	374	<div> <div>7%</div> <div> <div></div> <div>86%</div> <div>9%</div> <div>..</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6094 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Iridoid synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	359	Total	C	N	O	S	0	0	0
			2849	1835	469	526	19			
1	B	360	Total	C	N	O	S	0	0	0
			2858	1840	470	529	19			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	389	ALA	-	expression tag	UNP K7WDL7
A	390	SER	-	expression tag	UNP K7WDL7
A	391	ALA	-	expression tag	UNP K7WDL7
A	392	TRP	-	expression tag	UNP K7WDL7
A	393	SER	-	expression tag	UNP K7WDL7
A	394	HIS	-	expression tag	UNP K7WDL7
A	395	PRO	-	expression tag	UNP K7WDL7
A	396	GLN	-	expression tag	UNP K7WDL7
A	397	PHE	-	expression tag	UNP K7WDL7
A	398	GLU	-	expression tag	UNP K7WDL7
A	399	LYS	-	expression tag	UNP K7WDL7
B	389	ALA	-	expression tag	UNP K7WDL7
B	390	SER	-	expression tag	UNP K7WDL7
B	391	ALA	-	expression tag	UNP K7WDL7
B	392	TRP	-	expression tag	UNP K7WDL7
B	393	SER	-	expression tag	UNP K7WDL7
B	394	HIS	-	expression tag	UNP K7WDL7
B	395	PRO	-	expression tag	UNP K7WDL7
B	396	GLN	-	expression tag	UNP K7WDL7
B	397	PHE	-	expression tag	UNP K7WDL7
B	398	GLU	-	expression tag	UNP K7WDL7
B	399	LYS	-	expression tag	UNP K7WDL7


- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: C₂₁H₃₀N₇O₁₇P₃).




3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Iridoid synthase



• Molecule 1: Iridoid synthase



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	91.77 Å 94.89 Å 172.60 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 2.00 24.58 – 2.00	Depositor EDS
% Data completeness (in resolution range)	97.1 (25.00-2.00) 97.2 (24.58-2.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.18 (at 1.99 Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.192 , 0.246 0.200 , 0.248	Depositor DCC
R_{free} test set	2403 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	33.9	Xtriage
Anisotropy	0.848	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 54.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.116 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6094	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.71	0/2923	0.81	6/3969 (0.2%)
1	B	0.71	0/2932	0.79	5/3981 (0.1%)
All	All	0.71	0/5855	0.80	11/7950 (0.1%)

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	249	VAL	CB-CA-C	-7.84	96.50	111.40
1	B	249	VAL	CB-CA-C	-7.01	98.08	111.40
1	A	249	VAL	CG1-CB-CG2	6.26	120.91	110.90
1	A	110	ILE	C-N-CA	-5.64	110.44	122.30
1	B	249	VAL	CG1-CB-CG2	5.61	119.88	110.90
1	B	256	GLU	OE1-CD-OE2	-5.47	116.73	123.30
1	B	110	ILE	C-N-CA	-5.33	111.10	122.30
1	B	278	ASP	CB-CG-OD1	5.32	123.09	118.30
1	A	337	ILE	CB-CA-C	-5.14	101.32	111.60
1	A	385	ARG	NE-CZ-NH1	5.13	122.86	120.30
1	A	278	ASP	CB-CG-OD1	5.06	122.86	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2849	0	2812	16	1
1	B	2858	0	2818	17	0
2	A	48	0	26	0	0
2	B	48	0	26	1	0
3	A	144	0	0	1	0
3	B	147	0	0	1	0
All	All	6094	0	5682	33	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:110:ILE:HG23	1:A:121:MET:SD	2.29	0.73
1:B:159:VAL:O	1:B:159:VAL:HG22	1.92	0.69
1:B:249:VAL:HG13	1:B:275:ASN:HA	1.73	0.69
1:A:249:VAL:HG13	1:A:275:ASN:HA	1.79	0.64
1:B:92:ILE:O	1:B:96:SER:OG	2.16	0.56
1:B:97:PRO:HA	1:B:99:LYS:HE2	1.89	0.55
1:B:214:ASN:C	1:B:214:ASN:HD22	2.09	0.55
1:B:163:SER:HB2	1:B:164:PRO:HA	1.90	0.54
1:A:97:PRO:HA	1:A:99:LYS:HE2	1.90	0.54
1:A:163:SER:HB2	1:A:164:PRO:HA	1.91	0.52
1:A:327:ASN:O	1:A:385:ARG:NH2	2.44	0.50
1:A:114:ASP:HB3	1:A:117:THR:OG1	2.12	0.49
1:B:114:ASP:HB3	1:B:117:THR:OG1	2.12	0.49
1:A:214:ASN:HD22	1:A:214:ASN:C	2.16	0.49
1:B:328:ASN:O	1:B:385:ARG:NH2	2.48	0.47
1:A:41:VAL:HG11	1:A:142:GLN:HE22	1.80	0.47
1:B:249:VAL:HG13	1:B:275:ASN:CA	2.43	0.46
1:B:214:ASN:ND2	3:B:508:HOH:O	2.49	0.46
1:B:306:GLN:OE1	1:B:348:CYS:HB3	2.16	0.45
1:A:334:LEU:HA	1:A:337:ILE:HD11	1.98	0.45
1:A:284:HIS:HD2	3:A:588:HOH:O	1.99	0.45
1:A:296:GLU:HG2	1:A:297:MET:N	2.32	0.44
1:B:159:VAL:CG2	1:B:159:VAL:O	2.64	0.44
1:B:96:SER:N	1:B:97:PRO:CD	2.81	0.44
1:A:96:SER:OG	1:A:97:PRO:HD3	2.19	0.43
1:A:237:GLY:HA3	1:A:347:PHE:HB3	2.01	0.42
1:B:217:SER:HB3	1:B:381:MET:SD	2.59	0.42
1:B:85:VAL:HG13	2:B:401:NDP:H2A	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:306:GLN:OE1	1:A:348:CYS:HB3	2.19	0.42
1:B:95:LEU:HA	1:B:98:LEU:HD22	2.02	0.42
1:B:85:VAL:HG23	1:B:121:MET:HG2	2.03	0.41
1:A:217:SER:HB3	1:A:381:MET:SD	2.61	0.41
1:A:225:ILE:CD1	1:A:313:MET:CE	2.99	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:325:LYS:O	1:A:325:LYS:O[4_555]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	355/374 (95%)	345 (97%)	10 (3%)	0	100	100
1	B	356/374 (95%)	347 (98%)	9 (2%)	0	100	100
All	All	711/748 (95%)	692 (97%)	19 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	318/331 (96%)	298 (94%)	20 (6%)	18	13
1	B	319/331 (96%)	300 (94%)	19 (6%)	19	14
All	All	637/662 (96%)	598 (94%)	39 (6%)	18	14

All (39) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	33	LEU
1	A	73	LEU
1	A	78	VAL
1	A	98	LEU
1	A	110	ILE
1	A	159	VAL
1	A	172	LEU
1	A	185	LEU
1	A	214	ASN
1	A	234	VAL
1	A	249	VAL
1	A	289	LEU
1	A	315	ASP
1	A	317	ASP
1	A	326	LYS
1	A	329	LEU
1	A	334	LEU
1	A	337	ILE
1	A	362	LEU
1	A	387	ILE
1	B	29	LYS
1	B	33	LEU
1	B	73	LEU
1	B	96	SER
1	B	98	LEU
1	B	154	GLU
1	B	159	VAL
1	B	172	LEU
1	B	185	LEU
1	B	214	ASN
1	B	249	VAL
1	B	289	LEU
1	B	315	ASP
1	B	317	ASP
1	B	326	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	329	LEU
1	B	334	LEU
1	B	362	LEU
1	B	387	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	82	GLN
1	A	142	GLN
1	A	192	ASN
1	A	214	ASN
1	B	82	GLN
1	B	132	ASN
1	B	192	ASN
1	B	214	ASN
1	B	284	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NDP	B	401	-	45,52,52	1.18	5 (11%)	53,80,80	1.38	7 (13%)
2	NDP	A	401	-	45,52,52	1.02	1 (2%)	53,80,80	1.54	8 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	B	401	-	-	8/30/77/77	0/5/5/5
2	NDP	A	401	-	-	7/30/77/77	0/5/5/5

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	NDP	C5A-C4A	3.07	1.49	1.40
2	B	401	NDP	C6N-C5N	2.86	1.38	1.33
2	B	401	NDP	C5A-C4A	2.61	1.47	1.40
2	B	401	NDP	O4B-C1B	2.52	1.44	1.41
2	B	401	NDP	C4N-C5N	-2.52	1.42	1.48
2	B	401	NDP	C2A-N1A	2.45	1.38	1.33

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	NDP	N3A-C2A-N1A	-4.66	121.39	128.68
2	B	401	NDP	C4A-C5A-N7A	-4.01	105.22	109.40
2	A	401	NDP	C1B-N9A-C4A	-3.36	120.73	126.64
2	A	401	NDP	C2A-N1A-C6A	3.28	124.36	118.75
2	B	401	NDP	N3A-C2A-N1A	-3.15	123.75	128.68
2	A	401	NDP	O4D-C1D-N1N	3.09	114.10	108.06
2	A	401	NDP	C1D-N1N-C6N	-2.76	114.89	120.83
2	A	401	NDP	O3X-P2B-O2X	2.71	118.00	107.64
2	A	401	NDP	O2N-PN-O1N	2.65	125.35	112.24
2	B	401	NDP	O2B-P2B-O1X	-2.42	100.05	109.39
2	B	401	NDP	O2X-P2B-O1X	2.23	119.41	110.68
2	B	401	NDP	C1D-N1N-C6N	-2.23	116.03	120.83
2	A	401	NDP	C3D-C2D-C1D	-2.16	97.31	101.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	NDP	C5A-C6A-N1A	-2.13	115.53	120.35
2	B	401	NDP	O4B-C4B-C3B	2.08	109.23	105.11

There are no chirality outliers.

All (15) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	401	NDP	C5D-O5D-PN-O2N
2	A	401	NDP	C5D-O5D-PN-O2N
2	B	401	NDP	O4B-C4B-C5B-O5B
2	B	401	NDP	C2B-O2B-P2B-O2X
2	B	401	NDP	C5D-O5D-PN-O3
2	A	401	NDP	C2B-O2B-P2B-O2X
2	A	401	NDP	C5D-O5D-PN-O3
2	B	401	NDP	C5D-O5D-PN-O1N
2	B	401	NDP	C2D-C1D-N1N-C6N
2	B	401	NDP	O4D-C1D-N1N-C6N
2	A	401	NDP	O4D-C1D-N1N-C6N
2	A	401	NDP	O4B-C4B-C5B-O5B
2	B	401	NDP	C3B-C4B-C5B-O5B
2	A	401	NDP	C2D-C1D-N1N-C6N
2	A	401	NDP	C5D-O5D-PN-O1N


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	NDP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	359/374 (95%)	0.31	24 (6%) 17 17	30, 43, 65, 84	0
1	B	360/374 (96%)	0.27	27 (7%) 14 13	30, 44, 66, 94	0
All	All	719/748 (96%)	0.29	51 (7%) 16 15	30, 44, 66, 94	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	159	VAL	5.5
1	A	305	GLU	3.9
1	B	303	GLY	3.7
1	A	141	LEU	3.7
1	A	104	ILE	3.6
1	A	139	VAL	3.6
1	A	219	LEU	3.5
1	B	219	LEU	3.5
1	B	140	CYS	3.5
1	B	151	ILE	3.3
1	B	301	VAL	3.3
1	A	221	VAL	3.1
1	A	140	CYS	3.1
1	B	304	LYS	3.1
1	B	141	LEU	3.0
1	B	305	GLU	2.9
1	A	304	LYS	2.9
1	A	222	TYR	2.8
1	A	105	PHE	2.8
1	B	153	GLU	2.8
1	B	173	ASN	2.7
1	A	218	THR	2.6
1	A	318	GLN	2.6
1	B	172	LEU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	390	SER	2.5
1	A	106	TYR	2.5
1	A	35	VAL	2.4
1	B	106	TYR	2.4
1	B	35	VAL	2.4
1	B	37	VAL	2.3
1	B	159	VAL	2.3
1	A	173	ASN	2.3
1	B	33	LEU	2.3
1	A	193	ASN	2.3
1	A	33	LEU	2.2
1	B	105	PHE	2.2
1	B	311	GLU	2.2
1	B	218	THR	2.2
1	B	107	VAL	2.2
1	A	223	ALA	2.2
1	B	104	ILE	2.2
1	A	107	VAL	2.2
1	A	307	VAL	2.2
1	B	160	PRO	2.2
1	B	223	ALA	2.1
1	B	302	GLU	2.1
1	B	318	GLN	2.1
1	A	190	GLY	2.1
1	B	139	VAL	2.0
1	B	390	SER	2.0
1	A	160	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.


6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NDP	B	401	48/48	0.96	0.10	28,38,46,52	0
2	NDP	A	401	48/48	0.98	0.08	25,36,43,53	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.