



Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2020 – 12:57 am BST

PDB ID : 3DEL
Title : The structure of CT381, the arginine binding protein from the periplasm
Chlamydia trachomatis
Authors : Petit, P.; Vuillard, L.; Spinelli, S.
Deposited on : 2008-06-10
Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	7937 (1.94-1.90)
Clashscore	141614	8644 (1.94-1.90)
Ramachandran outliers	138981	8530 (1.94-1.90)
Sidechain outliers	138945	8530 (1.94-1.90)
RSRZ outliers	127900	7793 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	242	<div> <div>5%</div> <div> <div></div> <div>90%</div> <div>6%</div> <div></div> </div> </div>
1	C	242	<div> <div>5%</div> <div> <div></div> <div>89%</div> <div>6%</div> <div></div> </div> </div>
1	D	242	<div> <div>4%</div> <div> <div></div> <div>88%</div> <div>7%</div> <div></div> </div> </div>
1	F	242	<div> <div>4%</div> <div> <div></div> <div>90%</div> <div>6%</div> <div></div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 8711 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Arginine Binding Protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	232	Total	C	N	O	S	0	3	0
			1856	1196	312	343	5			
1	C	232	Total	C	N	O	S	8	3	0
			1861	1198	315	344	4			
1	D	232	Total	C	N	O	S	0	4	0
			1862	1199	312	347	4			
1	F	232	Total	C	N	O	S	0	1	0
			1848	1190	312	342	4			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	258	LEU	-	EXPRESSION TAG	UNP O84385
B	259	GLU	-	EXPRESSION TAG	UNP O84385
B	260	HIS	-	EXPRESSION TAG	UNP O84385
B	261	HIS	-	EXPRESSION TAG	UNP O84385
B	262	HIS	-	EXPRESSION TAG	UNP O84385
B	263	HIS	-	EXPRESSION TAG	UNP O84385
B	264	HIS	-	EXPRESSION TAG	UNP O84385
B	265	HIS	-	EXPRESSION TAG	UNP O84385
C	258	LEU	-	EXPRESSION TAG	UNP O84385
C	259	GLU	-	EXPRESSION TAG	UNP O84385
C	260	HIS	-	EXPRESSION TAG	UNP O84385
C	261	HIS	-	EXPRESSION TAG	UNP O84385
C	262	HIS	-	EXPRESSION TAG	UNP O84385
C	263	HIS	-	EXPRESSION TAG	UNP O84385
C	264	HIS	-	EXPRESSION TAG	UNP O84385
C	265	HIS	-	EXPRESSION TAG	UNP O84385
D	258	LEU	-	EXPRESSION TAG	UNP O84385
D	259	GLU	-	EXPRESSION TAG	UNP O84385
D	260	HIS	-	EXPRESSION TAG	UNP O84385
D	261	HIS	-	EXPRESSION TAG	UNP O84385
D	262	HIS	-	EXPRESSION TAG	UNP O84385

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	263	HIS	-	EXPRESSION TAG	UNP O84385
D	264	HIS	-	EXPRESSION TAG	UNP O84385
D	265	HIS	-	EXPRESSION TAG	UNP O84385
F	258	LEU	-	EXPRESSION TAG	UNP O84385
F	259	GLU	-	EXPRESSION TAG	UNP O84385
F	260	HIS	-	EXPRESSION TAG	UNP O84385
F	261	HIS	-	EXPRESSION TAG	UNP O84385
F	262	HIS	-	EXPRESSION TAG	UNP O84385
F	263	HIS	-	EXPRESSION TAG	UNP O84385
F	264	HIS	-	EXPRESSION TAG	UNP O84385
F	265	HIS	-	EXPRESSION TAG	UNP O84385


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	301	Total O 301 301	0	0
2	C	341	Total O 341 341	0	0
2	D	324	Total O 324 324	0	0
2	F	318	Total O 318 318	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Arginine Binding Protein




• Molecule 1: Arginine Binding Protein



• Molecule 1: Arginine Binding Protein



• Molecule 1: Arginine Binding Protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	40.07Å 121.88Å 170.10Å 90.00° 95.38° 90.00°	Depositor
Resolution (Å)	30.00 – 1.92 51.22 – 1.92	Depositor EDS
% Data completeness (in resolution range)	99.5 (30.00-1.92) 99.5 (51.22-1.92)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.78 (at 1.92Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.196 , 0.213 0.218 , 0.237	Depositor DCC
R_{free} test set	1237 reflections (1.00%)	wwPDB-VP
Wilson B-factor (Å ²)	20.7	Xtriage
Anisotropy	0.650	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 38.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.032 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8711	wwPDB-VP
Average B, all atoms (Å ²)	5.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.51	0/1906	0.54	0/2585
1	C	0.52	0/1911	0.56	0/2592
1	D	0.50	0/1915	0.57	0/2598
1	F	0.52	0/1892	0.57	0/2567
All	All	0.51	0/7624	0.56	0/10342

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1856	0	1881	9	0
1	C	1861	0	1884	15	0
1	D	1862	0	1882	14	0
1	F	1848	0	1867	16	0
2	B	301	0	0	4	0
2	C	341	0	0	2	0
2	D	324	0	0	3	0
2	F	318	0	0	4	0
All	All	8711	0	7514	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:133:LYS:HD2	1:D:133:LYS:H	1.17	1.10
1:C:171:LEU:HD13	1:F:171:LEU:HD13	1.45	0.98
1:D:133:LYS:CD	1:D:133:LYS:H	1.88	0.84
1:D:133:LYS:N	1:D:133:LYS:HD2	1.96	0.81
1:D:229:LEU:HA	1:D:232:LYS:HE2	1.69	0.74
1:F:240[B]:ILE:HD11	1:F:246:LEU:HD13	1.71	0.72
1:D:165:ARG:HD3	1:D:176:GLU:OE2	1.95	0.66
1:F:229:LEU:HA	1:F:232:LYS:HE2	1.76	0.66
1:C:171:LEU:HD13	1:F:171:LEU:CD1	2.24	0.64
1:F:263:HIS:HE1	2:F:558:HOH:O	1.83	0.62
1:F:264:HIS:HD2	2:F:508:HOH:O	1.84	0.60
1:C:171:LEU:HD11	1:F:199:PHE:CZ	2.37	0.59
1:D:263:HIS:HE1	2:D:835:HOH:O	1.87	0.58
1:C:262:HIS:O	1:C:265:HIS:HD2	1.86	0.58
1:F:140:GLN:NE2	2:F:658:HOH:O	2.37	0.58
1:D:264:HIS:HD2	2:D:637:HOH:O	1.87	0.57
1:F:262:HIS:O	1:F:265:HIS:HD2	1.88	0.56
1:F:256:ASN:H	1:F:256:ASN:ND2	2.04	0.56
1:D:43:THR:HG22	1:D:43:THR:O	2.06	0.56
1:C:171:LEU:CD1	1:F:171:LEU:HD13	2.29	0.55
1:F:256:ASN:H	1:F:256:ASN:HD22	1.54	0.54
1:D:262:HIS:O	1:D:265:HIS:HD2	1.92	0.52
1:C:171:LEU:HD11	1:F:199:PHE:HZ	1.73	0.52
1:C:264:HIS:HD2	2:C:352:HOH:O	1.93	0.52
1:B:163:HIS:NE2	1:D:165:ARG:HD2	2.25	0.52
1:B:229:LEU:HA	1:B:232:LYS:HE3	1.92	0.51
1:C:143:SER:HB2	1:C:165[B]:ARG:HG3	1.92	0.51
1:D:92:HIS:HE1	1:D:225:ASP:OD1	1.95	0.49
1:C:199:PHE:HZ	1:F:171:LEU:HD11	1.77	0.49
1:C:160:SER:O	1:C:161:GLU:HG2	2.13	0.49
1:B:140:GLN:NE2	2:B:527:HOH:O	2.41	0.48
1:C:229:LEU:HA	1:C:232:LYS:HE3	1.95	0.48
1:B:264:HIS:HD2	2:B:407:HOH:O	1.96	0.48
1:C:194:VAL:O	1:C:197:LYS:HG2	2.15	0.47
1:F:264:HIS:HE1	2:F:452:HOH:O	1.98	0.46
1:D:140:GLN:NE2	2:D:629:HOH:O	2.48	0.46
1:B:264:HIS:HE1	2:B:304:HOH:O	1.98	0.45
1:C:263:HIS:HE1	2:C:361:HOH:O	1.99	0.45
1:D:35:LYS:HD2	1:D:76:ASP:HB2	1.98	0.45
1:B:262:HIS:O	1:B:265:HIS:HD2	2.00	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:39:GLY:N	1:D:94[A]:ILE:HG21	2.31	0.45
1:C:160:SER:C	1:C:161:GLU:HG2	2.37	0.44
1:C:199:PHE:CZ	1:F:171:LEU:HD11	2.52	0.43
1:B:92:HIS:HE1	1:B:225:ASP:OD1	2.00	0.43
1:B:263:HIS:HE1	2:B:487:HOH:O	2.00	0.43
1:B:39:GLY:N	1:B:94[A]:ILE:HG21	2.35	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	233/242 (96%)	228 (98%)	5 (2%)	0	100	100
1	C	233/242 (96%)	230 (99%)	3 (1%)	0	100	100
1	D	234/242 (97%)	231 (99%)	3 (1%)	0	100	100
1	F	231/242 (96%)	227 (98%)	4 (2%)	0	100	100
All	All	931/968 (96%)	916 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	205/210 (98%)	202 (98%)	3 (2%)	65	61
1	C	205/210 (98%)	201 (98%)	4 (2%)	55	49
1	D	206/210 (98%)	203 (98%)	3 (2%)	65	61
1	F	203/210 (97%)	199 (98%)	4 (2%)	55	49
All	All	819/840 (98%)	805 (98%)	14 (2%)	62	55

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	34	GLU
1	B	109	LYS
1	B	142	ARG
1	C	34	GLU
1	C	93	ARG
1	C	165[A]	ARG
1	C	165[B]	ARG
1	D	34	GLU
1	D	93	ARG
1	D	133	LYS
1	F	34	GLU
1	F	93	ARG
1	F	142	ARG
1	F	256	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (32) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	41	ASN
1	B	88	ASN
1	B	92	HIS
1	B	140	GLN
1	B	257	ASN
1	B	263	HIS
1	B	264	HIS
1	B	265	HIS
1	C	41	ASN
1	C	88	ASN
1	C	92	HIS
1	C	140	GLN
1	C	257	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	263	HIS
1	C	264	HIS
1	C	265	HIS
1	D	88	ASN
1	D	92	HIS
1	D	134	HIS
1	D	140	GLN
1	D	257	ASN
1	D	263	HIS
1	D	264	HIS
1	D	265	HIS
1	F	41	ASN
1	F	88	ASN
1	F	140	GLN
1	F	256	ASN
1	F	257	ASN
1	F	263	HIS
1	F	264	HIS
1	F	265	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	232/242 (95%)	0.39	13 (5%) 24 27	2, 2, 8, 19	0
1	C	232/242 (95%)	0.29	13 (5%) 24 27	2, 2, 6, 16	0
1	D	232/242 (95%)	0.34	10 (4%) 35 38	2, 2, 7, 16	0
1	F	232/242 (95%)	0.31	9 (3%) 39 42	2, 2, 6, 15	0
All	All	928/968 (95%)	0.33	45 (4%) 30 34	2, 2, 7, 19	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	74	THR	3.5
1	D	34	GLU	3.3
1	B	175	MET	3.0
1	F	175	MET	2.9
1	D	134	HIS	2.8
1	C	133	LYS	2.7
1	B	72	GLY	2.6
1	B	131	GLU	2.6
1	B	133	LYS	2.5
1	B	256	ASN	2.5
1	B	35	LYS	2.5
1	D	175	MET	2.5
1	B	94[A]	ILE	2.4
1	B	34	GLU	2.4
1	C	178	MET	2.4
1	D	178	MET	2.4
1	F	34	GLU	2.4
1	F	178	MET	2.4
1	D	131	GLU	2.4
1	C	34	GLU	2.3
1	C	175	MET	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	74	THR	2.3
1	C	146	VAL	2.3
1	D	173	VAL	2.3
1	B	178	MET	2.2
1	F	173	VAL	2.2
1	C	171	LEU	2.2
1	B	173	VAL	2.2
1	B	87	LEU	2.2
1	F	35	LYS	2.2
1	C	87	LEU	2.1
1	C	177	VAL	2.1
1	F	131	GLU	2.1
1	F	177	VAL	2.1
1	C	93	ARG	2.1
1	B	242	LYS	2.1
1	C	165[A]	ARG	2.1
1	D	256	ASN	2.1
1	F	74	THR	2.1
1	C	131	GLU	2.0
1	C	173	VAL	2.0
1	D	177	VAL	2.0
1	F	163	HIS	2.0
1	D	93	ARG	2.0
1	D	160	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.