



Full wwPDB X-ray Structure Validation Report ⓘ

May 24, 2021 – 06:06 PM EDT

PDB ID : 2DFP
Title : X-RAY STRUCTURE OF AGED DI-ISOPROPYL-PHOSPHORO-FLUORIDE (DFP) BOUND TO ACETYLCHOLINESTERASE
Authors : Kryger, G.; Millard, C.B.; Silman, I.; Sussman, J.L.
Deposited on : 1998-12-07
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.18
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.18


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	534	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 19%, green 77%);"></div> <div style="display: flex; justify-content: space-between; font-size: 0.8em;"> 1% 77% 19% ... </div> </div>
2	B	2	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 50%, yellow 50%);"></div> <div style="display: flex; justify-content: space-between; font-size: 0.8em;"> 50% 50% </div> </div>
2	C	2	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, yellow 100%);"></div> <div style="text-align: center; font-size: 0.8em;">100%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	C	2	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4728 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (ACETYLCHOLINESTERASE).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	534	Total	C	N	O	P	S	0	0	0
			4270	2734	724	789	1	22			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	376	Total 376	O 376	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PROTEIN (ACETYLCHOLINESTERASE)



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	112.64Å 112.64Å 136.88Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.30 19.63 – 2.20	Depositor EDS
% Data completeness (in resolution range)	93.7 (20.00-2.30) 87.5 (19.63-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.60 (at 2.21Å)	Xtriage
Refinement program	CNS 0.4	Depositor
R, R_{free}	0.186 , 0.228 0.180 , 0.220	Depositor DCC
R_{free} test set	2015 reflections (4.49%)	wwPDB-VP
Wilson B-factor (Å ²)	38.0	Xtriage
Anisotropy	0.563	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 46.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.027 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4728	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, MES, MIS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.92	2/4380 (0.0%)	0.96	16/5944 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	228	SER	CB-OG	-6.02	1.34	1.42
1	A	145	SER	CB-OG	-5.32	1.35	1.42

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	426	ARG	NE-CZ-NH2	-9.54	115.53	120.30
1	A	221	ARG	NE-CZ-NH2	-8.42	116.09	120.30
1	A	531	LEU	CA-CB-CG	7.96	133.62	115.30
1	A	426	ARG	NE-CZ-NH1	7.72	124.16	120.30
1	A	216	ARG	NE-CZ-NH1	6.48	123.54	120.30
1	A	216	ARG	NE-CZ-NH2	-6.35	117.12	120.30
1	A	129	VAL	CB-CA-C	-6.21	99.60	111.40
1	A	494	LEU	CA-CB-CG	6.06	129.24	115.30
1	A	221	ARG	NE-CZ-NH1	6.04	123.32	120.30
1	A	518	VAL	CB-CA-C	-5.89	100.20	111.40
1	A	244	ARG	NE-CZ-NH2	-5.76	117.42	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	379	MET	CG-SD-CE	5.74	109.39	100.20
1	A	308	MET	CG-SD-CE	5.48	108.96	100.20
1	A	228	SER	CB-CA-C	-5.29	100.06	110.10
1	A	388	ARG	NE-CZ-NH1	5.15	122.88	120.30
1	A	47	ARG	NE-CZ-NH1	5.05	122.82	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	442	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4270	0	4111	69	0
2	B	28	0	25	0	0
2	C	28	0	25	0	0
3	A	14	0	13	1	0
4	A	12	0	13	0	0
5	A	376	0	0	9	0
All	All	4728	0	4187	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:22:VAL:HG13	1:A:133:LYS:HG3	1.63	0.79
1:A:468:ARG:HD3	5:A:2327:HOH:O	1.84	0.78
1:A:487:SER:OG	1:A:489:GLU:HG2	1.90	0.71
1:A:485:PRO:HA	5:A:2363:HOH:O	1.92	0.70
1:A:366:LEU:HD23	1:A:535:THR:HG21	1.77	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:223:ILE:HA	1:A:320:LEU:O	1.98	0.64
1:A:275:ILE:HA	1:A:278:GLU:OE1	1.99	0.63
1:A:345:SER:O	1:A:388:ARG:HG3	2.01	0.60
1:A:426:ARG:HD3	5:A:2085:HOH:O	2.02	0.60
1:A:5:GLU:OE2	1:A:104:PRO:HA	2.02	0.59
1:A:419:TYR:CZ	1:A:494:LEU:HD13	2.37	0.59
1:A:452:LEU:HD13	1:A:467:ARG:NH2	2.18	0.58
1:A:515:ARG:O	1:A:518:VAL:HG22	2.05	0.57
1:A:197:PHE:CB	1:A:223:ILE:HB	2.34	0.57
1:A:451:PRO:HA	1:A:458:TYR:CD2	2.41	0.55
1:A:63:TYR:CD1	1:A:126:THR:HG22	2.43	0.54
1:A:461:GLU:H	1:A:461:GLU:CD	2.10	0.54
1:A:366:LEU:CD2	1:A:535:THR:HG21	2.38	0.54
1:A:127:LEU:HD12	1:A:130:TYR:CE2	2.44	0.53
1:A:349:ARG:HD3	1:A:381:ASP:O	2.09	0.52
1:A:452:LEU:HD22	1:A:463:GLU:HG3	1.91	0.52
1:A:200:MIS:H33	1:A:233:TRP:CE2	2.45	0.52
1:A:23:LEU:HD11	1:A:452:LEU:HD12	1.93	0.50
1:A:204:ALA:O	1:A:208:MET:HG3	2.11	0.50
1:A:216:ARG:HB3	1:A:315:LYS:HB2	1.93	0.49
1:A:212:SER:O	1:A:216:ARG:HG2	2.13	0.49
1:A:405:MET:HA	1:A:408:VAL:HG12	1.94	0.48
1:A:132:GLY:HA3	1:A:143:LEU:HD22	1.94	0.48
1:A:244:ARG:HD3	1:A:291:SER:HB3	1.95	0.48
1:A:197:PHE:HB3	1:A:223:ILE:HB	1.95	0.48
1:A:88:ARG:CD	5:A:2264:HOH:O	2.62	0.47
1:A:197:PHE:HB2	1:A:223:ILE:HB	1.96	0.47
1:A:528:LEU:HB3	1:A:529:PRO:HD3	1.95	0.47
1:A:19:ARG:NH2	1:A:26:HIS:HB2	2.30	0.47
1:A:293:VAL:HB	1:A:294:PRO:HD2	1.96	0.47
1:A:454:LYS:HG3	1:A:455:GLU:N	2.30	0.47
1:A:484:GLU:HB2	1:A:487:SER:HB3	1.97	0.46
1:A:24:SER:HA	5:A:2179:HOH:O	2.16	0.46
1:A:199:GLU:HA	1:A:225:GLN:O	2.16	0.45
1:A:211:LEU:HD23	1:A:314:PHE:HB3	1.99	0.45
1:A:452:LEU:HD13	1:A:467:ARG:CZ	2.45	0.45
1:A:465:LEU:HD13	1:A:506:ASN:O	2.17	0.45
1:A:88:ARG:HD2	5:A:2264:HOH:O	2.16	0.45
1:A:481:ASN:OD1	1:A:482:PRO:HD2	2.17	0.45
1:A:22:VAL:O	1:A:23:LEU:C	2.56	0.44
1:A:487:SER:OG	1:A:488:GLN:N	2.51	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:450:LEU:HD23	1:A:450:LEU:N	2.33	0.43
1:A:508:GLU:HB3	1:A:509:PRO:HD2	1.99	0.43
1:A:82:GLU:HG2	5:A:2204:HOH:O	2.18	0.43
1:A:265:CYS:O	1:A:268:GLU:HB2	2.18	0.43
1:A:135:LEU:HD23	1:A:143:LEU:CD1	2.49	0.43
1:A:357:LYS:HG2	1:A:368:LEU:HD21	2.01	0.43
1:A:228:SER:HB3	1:A:230:ASN:OD1	2.19	0.43
1:A:439:ILE:HG22	1:A:440:HIS:O	2.19	0.43
1:A:206:VAL:CG1	1:A:222:ALA:HB1	2.49	0.42
1:A:353:MET:HG3	5:A:2106:HOH:O	2.18	0.42
1:A:216:ARG:HG2	1:A:216:ARG:H	1.49	0.42
1:A:286:SER:OG	1:A:288:PHE:O	2.38	0.41
1:A:349:ARG:HA	1:A:349:ARG:HD2	1.89	0.41
1:A:350:GLU:OE1	1:A:350:GLU:HA	2.20	0.41
1:A:508:GLU:HB3	1:A:509:PRO:CD	2.51	0.41
1:A:19:ARG:CZ	1:A:26:HIS:HB2	2.50	0.41
1:A:146:LEU:C	1:A:146:LEU:HD12	2.40	0.41
3:A:3001:NAG:O6	5:A:2368:HOH:O	2.22	0.41
1:A:105:ARG:HA	1:A:106:PRO:HD3	1.92	0.41
1:A:502:PHE:CZ	1:A:513:HIS:HB2	2.55	0.41
1:A:221:ARG:HD3	1:A:318:GLN:OE1	2.21	0.41
1:A:484:GLU:HB2	1:A:487:SER:CB	2.51	0.41
1:A:135:LEU:HD23	1:A:143:LEU:HD11	2.03	0.40
1:A:207:GLY:HA3	1:A:229:PRO:HD3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	531/534 (99%)	508 (96%)	21 (4%)	2 (0%)	34 42

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	3	HIS
1	A	485	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	466/466 (100%)	436 (94%)	30 (6%)	17 23

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	SER
1	A	24	SER
1	A	38	PRO
1	A	44	ARG
1	A	89	GLU
1	A	129	VAL
1	A	140	GLU
1	A	143	LEU
1	A	145	SER
1	A	197	PHE
1	A	216	ARG
1	A	228	SER
1	A	246	VAL
1	A	281	VAL
1	A	295	VAL
1	A	330	PHE
1	A	337	PRO
1	A	356	VAL
1	A	357	LYS
1	A	358	LEU
1	A	365	ASP
1	A	368	LEU
1	A	430	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	450	LEU
1	A	454	LYS
1	A	473	TRP
1	A	499	GLU
1	A	516	LEU
1	A	518	VAL
1	A	531	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	68	GLN
1	A	526	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	MIS	A	200	1	11,12,13	1.83	2 (18%)	10,16,18	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MIS	A	200	1	-	4/10/13/15	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	200	MIS	P-O3P	-4.99	1.47	1.60
1	A	200	MIS	P-OG	-2.70	1.48	1.59

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	200	MIS	N-CA-CB-OG
1	A	200	MIS	CB-OG-P-O3P
1	A	200	MIS	C1-O3P-P-OG
1	A	200	MIS	CB-OG-P-O2P

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	200	MIS	1	0

5.5 Carbohydrates

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	1,2	14,14,15	0.75	0	17,19,21	0.87	0
2	NAG	B	2	2	14,14,15	1.42	1 (7%)	17,19,21	1.19	1 (5%)
2	NAG	C	1	1,2	14,14,15	1.45	3 (21%)	17,19,21	0.90	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	2	2	14,14,15	1.16	1 (7%)	17,19,21	0.95	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1
2	NAG	C	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	C	2	2	-	1/6/23/26	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	NAG	C4-C5	3.12	1.59	1.53
2	C	1	NAG	C1-C2	2.83	1.56	1.52
2	C	1	NAG	O5-C5	2.24	1.48	1.43
2	C	1	NAG	O5-C1	2.23	1.47	1.43
2	C	2	NAG	C1-C2	2.01	1.55	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	NAG	C2-N2-C7	-4.21	116.91	122.90
2	C	1	NAG	C2-N2-C7	-2.33	119.59	122.90

There are no chirality outliers.


All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	2	NAG	O5-C5-C6-O6
2	B	1	NAG	O5-C5-C6-O6
2	B	1	NAG	C4-C5-C6-O6
2	C	1	NAG	C4-C5-C6-O6
2	B	2	NAG	C4-C5-C6-O6
2	C	1	NAG	O5-C5-C6-O6
2	C	2	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	MES	A	4001	-	12,12,12	8.30	7 (58%)	14,16,16	2.64	6 (42%)
3	NAG	A	3001	1	14,14,15	1.13	1 (7%)	17,19,21	1.01	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MES	A	4001	-	-	3/6/14/14	0/1/1/1
3	NAG	A	3001	1	-	2/6/23/26	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	4001	MES	C8-S	-21.48	1.47	1.77
4	A	4001	MES	O1S-S	11.50	1.79	1.45
4	A	4001	MES	O2S-S	10.50	1.76	1.45
4	A	4001	MES	O3S-S	8.79	1.78	1.47
4	A	4001	MES	C7-C8	-4.25	1.41	1.52
4	A	4001	MES	C7-N4	-2.82	1.40	1.47
4	A	4001	MES	C3-C2	-2.45	1.40	1.50
3	A	3001	NAG	O5-C5	2.07	1.47	1.43

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	4001	MES	O3S-S-C8	6.33	116.01	105.77
4	A	4001	MES	O1S-S-C8	4.79	112.69	106.92
4	A	4001	MES	O3S-S-O2S	-2.98	103.99	111.27
4	A	4001	MES	O2S-S-C8	2.56	110.00	106.92
4	A	4001	MES	O1-C2-C3	-2.44	106.43	111.80
3	A	3001	NAG	C2-N2-C7	-2.34	119.56	122.90
4	A	4001	MES	O3S-S-O1S	-2.34	105.55	111.27
3	A	3001	NAG	O5-C1-C2	-2.26	107.71	111.29

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	4001	MES	C7-C8-S-O2S
3	A	3001	NAG	O5-C5-C6-O6
3	A	3001	NAG	C4-C5-C6-O6
4	A	4001	MES	C7-C8-S-O3S
4	A	4001	MES	C7-C8-S-O1S

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	3001	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	533/534 (99%)	-0.47	7 (1%) 77 81	25, 39, 60, 91	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	ASP	6.9
1	A	3	HIS	6.4
1	A	486	HIS	4.5
1	A	488	GLN	4.0
1	A	485	PRO	3.4
1	A	55	SER	2.8
1	A	156	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	MIS	A	200	13/14	0.99	0.07	26,28,33,34	0


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	C	2	14/15	0.77	0.49	67,73,75,76	0
2	NAG	B	2	14/15	0.85	0.44	61,69,73,75	0
2	NAG	C	1	14/15	0.90	0.36	70,72,73,73	0
2	NAG	B	1	14/15	0.92	0.24	45,51,58,63	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NAG	A	3001	14/15	0.87	0.27	67,69,72,74	0
4	MES	A	4001	12/12	0.95	0.16	64,66,67,67	0

6.5 Other polymers [i](#)

There are no such residues in this entry.