



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 23, 2020 – 02:09 PM JST

PDB ID : 7DFM
Title : Crystal structure of glycoside hydrolase family 11 beta-xylanase from *Streptomyces olivaceoviridis* E-86
Authors : Fujimoto, Z.; Kishine, N.; Kaneko, S.
Deposited on : 2020-11-09
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.16
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.16


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	205	<div> <div> <div></div> <div>83%</div> <div>11%</div> <div>5%</div> </div> </div>
1	B	205	<div> <div> <div></div> <div>83%</div> <div>11%</div> <div>6%</div> </div> </div>
1	C	205	<div> <div> <div></div> <div>80%</div> <div>14%</div> <div>6%</div> </div> </div>
1	D	205	<div> <div> <div></div> <div>77%</div> <div>17%</div> <div>5%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6144 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endo-1,4-beta-xylanase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	194	Total	C	N	O	S	0	0	0
			1499	942	253	300	4			
1	B	193	Total	C	N	O	S	0	0	0
			1494	939	252	299	4			
1	C	193	Total	C	N	O	S	0	0	0
			1494	939	252	299	4			
1	D	194	Total	C	N	O	S	0	0	0
			1499	942	253	300	4			

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	4	Total	Cl	0	0
			4	4		
2	A	7	Total	Cl	0	0
			7	7		
2	D	5	Total	Cl	0	0
			5	5		
2	C	6	Total	Cl	0	0
			6	6		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	37	Total	O	0	0
			37	37		
3	B	37	Total	O	0	0
			37	37		
3	C	28	Total	O	0	0
			28	28		
3	D	34	Total	O	0	0
			34	34		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Endo-1,4-beta-xylanase




- Molecule 1: Endo-1,4-beta-xylanase



- Molecule 1: Endo-1,4-beta-xylanase



- Molecule 1: Endo-1,4-beta-xylanase



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	135.04Å 135.04Å 72.34Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.41 – 2.40 49.36 – 2.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.41-2.40) 100.0 (49.36-2.40)	Depositor EDS
R_{merge}	0.23	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.88 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.168 , 0.245 0.175 , 0.246	Depositor DCC
R_{free} test set	1571 reflections (5.24%)	wwPDB-VP
Wilson B-factor (Å ²)	29.2	Xtriage
Anisotropy	0.011	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 36.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.019 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6144	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.81	0/1543	1.04	1/2102 (0.0%)
1	B	0.77	0/1538	1.05	2/2095 (0.1%)
1	C	0.80	0/1538	1.07	1/2095 (0.0%)
1	D	0.81	0/1543	1.06	1/2102 (0.0%)
All	All	0.80	0/6162	1.06	5/8394 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	140	ARG	NE-CZ-NH2	-7.85	116.38	120.30
1	B	140	ARG	NE-CZ-NH1	6.16	123.38	120.30
1	A	59	ARG	NE-CZ-NH1	-5.97	117.31	120.30
1	D	140	ARG	NE-CZ-NH2	-5.25	117.67	120.30
1	B	140	ARG	NE-CZ-NH2	-5.03	117.79	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1499	0	1371	12	0
1	B	1494	0	1366	10	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1494	0	1366	15	0
1	D	1499	0	1371	21	0
2	A	7	0	0	0	0
2	B	4	0	0	0	0
2	C	6	0	0	0	0
2	D	5	0	0	0	0
3	A	37	0	0	0	0
3	B	37	0	0	0	0
3	C	28	0	0	0	0
3	D	34	0	0	0	0
All	All	6144	0	5474	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:29:MET:HE2	1:C:37:TYR:CD2	2.22	0.73
1:B:63:TYR:HA	1:B:188:THR:O	1.96	0.65
1:A:68:TYR:CD1	1:D:119:THR:HB	2.33	0.64
1:A:6:THR:O	1:A:8:GLN:HG2	1.99	0.63
1:C:79:GLY:HA3	1:C:88:TYR:CZ	2.35	0.61
1:C:80:TRP:HA	1:C:86:VAL:O	1.99	0.60
1:B:5:THR:H	1:C:55:ASN:ND2	1.98	0.60
1:C:160:ARG:HD2	1:D:126:SER:O	2.02	0.58
1:D:34:GLY:O	1:D:51:LYS:NZ	2.35	0.58
1:C:108:THR:O	1:D:128:GLU:OE2	2.25	0.55
1:C:29:MET:CE	1:C:37:TYR:CD2	2.93	0.51
1:B:77:LEU:HD13	1:B:77:LEU:C	2.32	0.49
1:B:61:VAL:O	1:B:148:THR:HA	2.14	0.48
1:B:72:ASN:HD22	1:B:179:TYR:HB3	1.78	0.48
1:D:62:GLN:NE2	1:D:148:THR:OG1	2.44	0.47
1:C:55:ASN:HA	1:C:170:TYR:O	2.15	0.47
1:A:29:MET:HE2	1:A:37:TYR:HB2	1.97	0.47
1:B:80:TRP:CZ3	1:B:87:GLU:HB2	2.51	0.46
1:D:94:TRP:CE2	1:D:143:LYS:HE2	2.51	0.46
1:D:87:GLU:O	1:D:135:GLN:HA	2.15	0.46
1:C:87:GLU:O	1:C:135:GLN:HA	2.15	0.46
1:A:68:TYR:CE1	1:D:119:THR:HB	2.51	0.46
1:C:82:SER:O	1:C:166:GLY:HA3	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:80:TRP:HA	1:B:86:VAL:O	2.16	0.46
1:C:160:ARG:CD	1:D:126:SER:O	2.65	0.45
1:D:94:TRP:NE1	1:D:143:LYS:HE2	2.31	0.45
1:B:87:GLU:O	1:B:135:GLN:HA	2.16	0.45
1:B:46:ASN:HA	1:B:178:GLY:O	2.17	0.45
1:C:48:VAL:HG11	1:C:78:TYR:CE2	2.51	0.45
1:D:66:SER:HB2	1:D:68:TYR:CE2	2.51	0.45
1:A:61:VAL:O	1:A:148:THR:HA	2.18	0.44
1:A:186:ASN:HD21	1:D:72:ASN:ND2	2.14	0.44
1:B:157:ALA:O	1:B:158:TRP:C	2.56	0.44
1:D:80:TRP:NE1	1:D:127:VAL:HG13	2.33	0.44
1:A:73:GLY:O	1:A:94:TRP:HA	2.18	0.43
1:A:84:PRO:HD2	1:A:166:GLY:CA	2.48	0.43
1:A:77:LEU:HD11	1:A:172:MET:SD	2.58	0.43
1:D:73:GLY:O	1:D:94:TRP:HA	2.18	0.43
1:C:25:GLY:HA3	1:C:43:ASN:O	2.19	0.43
1:D:93:ASN:HB2	1:D:144:ARG:O	2.19	0.42
1:D:65:GLY:HA3	1:D:186:ASN:O	2.19	0.42
1:D:32:ASN:HB3	1:D:36:ASN:HB3	2.01	0.42
1:D:11:THR:HA	1:D:15:PHE:O	2.20	0.42
1:D:157:ALA:HA	1:D:160:ARG:HD3	2.00	0.42
1:D:18:SER:O	1:D:49:ALA:HA	2.19	0.42
1:C:63:TYR:HA	1:C:188:THR:O	2.19	0.41
1:A:35:GLY:HA3	1:A:189:VAL:O	2.21	0.41
1:A:18:SER:O	1:A:49:ALA:HA	2.21	0.41
1:C:1:ALA:HA	1:C:31:LEU:O	2.21	0.40
1:A:186:ASN:HD21	1:D:72:ASN:HD21	1.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	192/205 (94%)	187 (97%)	5 (3%)	0	100	100
1	B	191/205 (93%)	186 (97%)	5 (3%)	0	100	100
1	C	191/205 (93%)	183 (96%)	8 (4%)	0	100	100
1	D	192/205 (94%)	188 (98%)	4 (2%)	0	100	100
All	All	766/820 (93%)	744 (97%)	22 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	156/165 (94%)	152 (97%)	4 (3%)	46	66
1	B	156/165 (94%)	150 (96%)	6 (4%)	33	51
1	C	156/165 (94%)	151 (97%)	5 (3%)	39	59
1	D	156/165 (94%)	148 (95%)	8 (5%)	24	39
All	All	624/660 (94%)	601 (96%)	23 (4%)	34	53

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	62	GLN
1	A	83	ASN
1	A	126	SER
1	A	192	LYS
1	B	26	SER
1	B	33	SER
1	B	142	SER
1	B	169	SER
1	B	184	SER
1	B	192	LYS
1	C	40	SER
1	C	54	SER
1	C	167	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	174	LEU
1	C	192	LYS
1	D	32	ASN
1	D	33	SER
1	D	108	THR
1	D	121	ARG
1	D	126	SER
1	D	167	SER
1	D	177	GLU
1	D	184	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	118	GLN
1	A	180	GLN
1	C	46	ASN
1	C	55	ASN
1	C	141	GLN
1	D	60	ASN
1	D	62	GLN
1	D	72	ASN
1	D	141	GLN
1	D	180	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 22 ligands modelled in this entry, 22 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	194/205 (94%)	-0.53	2 (1%) 82 80	18, 26, 43, 71	0
1	B	193/205 (94%)	-0.36	3 (1%) 72 70	20, 28, 42, 84	0
1	C	193/205 (94%)	-0.51	0 100 100	17, 27, 43, 71	0
1	D	194/205 (94%)	-0.36	4 (2%) 63 61	22, 29, 42, 84	0
All	All	774/820 (94%)	-0.44	9 (1%) 79 77	17, 28, 43, 84	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	192	LYS	4.1
1	D	191	GLY	3.9
1	D	192	LYS	3.8
1	D	194	ALA	3.5
1	B	191	GLY	3.0
1	A	193	LEU	2.8
1	B	193	LEU	2.6
1	D	193	LEU	2.6
1	A	192	LYS	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CL	D	304	1/1	0.93	0.08	41,41,41,41	0
2	CL	B	401	1/1	0.96	0.12	34,34,34,34	0
2	CL	A	404	1/1	0.96	0.07	39,39,39,39	0
2	CL	B	403	1/1	0.97	0.07	33,33,33,33	0
2	CL	C	404	1/1	0.97	0.07	42,42,42,42	0
2	CL	C	406	1/1	0.97	0.08	38,38,38,38	0
2	CL	D	301	1/1	0.97	0.11	39,39,39,39	0
2	CL	A	405	1/1	0.97	0.25	44,44,44,44	0
2	CL	C	403	1/1	0.97	0.09	43,43,43,43	0
2	CL	D	305	1/1	0.98	0.09	41,41,41,41	0
2	CL	A	407	1/1	0.98	0.11	29,29,29,29	0
2	CL	D	302	1/1	0.98	0.16	39,39,39,39	0
2	CL	A	402	1/1	0.98	0.06	25,25,25,25	0
2	CL	B	404	1/1	0.98	0.14	38,38,38,38	0
2	CL	C	401	1/1	0.99	0.06	29,29,29,29	0
2	CL	C	402	1/1	0.99	0.08	29,29,29,29	0
2	CL	D	303	1/1	0.99	0.20	42,42,42,42	0
2	CL	A	401	1/1	0.99	0.09	26,26,26,26	0
2	CL	A	406	1/1	0.99	0.03	28,28,28,28	0
2	CL	C	405	1/1	0.99	0.04	36,36,36,36	0
2	CL	B	402	1/1	1.00	0.05	32,32,32,32	0
2	CL	A	403	1/1	1.00	0.07	35,35,35,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.