



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 1, 2021 – 04:58 PM EDT

PDB ID : 3DGS  
Title : Changing the determinants of protein stability from covalent to non-covalent interactions by in-vitro evolution: a structural and energetic analysis  
Authors : Jakob, R.P.; Kather, I.; Dobbek, H.; Schmid, F.X.  
Deposited on : 2008-06-16  
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.23.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	227	
1	B	227	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3213 atoms, of which 2 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Coat protein A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	186	Total	C	H	N	O	S	4	0	0
			1483	955	1	238	287	2			
1	B	186	Total	C	H	N	O	S	15	0	0
			1472	947	1	237	285	2			

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP P03661
A	7	SER	CYS	engineered mutation	UNP P03661
A	11	SER	PRO	engineered mutation	UNP P03661
A	13	ILE	THR	engineered mutation	UNP P03661
A	15	GLY	ASN	engineered mutation	UNP P03661
A	29	TRP	ARG	engineered mutation	UNP P03661
A	36	ILE	CYS	engineered mutation	UNP P03661
A	39	LYS	ASN	engineered mutation	UNP P03661
A	46	ILE	CYS	engineered mutation	UNP P03661
A	53	VAL	CYS	engineered mutation	UNP P03661
A	55	ALA	GLY	engineered mutation	UNP P03661
A	56	ILE	THR	engineered mutation	UNP P03661
A	60	VAL	ILE	engineered mutation	UNP P03661
A	101	ILE	THR	engineered mutation	UNP P03661
A	129	HIS	GLN	engineered mutation	UNP P03661
A	138	GLY	ASN	engineered mutation	UNP P03661
A	188	VAL	CYS	engineered mutation	UNP P03661
A	199	LEU	PHE	engineered mutation	UNP P03661
A	201	ALA	CYS	engineered mutation	UNP P03661
A	207	LEU	SER	engineered mutation	UNP P03661
A	209	TYR	ASP	engineered mutation	UNP P03661
A	218	PRO	-	expression tag	UNP P03661
A	219	SER	-	expression tag	UNP P03661
A	220	GLY	-	expression tag	UNP P03661
A	221	HIS	-	expression tag	UNP P03661

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	222	HIS	-	expression tag	UNP P03661
A	223	HIS	-	expression tag	UNP P03661
A	224	HIS	-	expression tag	UNP P03661
A	225	HIS	-	expression tag	UNP P03661
A	226	HIS	-	expression tag	UNP P03661
B	0	MET	-	initiating methionine	UNP P03661
B	7	SER	CYS	engineered mutation	UNP P03661
B	11	SER	PRO	engineered mutation	UNP P03661
B	13	ILE	THR	engineered mutation	UNP P03661
B	15	GLY	ASN	engineered mutation	UNP P03661
B	29	TRP	ARG	engineered mutation	UNP P03661
B	36	ILE	CYS	engineered mutation	UNP P03661
B	39	LYS	ASN	engineered mutation	UNP P03661
B	46	ILE	CYS	engineered mutation	UNP P03661
B	53	VAL	CYS	engineered mutation	UNP P03661
B	55	ALA	GLY	engineered mutation	UNP P03661
B	56	ILE	THR	engineered mutation	UNP P03661
B	60	VAL	ILE	engineered mutation	UNP P03661
B	101	ILE	THR	engineered mutation	UNP P03661
B	129	HIS	GLN	engineered mutation	UNP P03661
B	138	GLY	ASN	engineered mutation	UNP P03661
B	188	VAL	CYS	engineered mutation	UNP P03661
B	199	LEU	PHE	engineered mutation	UNP P03661
B	201	ALA	CYS	engineered mutation	UNP P03661
B	207	LEU	SER	engineered mutation	UNP P03661
B	209	TYR	ASP	engineered mutation	UNP P03661
B	218	PRO	-	expression tag	UNP P03661
B	219	SER	-	expression tag	UNP P03661
B	220	GLY	-	expression tag	UNP P03661
B	221	HIS	-	expression tag	UNP P03661
B	222	HIS	-	expression tag	UNP P03661
B	223	HIS	-	expression tag	UNP P03661
B	224	HIS	-	expression tag	UNP P03661
B	225	HIS	-	expression tag	UNP P03661
B	226	HIS	-	expression tag	UNP P03661

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	151	Total O 151 151	0	0
2	B	107	Total O 107 107	0	0



- Molecule 1: Coat protein A



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.86Å 92.21Å 94.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.90 19.63 – 1.80	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-1.90) 100.0 (19.63-1.80)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.56 (at 1.80Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.216 , 0.263 0.222 , 0.268	Depositor DCC
$R_{free}$ test set	2012 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.4	Xtriage
Anisotropy	0.197	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 48.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.026 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3213	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.11% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.43	0/1533	0.63	0/2105
1	B	0.42	0/1520	0.65	0/2086
All	All	0.43	0/3053	0.64	0/4191

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1482	1	1383	38	1
1	B	1471	1	1381	42	1
2	A	151	0	0	5	0
2	B	107	0	0	4	0
All	All	3211	2	2764	80	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 80 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:132:ASN:OD1	1:B:193:GLY:N	1.97	0.97

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:18:THR:H	1:A:216:ASN:HD21	1.10	0.91
1:A:162:VAL:HG22	2:A:344:HOH:O	1.68	0.91
1:A:155:VAL:HG13	1:A:165:TYR:HB2	1.52	0.89
1:B:132:ASN:ND2	1:B:192:SER:HA	1.96	0.81

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:123:PRO:O	1:B:192:SER:OG[3_555]	2.14	0.06

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	182/227 (80%)	168 (92%)	12 (7%)	2 (1%)	14	5
1	B	182/227 (80%)	164 (90%)	12 (7%)	6 (3%)	4	0
All	All	364/454 (80%)	332 (91%)	24 (7%)	8 (2%)	6	1

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	161	PRO
1	B	6	SER
1	B	161	PRO
1	B	162	VAL
1	B	197	ASP



### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	159/185 (86%)	156 (98%)	3 (2%)	57	53
1	B	158/185 (85%)	157 (99%)	1 (1%)	86	87
All	All	317/370 (86%)	313 (99%)	4 (1%)	69	68

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	94	ASP
1	A	162	VAL
1	A	216	ASN
1	B	157	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	216	ASN
1	B	19	ASN
1	B	206	GLN
1	B	129	HIS
1	B	167	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	186/227 (81%)	-0.04	9 (4%) 30 33	14, 23, 62, 124	1 (0%)
1	B	186/227 (81%)	0.48	23 (12%) 4 4	12, 26, 86, 136	6 (3%)
All	All	372/454 (81%)	0.22	32 (8%) 10 12	12, 24, 79, 136	7 (1%)

The worst 5 of 32 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	194	PHE	20.1
1	B	159	THR	11.2
1	B	195	ASN	8.3
1	B	161	PRO	7.1
1	A	159	THR	6.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.