



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 02:14 pm BST

PDB ID : 2DH2
Title : Crystal Structure of human ED-4F2hc
Authors : Fort, J.; Fita, I.; Palacin, M.
Deposited on : 2006-03-21
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	424	<div> <div>11%</div> <div>84%</div> <div>14%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ACT	A	601	-	-	X	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3412 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 4F2 cell-surface antigen heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	421	3269	2077	557	630	1	4	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	106	ASP	-	EXPRESSION TAG	UNP P08195
A	107	ARG	-	EXPRESSION TAG	UNP P08195
A	108	TRP	-	EXPRESSION TAG	UNP P08195
A	109	GLY	-	EXPRESSION TAG	UNP P08195
A	110	SER	-	EXPRESSION TAG	UNP P08195

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	4	2	2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	139	Total 139	O 139	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 4F2 cell-surface antigen heavy chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	39.67Å 67.85Å 73.95Å 90.00° 98.49° 90.00°	Depositor
Resolution (Å)	17.83 – 2.10 19.80 – 1.99	Depositor EDS
% Data completeness (in resolution range)	97.3 (17.83-2.10) 97.7 (19.80-1.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.28 (at 1.99Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.175 , 0.230 0.189 , 0.234	Depositor DCC
R_{free} test set	1157 reflections (4.47%)	wwPDB-VP
Wilson B-factor (Å ²)	29.4	Xtriage
Anisotropy	0.072	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 55.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3412	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	0/3338	0.73	3/4520 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	496	LEU	CA-CB-CG	6.72	130.76	115.30
1	A	350	ARG	NE-CZ-NH2	-6.32	117.14	120.30
1	A	308	GLY	N-CA-C	-5.09	100.38	113.10

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	220	THR	Peptide
1	A	221	GLN	Peptide
1	A	307	SER	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3269	0	3226	29	0
2	A	4	0	3	3	0
3	A	139	0	0	0	0
All	All	3412	0	3229	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:151:SER:HA	2:A:601:ACT:H1	1.68	0.76
1:A:155:VAL:O	2:A:601:ACT:H3	1.87	0.74
1:A:453:ILE:HD11	1:A:488:LEU:CD1	2.22	0.68
1:A:250:GLU:OE1	1:A:280:ASN:ND2	2.24	0.67
1:A:453:ILE:HD11	1:A:488:LEU:HD11	1.78	0.65
1:A:321:TYR:HE2	1:A:330:CYS:HG	1.45	0.62
1:A:449:LEU:CD1	1:A:475:ALA:HA	2.30	0.61
1:A:339:LEU:HD11	1:A:386:PRO:HG3	1.86	0.57
1:A:456:TRP:HD1	1:A:459:ASN:HD22	1.50	0.57
1:A:528:ALA:O	1:A:529:ALA:C	2.42	0.57
1:A:382:PRO:HB2	1:A:385:ALA:H	1.72	0.54
1:A:479:ALA:HB1	1:A:482:LEU:HD22	1.92	0.52
1:A:193:LEU:O	1:A:197:LYS:HG2	2.10	0.51
1:A:281:SER:OG	1:A:286:GLN:NE2	2.43	0.50
1:A:164:HIS:CD2	1:A:176:LEU:HD22	2.47	0.50
1:A:449:LEU:HD11	1:A:475:ALA:HA	1.96	0.47
1:A:302:SER:O	1:A:305:SER:HB3	2.14	0.47
1:A:130:GLN:HE21	1:A:130:GLN:H	1.63	0.46
1:A:449:LEU:HD23	1:A:467:ASN:HA	1.96	0.46
1:A:405:ASN:ND2	1:A:405:ASN:H	2.14	0.45
1:A:129:LEU:HD21	1:A:160:LEU:HD11	1.98	0.45
1:A:173:GLN:HA	1:A:173:GLN:HE21	1.82	0.45
1:A:321:TYR:HE2	1:A:330:CYS:SG	2.40	0.44
1:A:284:LEU:O	1:A:288:LEU:HG	2.15	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:150:LEU:HD13	1:A:158:LEU:HD21	1.99	0.43
1:A:154:LYS:HA	2:A:601:ACT:H2	1.99	0.43
1:A:479:ALA:CB	1:A:482:LEU:HD22	2.49	0.42
1:A:124:TYR:CZ	1:A:367:TYR:HA	2.55	0.41
1:A:325:THR:HB	1:A:328:ARG:HD3	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	419/424 (99%)	394 (94%)	22 (5%)	3 (1%)	22 18

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	306	ASP
1	A	479	ALA
1	A	380	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	354/353 (100%)	332 (94%)	22 (6%)	18 15

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	130	GLN
1	A	135	HIS
1	A	167	GLN
1	A	186	LYS
1	A	187	GLU
1	A	219	SER
1	A	221	GLN
1	A	223	ASP
1	A	273	ARG
1	A	290	LEU
1	A	296	ASP
1	A	305	SER
1	A	306	ASP
1	A	347	GLN
1	A	378	LEU
1	A	383	MSE
1	A	482	LEU
1	A	488	LEU
1	A	495	LEU
1	A	496	LEU
1	A	509	GLU
1	A	515	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	130	GLN
1	A	194	GLN
1	A	221	GLN
1	A	238	GLN
1	A	286	GLN
1	A	381	GLN
1	A	405	ASN
1	A	458	GLN
1	A	459	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACT	A	601	-	1,3,3	0.25	0	0,3,3	0.00	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	ACT	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	417/424 (98%)	0.60	47 (11%) 5 6	21, 32, 50, 63	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	327	ASN	9.1
1	A	109	GLY	8.2
1	A	110	SER	7.6
1	A	135	HIS	7.6
1	A	111	GLU	6.8
1	A	220	THR	5.8
1	A	482	LEU	5.6
1	A	136	GLY	5.2
1	A	308	GLY	4.9
1	A	112	LEU	4.6
1	A	529	ALA	4.4
1	A	114	ALA	4.3
1	A	480	SER	4.3
1	A	306	ASP	4.2
1	A	481	ASP	4.2
1	A	399	PRO	4.0
1	A	382	PRO	4.0
1	A	223	ASP	3.7
1	A	384	GLU	3.6
1	A	137	ALA	3.4
1	A	447	PRO	3.2
1	A	309	SER	3.2
1	A	484	ALA	3.2
1	A	292	GLU	3.1
1	A	307	SER	3.1
1	A	323	ASN	3.1
1	A	464	VAL	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	219	SER	2.8
1	A	295	LYS	2.8
1	A	380	GLY	2.7
1	A	371	ILE	2.7
1	A	479	ALA	2.6
1	A	486	ALA	2.6
1	A	465	VAL	2.6
1	A	113	PRO	2.6
1	A	214	GLU	2.6
1	A	446	GLY	2.5
1	A	483	PRO	2.5
1	A	134	GLY	2.5
1	A	501	GLY	2.5
1	A	478	GLN	2.4
1	A	221	GLN	2.4
1	A	521	LEU	2.3
1	A	522	LEU	2.3
1	A	385	ALA	2.3
1	A	463	LEU	2.3
1	A	523	LEU	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ACT	A	601	4/4	0.85	0.16	43,43,44,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.