



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 10:25 am BST

PDB ID : 4DJH
Title : Structure of the human kappa opioid receptor in complex with JDTic
Authors : Wu, H.; Wacker, D.; Katritch, V.; Mileni, M.; Han, G.W.; Vardy, E.; Liu, W.; Thompson, A.A.; Huang, X.P.; Carroll, F.I.; Mascarella, S.W.; Westkaemper, R.B.; Mosier, P.D.; Roth, B.L.; Cherezov, V.; Stevens, R.C.; GPCR Network (GPCR)
Deposited on : 2012-02-01
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	480	
1	B	480	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7089 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Kappa-type opioid receptor, Lysozyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	447	Total	C	N	O	S	0	0	0
			3480	2277	564	616	23			
1	B	448	Total	C	N	O	S	0	0	0
			3452	2263	555	611	23			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	38	GLY	-	EXPRESSION TAG	UNP P41145
A	39	GLY	-	EXPRESSION TAG	UNP P41145
A	40	THR	-	EXPRESSION TAG	UNP P41145
A	41	THR	-	EXPRESSION TAG	UNP P41145
A	42	MET	-	EXPRESSION TAG	UNP P41145
A	135	LEU	ILE	engineered mutation	UNP P41145
A	1054	THR	CYS	engineered mutation	UNP P00720
A	1097	ALA	CYS	engineered mutation	UNP P00720
B	38	GLY	-	EXPRESSION TAG	UNP P41145
B	39	GLY	-	EXPRESSION TAG	UNP P41145
B	40	THR	-	EXPRESSION TAG	UNP P41145
B	41	THR	-	EXPRESSION TAG	UNP P41145
B	42	MET	-	EXPRESSION TAG	UNP P41145
B	135	LEU	ILE	engineered mutation	UNP P41145
B	1054	THR	CYS	engineered mutation	UNP P00720
B	1097	ALA	CYS	engineered mutation	UNP P00720

- Molecule 2 is (3R)-7-hydroxy-N-{(2S)-1-[(3R,4R)-4-(3-hydroxyphenyl)-3,4-dimethylpiperidin-1-yl]-3-methylbutan-2-yl}-1,2,3,4-tetrahydroisoquinoline-3-carboxamide (three-letter code: JDC) (formula: C₂₈H₃₉N₃O₃).




- CIT
-
- The chemical structure of Citric acid (CIT) is shown. It consists of a central carbon atom (C3) bonded to three hydroxyl groups (OH) and a carboxyl group (COOH). The carboxyl group is further bonded to a methylene group (CH2), which is then bonded to another carboxyl group (COOH). The atoms are labeled with green text: C1, C2, C3, C4, C5, C6, C7, O1, O2, O3, O4, O5, O6, O7. The hydroxyl groups are shown in red, and the carboxyl groups are shown in red with double bonds to the oxygen atoms.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	6	7		


- Molecule 4 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLC)

(formula: C₂₁H₄₀O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			16	12	4		
4	B	1	Total	C	O	0	0
			21	17	4		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			7	4	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			7	4	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	11	Total	O	0	0
			11	11		
6	B	14	Total	O	0	0
			14	14		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Kappa-type opioid receptor, Lysozyme



- Molecule 1: Kappa-type opioid receptor, Lysozyme



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	54.90 Å 147.30 Å 205.29 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.86 – 2.90 35.86 – 2.86	Depositor EDS
% Data completeness (in resolution range)	97.8 (35.86-2.90) 96.5 (35.86-2.86)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.51 (at 2.85 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, R_{free}	0.227 , 0.265 0.223 , 0.258	Depositor DCC
R_{free} test set	1909 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	75.2	Xtriage
Anisotropy	0.213	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 57.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7089	wwPDB-VP
Average B, all atoms (Å ²)	84.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, OLC, JDC, CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.29	0/3555	0.42	0/4843
1	B	0.30	0/3528	0.45	0/4818
All	All	0.29	0/7083	0.44	0/9661

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3480	0	3491	49	0
1	B	3452	0	3430	45	0
2	A	34	0	38	0	0
2	B	34	0	38	2	0
3	A	13	0	5	0	0
4	B	37	0	50	2	0
5	B	14	0	20	0	0
6	A	11	0	0	0	0
6	B	14	0	0	0	0
All	All	7089	0	7072	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 7.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:124:TRP:CH2	1:B:134:VAL:HG11	2.23	0.73
1:A:1011:GLU:HG2	1:A:1030:GLY:HA3	1.76	0.67
1:B:138:ASP:OD1	2:B:1300:JDC:N2	2.31	0.64
1:A:122:ASN:ND2	1:A:122:ASN:O	2.32	0.63
1:A:161:CYS:SG	1:A:252:ARG:NH2	2.74	0.61
1:B:121:MET:HE1	1:B:125:PRO:HB3	1.83	0.60
1:B:138:ASP:OD2	2:B:1300:JDC:N2A	2.34	0.60
1:B:255:SER:O	1:B:258:LEU:CD2	2.50	0.60
1:A:139:TYR:HB3	1:A:189:VAL:CG2	2.34	0.57
1:A:112:MET:HE1	1:A:320:TYR:HB3	1.87	0.56
1:A:326:ASN:HB2	1:A:327:PRO:HD3	1.87	0.56
1:A:126:PHE:HB3	1:A:130:LEU:HD12	1.88	0.55
1:A:90:MET:HG3	1:A:90:MET:O	2.06	0.55
1:B:178:ILE:HD11	4:B:1302:OLC:H2	1.88	0.55
1:A:139:TYR:HB3	1:A:189:VAL:HG23	1.88	0.54
1:B:1095:ARG:NH1	1:B:1153:PHE:O	2.41	0.54
1:B:79:LEU:HD11	1:B:344:PHE:CD1	2.44	0.53
1:A:1085:LYS:N	1:A:1086:PRO:HD2	2.22	0.53
1:A:121:MET:HE1	1:A:125:PRO:HB3	1.91	0.52
1:B:115:GLN:OE1	1:B:316:ILE:CD1	2.57	0.52
1:B:112:MET:N	1:B:113:PRO:HD2	2.25	0.51
1:B:326:ASN:HB2	1:B:327:PRO:HD3	1.92	0.51
1:A:199:THR:HG22	1:A:208:ILE:CG2	2.41	0.50
1:B:157:TYR:N	1:B:249:MET:HE2	2.26	0.50
1:A:112:MET:N	1:A:113:PRO:HD2	2.26	0.50
1:B:112:MET:HG2	1:B:320:TYR:CE1	2.47	0.49
1:B:1048:LYS:O	1:B:1049:ALA:HB3	2.12	0.49
1:B:275:LEU:HD13	1:B:333:LEU:HB2	1.94	0.49
1:B:288:THR:N	1:B:289:PRO:CD	2.76	0.49
1:A:1008:ARG:NH2	1:A:1013:LEU:HD23	2.29	0.48
1:A:152:MET:HE2	1:A:276:VAL:HG22	1.95	0.48
1:B:119:TYR:CD2	1:B:119:TYR:C	2.86	0.48
1:A:257:ARG:HG2	1:A:258:LEU:HD12	1.95	0.48
1:B:1024:TYR:HB3	1:B:1032:LEU:HD11	1.97	0.47
1:B:266:ASP:OD1	1:B:266:ASP:N	2.47	0.47
1:A:92:THR:O	1:A:93:ALA:C	2.51	0.47
1:B:208:ILE:N	1:B:208:ILE:HD12	2.30	0.47
1:B:1011:GLU:CG	1:B:1030:GLY:HA3	2.45	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:124:TRP:CG	1:A:210:CYS:HB2	2.51	0.46
1:A:205:VAL:HB	1:A:207:VAL:HG13	1.98	0.46
1:A:199:THR:HG22	1:A:208:ILE:HG23	1.97	0.46
1:B:100:ASN:HB2	1:B:179:ASN:ND2	2.30	0.46
1:B:157:TYR:HE2	4:B:1303:OLC:H8A	1.81	0.46
1:A:1088:TYR:CZ	1:A:1096:ARG:HD3	2.51	0.45
1:A:274:ARG:O	1:A:278:VAL:HG23	2.16	0.45
1:A:171:THR:HB	1:A:172:PRO:HD3	1.98	0.45
1:A:58:ILE:HG22	1:A:59:PRO:N	2.32	0.45
1:B:1026:THR:HG22	1:B:1027:ILE:N	2.32	0.45
1:A:1119:ARG:HG3	1:A:1120:MET:N	2.32	0.45
1:A:156:ARG:C	1:A:249:MET:HE2	2.37	0.45
1:A:307:ALA:O	1:A:308:ALA:HB3	2.17	0.45
1:B:143:PHE:CD1	1:B:189:VAL:HG21	2.52	0.45
1:A:58:ILE:HB	1:A:59:PRO:HD3	1.98	0.45
1:B:58:ILE:HB	1:B:59:PRO:HD3	1.99	0.44
1:B:243:ILE:O	1:B:247:THR:HG23	2.18	0.44
1:A:288:THR:N	1:A:289:PRO:CD	2.80	0.44
1:B:124:TRP:CG	1:B:210:CYS:HB2	2.52	0.43
1:B:1004:PHE:O	1:B:1008:ARG:HG3	2.17	0.43
1:B:85:ILE:O	1:B:91:LYS:NZ	2.46	0.43
1:B:1085:LYS:N	1:B:1086:PRO:HD2	2.33	0.43
1:A:121:MET:HE1	1:A:125:PRO:CG	2.49	0.43
1:A:124:TRP:CE2	1:A:126:PHE:HB2	2.53	0.43
1:B:124:TRP:CZ2	1:B:131:CYS:HA	2.54	0.43
1:A:288:THR:HB	1:A:289:PRO:HD3	2.01	0.42
1:B:1024:TYR:CB	1:B:1032:LEU:HD11	2.49	0.42
1:A:199:THR:CG2	1:A:208:ILE:CG2	2.97	0.42
1:A:216:ASP:O	1:A:217:ASP:CB	2.65	0.42
1:A:272:ILE:O	1:A:275:LEU:HB3	2.18	0.42
1:A:65:VAL:O	1:A:69:VAL:HG23	2.18	0.42
1:A:1095:ARG:NE	1:A:1126:TRP:CH2	2.88	0.42
1:B:199:THR:HA	1:B:209:GLU:O	2.19	0.42
1:A:152:MET:CE	1:A:276:VAL:HG22	2.49	0.42
1:A:214:PHE:HB2	1:A:219:TYR:HA	2.02	0.42
1:A:1013:LEU:HD13	1:A:1029:ILE:HG13	2.01	0.42
1:A:93:ALA:HB2	1:A:171:THR:HG22	2.02	0.42
1:B:203:GLU:O	1:B:204:ASP:CB	2.68	0.42
1:A:140:TYR:O	1:A:144:THR:HB	2.20	0.41
1:A:345:CYS:O	1:A:346:PHE:CD2	2.73	0.41
1:A:81:MET:SD	1:A:103:LEU:HA	2.60	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:172:PRO:O	1:B:176:LYS:HG3	2.21	0.41
1:A:1013:LEU:HA	1:A:1028:GLY:O	2.21	0.41
1:A:78:SER:HA	1:A:81:MET:CE	2.50	0.41
1:B:1106:MET:HE1	1:B:1138:TRP:HB2	2.02	0.41
1:B:131:CYS:O	1:B:134:VAL:HG12	2.21	0.41
1:B:152:MET:O	1:B:156:ARG:HG2	2.21	0.41
1:B:214:PHE:HB3	1:B:215:PRO:HD2	2.03	0.41
1:B:161:CYS:HA	1:B:252:ARG:HD3	2.03	0.41
1:B:250:ILE:HG13	1:B:251:LEU:N	2.36	0.41
1:A:1073:ALA:HA	1:A:1076:ARG:NH1	2.36	0.41
1:B:258:LEU:HA	1:B:1158:TRP:CD1	2.55	0.41
1:B:331:ALA:O	1:B:338:LYS:HB2	2.22	0.40
1:B:261:GLY:HA3	1:B:1158:TRP:CE2	2.57	0.40
1:A:214:PHE:CD1	1:A:222:TRP:HB3	2.56	0.40
1:A:58:ILE:CB	1:A:59:PRO:HD3	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	443/480 (92%)	427 (96%)	16 (4%)	0	100	100
1	B	444/480 (92%)	434 (98%)	10 (2%)	0	100	100
All	All	887/960 (92%)	861 (97%)	26 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	370/422 (88%)	353 (95%)	17 (5%)	27	60
1	B	363/422 (86%)	353 (97%)	10 (3%)	43	76
All	All	733/844 (87%)	706 (96%)	27 (4%)	34	68

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	83	VAL
1	A	117	THR
1	A	128	ASP
1	A	158	ILE
1	A	168	ASP
1	A	189	VAL
1	A	199	THR
1	A	203	GLU
1	A	207	VAL
1	A	1026	THR
1	A	1066	LEU
1	A	1104	PHE
1	A	1115	THR
1	A	1127	ASP
1	A	318	LEU
1	A	340	CYS
1	A	345	CYS
1	B	68	VAL
1	B	96	ILE
1	B	115	GLN
1	B	128	ASP
1	B	258	LEU
1	B	1022	GLU
1	B	1055	ASN
1	B	1125	ARG
1	B	266	ASP
1	B	270	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	JDC	B	1300	-	37,37,37	2.73	15 (40%)	46,54,54	1.71	10 (21%)
4	OLC	B	1302	-	15,15,24	1.13	1 (6%)	16,16,25	1.13	2 (12%)
5	PEG	B	1304	-	6,6,6	0.61	0	5,5,5	0.73	0
5	PEG	B	1305	-	6,6,6	0.72	0	5,5,5	0.72	0
4	OLC	B	1303	-	20,20,24	1.06	1 (5%)	21,21,25	0.92	2 (9%)
3	CIT	A	1301	-	3,12,12	1.47	0	3,17,17	2.52	2 (66%)
2	JDC	A	1300	-	37,37,37	2.75	16 (43%)	46,54,54	1.60	9 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	JDC	B	1300	-	-	5/22/47/47	0/4/4/4
4	OLC	B	1302	-	-	7/15/15/24	-
5	PEG	B	1304	-	-	1/4/4/4	-
5	PEG	B	1305	-	-	0/4/4/4	-
4	OLC	B	1303	-	-	4/20/20/24	-
3	CIT	A	1301	-	-	3/6/16/16	-
2	JDC	A	1300	-	-	1/22/47/47	0/4/4/4

All (33) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1300	JDC	C41-C31	-7.76	1.46	1.56
2	A	1300	JDC	C41-C31	-7.56	1.46	1.56
2	A	1300	JDC	C51-C41	-6.53	1.44	1.54
2	B	1300	JDC	C51-C41	-6.41	1.44	1.54
2	A	1300	JDC	C1A-N2A	5.15	1.45	1.34
2	B	1300	JDC	C1A-N2A	4.83	1.44	1.34
4	B	1303	OLC	O20-C1	4.48	1.46	1.33
2	B	1300	JDC	C8-C8A	4.36	1.47	1.39
2	A	1300	JDC	C8-C8A	4.24	1.46	1.39
4	B	1302	OLC	O20-C1	4.12	1.45	1.33
2	A	1300	JDC	C4D-N11	3.93	1.54	1.47
2	B	1300	JDC	C5-C4A	3.82	1.46	1.39
2	A	1300	JDC	C1-N2	-3.79	1.41	1.46
2	A	1300	JDC	C5-C4A	3.77	1.46	1.39
2	A	1300	JDC	C21-C31	-3.74	1.48	1.52
2	A	1300	JDC	O3C-C3C	3.69	1.45	1.37
2	B	1300	JDC	C21-C31	-3.56	1.48	1.52
2	B	1300	JDC	C8A-C4A	-3.55	1.33	1.40
2	B	1300	JDC	O3C-C3C	3.53	1.45	1.37
2	A	1300	JDC	C8A-C4A	-3.50	1.33	1.40
2	B	1300	JDC	C4D-N11	3.50	1.53	1.47
2	B	1300	JDC	C4D-C2A	-3.23	1.51	1.53
2	B	1300	JDC	C1-N2	-3.23	1.42	1.46
2	B	1300	JDC	C6C-C1C	2.74	1.43	1.39
2	A	1300	JDC	C6C-C1C	2.74	1.43	1.39
2	A	1300	JDC	C4C-C3C	2.72	1.44	1.38
2	B	1300	JDC	C4C-C3C	2.70	1.44	1.38
2	B	1300	JDC	C2A-N2A	-2.40	1.42	1.46
2	A	1300	JDC	C34-C31	2.20	1.58	1.53
2	A	1300	JDC	C4D-C2A	-2.13	1.51	1.53
2	B	1300	JDC	C34-C31	2.12	1.57	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1300	JDC	C2A-N2A	-2.05	1.42	1.46
2	A	1300	JDC	C6-C5	2.02	1.42	1.38

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1300	JDC	C61-N11-C4D	4.31	123.72	111.20
2	A	1300	JDC	C61-N11-C4D	4.31	123.71	111.20
2	A	1300	JDC	C4E-C41-C51	-4.09	104.31	109.11
2	B	1300	JDC	C4E-C41-C31	-3.81	106.48	111.00
3	A	1301	CIT	C3-C4-C5	-3.81	108.89	114.98
2	B	1300	JDC	C3A-C2A-N2A	-3.77	103.44	110.93
2	B	1300	JDC	C61-N11-C21	3.75	115.25	109.52
2	B	1300	JDC	C34-C31-C41	-3.53	110.44	113.87
2	A	1300	JDC	C3A-C2A-N2A	-3.35	104.29	110.93
2	A	1300	JDC	C34-C31-C41	-3.25	110.71	113.87
2	B	1300	JDC	C51-C41-C31	3.15	112.25	108.06
2	A	1300	JDC	C51-C41-C31	3.02	112.08	108.06
4	B	1302	OLC	O20-C1-C2	2.97	121.23	111.91
2	B	1300	JDC	C4D-N11-C21	2.90	115.56	111.56
4	B	1303	OLC	O20-C1-C2	2.76	120.56	111.91
2	A	1300	JDC	C51-C61-N11	2.33	113.53	111.23
2	B	1300	JDC	C4E-C41-C51	-2.33	106.38	109.11
2	B	1300	JDC	C2A-N2A-C1A	-2.17	119.24	123.07
2	A	1300	JDC	C4E-C41-C31	-2.10	108.51	111.00
3	A	1301	CIT	C3-C2-C1	-2.08	111.65	114.98
2	A	1300	JDC	C34-C31-C21	-2.06	107.88	110.22
2	A	1300	JDC	C61-N11-C21	2.03	112.62	109.52
4	B	1303	OLC	O20-C1-O19	-2.03	118.48	123.59
2	B	1300	JDC	C4E-C41-C1C	-2.02	105.28	109.38
4	B	1302	OLC	O20-C1-O19	-2.00	118.54	123.59

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1300	JDC	O1-C1A-C3-N2
2	B	1300	JDC	N2A-C1A-C3-N2
4	B	1303	OLC	O20-C21-C22-O23
3	A	1301	CIT	C2-C3-C4-C5
3	A	1301	CIT	O7-C3-C4-C5
3	A	1301	CIT	C6-C3-C4-C5

Continued on next page...

Continued from previous page...




Mol	Chain	Res	Type	Atoms
4	B	1302	OLC	C2-C1-O20-C21
4	B	1302	OLC	O19-C1-O20-C21
4	B	1302	OLC	O20-C21-C22-O23
4	B	1302	OLC	O20-C21-C22-C24
4	B	1303	OLC	O20-C21-C22-C24
4	B	1302	OLC	C2-C3-C4-C5
4	B	1303	OLC	C1-C2-C3-C4
4	B	1302	OLC	C5-C6-C7-C8
4	B	1302	OLC	C4-C5-C6-C7
2	A	1300	JDC	O1-C1A-C3-N2
2	B	1300	JDC	C2A-C4D-N11-C21
5	B	1304	PEG	O2-C3-C4-O4
2	B	1300	JDC	O1-C1A-C3-C4
2	B	1300	JDC	N2A-C1A-C3-C4
4	B	1303	OLC	C9-C10-C11-C12


There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1300	JDC	2	0
4	B	1302	OLC	1	0
4	B	1303	OLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	447/480 (93%)	-0.26	2 (0%) 92 93	48, 79, 139, 260	0
1	B	448/480 (93%)	-0.16	14 (3%) 49 44	39, 72, 141, 239	0
All	All	895/960 (93%)	-0.21	16 (1%) 68 67	39, 76, 142, 260	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	204	ASP	4.4
1	B	217	ASP	3.3
1	A	206	ASP	3.3
1	B	201	VAL	3.2
1	B	305	SER	3.1
1	B	206	ASP	2.8
1	B	120	LEU	2.8
1	B	216	ASP	2.7
1	B	199	THR	2.6
1	B	207	VAL	2.6
1	B	266	ASP	2.6
1	B	205	VAL	2.4
1	A	204	ASP	2.2
1	B	203	GLU	2.2
1	B	219	TYR	2.2
1	B	306	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	PEG	B	1304	7/7	0.74	0.19	75,93,104,105	0
3	CIT	A	1301	13/13	0.76	0.35	129,148,161,166	0
5	PEG	B	1305	7/7	0.82	0.35	93,94,100,101	0
4	OLC	B	1303	21/25	0.87	0.22	52,75,134,145	0
4	OLC	B	1302	16/25	0.93	0.23	54,68,72,78	0
2	JDC	B	1300	34/34	0.95	0.18	48,65,92,99	0
2	JDC	A	1300	34/34	0.96	0.16	43,62,79,90	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around OLC B 1303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLC B 1302:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around JDC B 1300:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.