



Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 11:13 pm BST

PDB ID : 5DJB
Title : Structure of the Haliangium ochraceum BMC-H shell protein
Authors : Sutter, M.; Kerfeld, C.A.
Deposited on : 2015-09-01
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	99	<div><div></div><div>93%6%.</div></div>
1	B	99	<div><div></div><div>93%6%.</div></div>
1	C	99	<div><div></div><div>91%5%.</div></div>
1	D	99	<div><div></div><div>97%..</div></div>
1	E	99	<div><div></div><div>86%9%5%</div></div>
1	F	99	<div><div></div><div>90%5%5%</div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	99	<div><div></div><div>86%</div><div>9%5%</div></div>
1	H	99	<div><div></div><div>91%</div><div>• 5%</div></div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 12048 atoms, of which 5605 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Microcompartments protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	98	Total	C	H	N	O	S	0	0	0
			1415	436	717	127	131	4			
1	B	98	Total	C	H	N	O	S	0	0	0
			1415	436	717	127	131	4			
1	C	95	Total	C	H	N	O	S	0	0	0
			1372	424	694	123	127	4			
1	D	98	Total	C	H	N	O	S	0	0	0
			1415	436	717	127	131	4			
1	E	94	Total	C	H	N	O	S	0	0	0
			1360	420	690	122	124	4			
1	F	94	Total	C	H	N	O	S	0	0	0
			1360	420	690	122	124	4			
1	G	94	Total	C	H	N	O	S	0	0	0
			1360	420	690	122	124	4			
1	H	94	Total	C	H	N	O	S	0	0	0
			1360	420	690	122	124	4			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	114	Total	O	0	0
			114	114		
2	B	134	Total	O	0	0
			134	134		
2	C	107	Total	O	0	0
			107	107		
2	D	121	Total	O	0	0
			121	121		
2	E	142	Total	O	0	0
			142	142		
2	F	115	Total	O	0	0
			115	115		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	117	Total	O	0	0
			117	117		
2	H	141	Total	O	0	0
			141	141		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Microcompartments protein

Chain A:  93% 6%



- Molecule 1: Microcompartments protein

Chain B:  93% 6%



- Molecule 1: Microcompartments protein

Chain C:  91% 5%




- Molecule 1: Microcompartments protein

Chain D:  97%




- Molecule 1: Microcompartments protein

Chain E:  86% 9% 5%




- Molecule 1: Microcompartments protein




Chain F:  90% 5% 5%



- Molecule 1: Microcompartments protein



- Molecule 1: Microcompartments protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 3	Depositor
Cell constants a, b, c, α , β , γ	69.20 Å 69.20 Å 120.42 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	33.35 – 1.80 33.35 – 1.80	Depositor EDS
% Data completeness (in resolution range)	98.7 (33.35-1.80) 98.7 (33.35-1.80)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 1.81 Å)	Xtriage
Refinement program	PHENIX 1.9_1690	Depositor
R, R_{free}	0.170 , 0.204 0.150 , 0.175	Depositor DCC
R_{free} test set	2007 reflections (3.40%)	wwPDB-VP
Wilson B-factor (Å ²)	15.1	Xtriage
Anisotropy	0.731	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 46.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	0.468 for -h,-k,l 0.469 for h,-h-k,-l 0.470 for -k,-h,-l	Xtriage
Reported twinning fraction	0.500 for -k,-h,-l	Depositor
Outliers	0 of 59085 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12048	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 21.59 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.7895e-03.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.22	0/706	0.43	0/956
1	B	0.22	0/706	0.42	0/956
1	C	0.22	0/686	0.43	0/930
1	D	0.22	0/706	0.41	0/956
1	E	0.23	0/678	0.45	0/919
1	F	0.22	0/678	0.44	0/919
1	G	0.24	0/678	0.45	0/919
1	H	0.23	0/678	0.44	0/919
All	All	0.22	0/5516	0.43	0/7474

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	698	717	717	5	0
1	B	698	717	717	6	0
1	C	678	694	694	6	0
1	D	698	717	717	1	0
1	E	670	690	690	10	0
1	F	670	690	690	5	0
1	G	670	690	690	9	0
1	H	670	690	690	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	114	0	0	4	3
2	B	134	0	0	4	2
2	C	107	0	0	6	0
2	D	121	0	0	1	0
2	E	142	0	0	8	1
2	F	115	0	0	5	1
2	G	117	0	0	6	1
2	H	141	0	0	4	1
All	All	6443	5605	5605	44	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:16:MET:SD	2:E:113:HOH:O	2.31	0.89
1:E:66:ARG:NH1	1:G:66:ARG:O	2.10	0.84
1:F:42:VAL:O	2:F:101:HOH:O	2.01	0.76
1:H:87:LEU:O	2:H:101:HOH:O	2.03	0.76
1:E:64:ALA:O	2:E:101:HOH:O	2.07	0.73
1:G:69:GLU:O	2:G:101:HOH:O	2.08	0.69
1:H:83:VAL:O	2:H:102:HOH:O	2.10	0.68
1:B:7:MET:SD	2:B:208:HOH:O	2.52	0.67
1:G:66:ARG:NH1	2:G:106:HOH:O	2.29	0.66
1:A:99:ALA:O	2:A:101:HOH:O	2.13	0.66
1:E:62:ARG:NH2	2:E:108:HOH:O	2.30	0.64
1:C:64:ALA:N	2:C:105:HOH:O	2.30	0.63
1:B:60:GLY:N	2:B:102:HOH:O	2.32	0.61
1:C:66:ARG:NH1	2:C:106:HOH:O	2.33	0.61
1:A:11:ARG:NH1	2:A:109:HOH:O	2.34	0.59
1:A:64:ALA:N	2:A:108:HOH:O	2.34	0.59
1:E:6:GLY:N	2:E:105:HOH:O	2.36	0.58
1:D:49:ASP:OD1	2:D:101:HOH:O	2.17	0.58
1:E:66:ARG:NH2	2:E:110:HOH:O	2.35	0.58
1:E:68:GLY:N	2:E:101:HOH:O	2.35	0.58
1:F:95:MET:SD	2:F:122:HOH:O	2.58	0.56
1:E:66:ARG:O	1:G:66:ARG:NH2	2.40	0.54
1:G:49:ASP:OD2	2:G:102:HOH:O	2.17	0.54
1:E:54:LYS:NZ	2:E:111:HOH:O	2.37	0.53
1:C:78:ARG:NH2	2:C:111:HOH:O	2.42	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:66:ARG:NE	2:C:112:HOH:O	2.42	0.52
1:F:91:ARG:NH1	2:F:109:HOH:O	2.45	0.50
1:C:84:ASP:OD2	2:C:101:HOH:O	2.19	0.49
1:C:43:THR:N	2:C:118:HOH:O	2.45	0.49
1:E:61:GLN:OE1	2:E:102:HOH:O	2.20	0.49
1:H:87:LEU:N	2:H:102:HOH:O	2.48	0.47
1:G:11:ARG:NH1	2:G:118:HOH:O	2.47	0.46
1:B:7:MET:HE1	1:B:89:LEU:HD11	1.97	0.46
1:H:49:ASP:OD2	2:H:103:HOH:O	2.20	0.45
1:G:19:ALA:HB2	1:G:64:ALA:HB2	1.98	0.45
1:B:61:GLN:NE2	2:B:113:HOH:O	2.49	0.44
1:B:7:MET:CE	1:B:89:LEU:HD11	2.46	0.44
1:A:49:ASP:OD2	2:A:102:HOH:O	2.21	0.44
1:B:49:ASP:OD1	2:B:101:HOH:O	2.21	0.43
1:F:49:ASP:OD1	2:F:102:HOH:O	2.21	0.43
1:G:53:VAL:HG12	2:G:132:HOH:O	2.17	0.43
1:G:65:GLU:OE1	2:G:103:HOH:O	2.22	0.42
1:A:87:LEU:HB2	1:A:89:LEU:HD23	2.01	0.41
1:F:10:VAL:N	2:F:101:HOH:O	2.48	0.41

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:154:HOH:O	2:B:129:HOH:O[3_775]	2.09	0.11
2:A:101:HOH:O	2:G:113:HOH:O[1_556]	2.15	0.05
2:E:129:HOH:O	2:F:120:HOH:O[3_775]	2.15	0.05
2:H:105:HOH:O	2:H:132:HOH:O[3_675]	2.17	0.03
2:A:155:HOH:O	2:B:135:HOH:O[3_775]	2.18	0.02

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	96/99 (97%)	93 (97%)	3 (3%)	0	100	100
1	B	96/99 (97%)	93 (97%)	3 (3%)	0	100	100
1	C	93/99 (94%)	90 (97%)	3 (3%)	0	100	100
1	D	96/99 (97%)	93 (97%)	3 (3%)	0	100	100
1	E	92/99 (93%)	89 (97%)	3 (3%)	0	100	100
1	F	92/99 (93%)	89 (97%)	3 (3%)	0	100	100
1	G	92/99 (93%)	89 (97%)	3 (3%)	0	100	100
1	H	92/99 (93%)	89 (97%)	3 (3%)	0	100	100
All	All	749/792 (95%)	725 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	67/68 (98%)	67 (100%)	0	100	100
1	B	67/68 (98%)	66 (98%)	1 (2%)	65	56
1	C	65/68 (96%)	65 (100%)	0	100	100
1	D	67/68 (98%)	66 (98%)	1 (2%)	65	56
1	E	64/68 (94%)	63 (98%)	1 (2%)	62	54
1	F	64/68 (94%)	64 (100%)	0	100	100
1	G	64/68 (94%)	63 (98%)	1 (2%)	62	54
1	H	64/68 (94%)	63 (98%)	1 (2%)	62	54
All	All	522/544 (96%)	517 (99%)	5 (1%)	76	71

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	82	ASN
1	D	82	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	82	ASN
1	G	82	ASN
1	H	89	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	98/99 (98%)	-0.74	0 100 100	11, 15, 20, 23	0
1	B	98/99 (98%)	-0.72	0 100 100	11, 15, 20, 29	0
1	C	95/99 (95%)	-0.65	0 100 100	13, 17, 22, 26	0
1	D	98/99 (98%)	-0.69	0 100 100	12, 16, 23, 28	0
1	E	94/99 (94%)	-0.74	0 100 100	12, 15, 20, 26	0
1	F	94/99 (94%)	-0.74	0 100 100	13, 15, 21, 24	0
1	G	94/99 (94%)	-0.74	0 100 100	10, 14, 20, 25	0
1	H	94/99 (94%)	-0.75	0 100 100	10, 13, 19, 21	0
All	All	765/792 (96%)	-0.72	0 100 100	10, 15, 21, 29	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.