



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 20, 2020 – 12:19 PM BST

PDB ID : 4DKW  
Title : Structure of P22 Large terminase nuclease domain  
Authors : Roy, A.; Cingolani, G.  
Deposited on : 2012-02-04  
Resolution : 2.02 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	10434 (2.04-2.00)
Clashscore	141614	11643 (2.04-2.00)
Ramachandran outliers	138981	11493 (2.04-2.00)
Sidechain outliers	138945	11492 (2.04-2.00)
RSRZ outliers	127900	10220 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	211	<div> <div>6%</div> <div> <div></div> <div>76%</div> <div>14%</div> <div>•</div> <div>8%</div> </div> </div>
1	B	211	<div> <div>6%</div> <div> <div></div> <div>75%</div> <div>15%</div> <div>•</div> <div>8%</div> </div> </div>
1	C	211	<div> <div>6%</div> <div> <div></div> <div>75%</div> <div>15%</div> <div>•</div> <div>8%</div> </div> </div>
1	D	211	<div> <div>6%</div> <div> <div></div> <div>76%</div> <div>13%</div> <div>•</div> <div>8%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	B	503	-	-	X	-

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7516 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Large terminase protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	194	Total	C	N	O	S	0	0	0
			1594	1014	281	289	10			
1	B	194	Total	C	N	O	S	0	0	0
			1594	1014	281	289	10			
1	C	194	Total	C	N	O	S	0	0	0
			1594	1014	281	289	10			
1	D	194	Total	C	N	O	S	0	0	0
			1594	1014	281	289	10			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Mg	0	0
			2	2		
2	A	2	Total	Mg	0	0
			2	2		
2	D	2	Total	Mg	0	0
			2	2		
2	C	2	Total	Mg	0	0
			2	2		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	287	Total	O	0	0
			287	287		
4	B	267	Total	O	0	0
			267	267		
4	C	296	Total	O	0	0
			296	296		
4	D	262	Total	O	0	0
			262	262		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Large terminase protein




- Molecule 1: Large terminase protein



- Molecule 1: Large terminase protein



- Molecule 1: Large terminase protein



K452
PRO
LYS
GLU
LYS
LYS
LYS
ILE
PRO
ALA
PRO
ILE
ARG
PRO
VAL
ARG
ARG
GLY
ARG

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	60.92Å 139.79Å 61.01Å 90.00° 95.10° 90.00°	Depositor
Resolution (Å)	14.97 – 2.02 14.97 – 2.01	Depositor EDS
% Data completeness (in resolution range)	81.7 (14.97-2.02) 81.8 (14.97-2.01)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.66 (at 2.01Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, $R_{free}$	0.166 , 0.197 0.160 , 0.196	Depositor DCC
$R_{free}$ test set	1960 reflections (3.30%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.0	Xtriage
Anisotropy	0.098	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 47.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.457 for l,-k,h	Xtriage
Reported twinning fraction	0.493 for l,-k,h	Depositor
Outliers	0 of 59308 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7516	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.84% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.38	0/1643	0.53	0/2220
1	B	0.38	0/1643	0.54	0/2220
1	C	0.37	0/1643	0.54	0/2220
1	D	0.37	0/1643	0.53	0/2220
All	All	0.38	0/6572	0.53	0/8880

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1594	0	1499	29	0
1	B	1594	0	1499	35	0
1	C	1594	0	1499	27	1
1	D	1594	0	1499	30	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	5	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	5	0	0	2	0
3	C	5	0	0	0	0
3	D	5	0	0	1	0
4	A	287	0	0	12	1
4	B	267	0	0	12	3
4	C	296	0	0	13	4
4	D	262	0	0	19	1
All	All	7516	0	5996	120	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 120 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:383:GLY:CA	1:B:387:GLN:HG3	1.57	1.35
1:B:383:GLY:HA2	1:B:387:GLN:HG3	1.26	1.05
1:D:301:GLU:HG3	4:D:712:HOH:O	1.60	1.01
1:B:383:GLY:HA3	1:B:387:GLN:HG3	1.45	0.98
1:C:378:HIS:NE2	4:C:875:HOH:O	1.99	0.96

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:701:HOH:O	4:C:722:HOH:O[2_354]	2.06	0.14
4:B:723:HOH:O	4:C:734:HOH:O[2_354]	2.09	0.11
1:C:382:LYS:NZ	4:D:719:HOH:O[1_556]	2.11	0.09
4:A:782:HOH:O	4:C:677:HOH:O[1_554]	2.12	0.08
4:B:623:HOH:O	4:C:771:HOH:O[2_354]	2.17	0.03

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	192/211 (91%)	177 (92%)	13 (7%)	2 (1%)	15	9
1	B	192/211 (91%)	179 (93%)	11 (6%)	2 (1%)	15	9
1	C	192/211 (91%)	178 (93%)	12 (6%)	2 (1%)	15	9
1	D	192/211 (91%)	178 (93%)	12 (6%)	2 (1%)	15	9
All	All	768/844 (91%)	712 (93%)	48 (6%)	8 (1%)	15	9

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	375	HIS
1	A	480	ILE
1	B	375	HIS
1	B	480	ILE
1	C	480	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	165/180 (92%)	158 (96%)	7 (4%)	30	26
1	B	165/180 (92%)	159 (96%)	6 (4%)	35	32
1	C	165/180 (92%)	159 (96%)	6 (4%)	35	32
1	D	165/180 (92%)	158 (96%)	7 (4%)	30	26
All	All	660/720 (92%)	634 (96%)	26 (4%)	32	29

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	387	GLN
1	C	301	GLU
1	D	445	TYR
1	B	389	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	289	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	313	HIS
1	B	313	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	A	503	-	4,4,4	0.15	0	6,6,6	0.05	0
3	SO4	B	503	-	4,4,4	0.14	0	6,6,6	0.05	0
3	SO4	C	503	-	4,4,4	0.16	0	6,6,6	0.21	0
3	SO4	D	503	-	4,4,4	0.15	0	6,6,6	0.07	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	503	SO4	2	0
3	D	503	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	194/211 (91%)	-0.36	12 (6%) 20 19	17, 30, 99, 171	0
1	B	194/211 (91%)	-0.32	12 (6%) 20 19	17, 30, 100, 171	0
1	C	194/211 (91%)	-0.30	12 (6%) 20 19	17, 30, 100, 171	0
1	D	194/211 (91%)	-0.34	13 (6%) 17 17	17, 30, 99, 171	0
All	All	776/844 (91%)	-0.33	49 (6%) 20 19	17, 30, 114, 171	0

The worst 5 of 49 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	380	HIS	12.4
1	C	380	HIS	12.4
1	B	383	GLY	12.2
1	D	380	HIS	11.9
1	C	382	LYS	11.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	SO4	B	503	5/5	0.85	0.15	111,116,118,129	0
3	SO4	A	503	5/5	0.89	0.16	112,118,120,122	0
2	MG	B	502	1/1	0.92	0.14	32,32,32,32	0
2	MG	B	501	1/1	0.95	0.06	28,28,28,28	0
2	MG	A	502	1/1	0.96	0.18	33,33,33,33	0
2	MG	C	502	1/1	0.97	0.17	24,24,24,24	0
2	MG	D	501	1/1	0.97	0.10	42,42,42,42	0
2	MG	C	501	1/1	0.97	0.05	22,22,22,22	0
2	MG	A	501	1/1	0.98	0.10	36,36,36,36	0
3	SO4	C	503	5/5	0.98	0.07	57,65,70,72	0
3	SO4	D	503	5/5	0.99	0.06	38,51,62,70	0
2	MG	D	502	1/1	0.99	0.21	29,29,29,29	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.