



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 29, 2020 – 03:39 PM BST

PDB ID : 6DK7
Title : RetS histidine kinase region with cobalt
Authors : Mancl, J.M.; Schubot, F.D.
Deposited on : 2018-05-29
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.13
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	227	<div> <div>4%</div> <div> <div></div> <div>67%</div> <div>29%</div> <div>..</div> </div> </div>
1	B	227	<div> <div>6%</div> <div> <div></div> <div>78%</div> <div>19%</div> <div>..</div> </div> </div>
1	C	227	<div> <div>3%</div> <div> <div></div> <div>78%</div> <div>19%</div> <div>..</div> </div> </div>
1	D	227	<div> <div>4%</div> <div> <div></div> <div>78%</div> <div>19%</div> <div>..</div> </div> </div>
1	E	227	<div> <div>%</div> <div> <div></div> <div>79%</div> <div>19%</div> <div>..</div> </div> </div>
1	F	227	<div> <div>4%</div> <div> <div></div> <div>77%</div> <div>19%</div> <div>..</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	227	<div><div></div><div>2%</div><div>80%</div><div>17%</div><div>••</div></div>
1	H	227	<div><div></div><div>7%</div><div>78%</div><div>21%</div><div>•</div></div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 14138 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RetS (Regulator of Exopolysaccharide and Type III Secretion).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	223	Total	C	N	O	S	0	0	0
			1724	1087	296	337	4			
1	C	225	Total	C	N	O	S	0	0	0
			1735	1093	298	340	4			
1	D	225	Total	C	N	O	S	0	0	0
			1735	1093	298	340	4			
1	B	224	Total	C	N	O	S	0	0	0
			1731	1091	297	339	4			
1	E	225	Total	C	N	O	S	0	0	0
			1736	1093	298	341	4			
1	F	223	Total	C	N	O	S	0	0	0
			1726	1087	297	338	4			
1	G	225	Total	C	N	O	S	0	0	0
			1739	1095	299	341	4			
1	H	225	Total	C	N	O	S	0	0	0
			1738	1095	299	340	4			

- Molecule 2 is COBALT (II) ION (three-letter code: CO) (formula: Co).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	2	Total	Co	0	0
			2	2		
2	D	3	Total	Co	0	0
			3	3		
2	E	4	Total	Co	0	0
			4	4		
2	H	3	Total	Co	0	0
			3	3		
2	B	4	Total	Co	0	0
			4	4		
2	C	4	Total	Co	0	0
			4	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Co	0	0
			2	2		
2	F	2	Total	Co	0	0
			2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	29	Total	O	0	0
			29	29		
3	C	23	Total	O	0	0
			23	23		
3	D	41	Total	O	0	0
			41	41		
3	B	33	Total	O	0	0
			33	33		
3	E	32	Total	O	0	0
			32	32		
3	F	38	Total	O	0	0
			38	38		
3	G	22	Total	O	0	0
			22	22		
3	H	32	Total	O	0	0
			32	32		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RetS (Regulator of Exopolysaccharide and Type III Secretion)




- Molecule 1: RetS (Regulator of Exopolysaccharide and Type III Secretion)




- Molecule 1: RetS (Regulator of Exopolysaccharide and Type III Secretion)




- Molecule 1: RetS (Regulator of Exopolysaccharide and Type III Secretion)




- Molecule 1: RetS (Regulator of Exopolysaccharide and Type III Secretion)




- Molecule 1: RetS (Regulator of Exopolysaccharide and Type III Secretion)



- Molecule 1: RetS (Regulator of Exopolysaccharide and Type III Secretion)



- Molecule 1: RetS (Regulator of Exopolysaccharide and Type III Secretion)





4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	158.71Å 158.71Å 243.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	56.78 – 2.60 56.78 – 2.60	Depositor EDS
% Data completeness (in resolution range)	96.3 (56.78-2.60) 96.3 (56.78-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.42 (at 2.61Å)	Xtriage
Refinement program	PHENIX 1.10.1_2155	Depositor
R, R_{free}	0.211 , 0.261 0.212 , 0.258	Depositor DCC
R_{free} test set	2519 reflections (2.72%)	wwPDB-VP
Wilson B-factor (Å ²)	51.2	Xtriage
Anisotropy	0.226	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 52.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	14138	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section:
CO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/1745	0.60	2/2360 (0.1%)
1	B	0.36	0/1752	0.53	0/2370
1	C	0.37	0/1756	0.55	1/2375 (0.0%)
1	D	0.49	1/1756 (0.1%)	0.59	2/2375 (0.1%)
1	E	0.32	0/1758	0.57	3/2379 (0.1%)
1	F	0.46	2/1746 (0.1%)	0.55	1/2360 (0.0%)
1	G	0.47	0/1760	0.62	2/2380 (0.1%)
1	H	0.43	1/1759 (0.1%)	0.57	1/2379 (0.0%)
All	All	0.43	4/14032 (0.0%)	0.57	12/18978 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	548	GLU	CD-OE1	-6.95	1.18	1.25
1	F	548	GLU	CD-OE2	-6.44	1.18	1.25
1	H	620	GLU	CD-OE1	-5.91	1.19	1.25
1	D	518	PRO	N-CD	5.10	1.54	1.47

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	442	GLY	C-N-CA	7.24	139.80	121.70
1	H	563	THR	C-N-CD	6.95	142.99	128.40
1	E	442	GLY	C-N-CA	6.91	138.97	121.70
1	F	548	GLU	OE1-CD-OE2	-6.71	115.25	123.30
1	D	563	THR	C-N-CD	5.88	140.75	128.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1724	0	1758	47	0
1	B	1731	0	1765	27	0
1	C	1735	0	1767	32	0
1	D	1735	0	1768	33	0
1	E	1736	0	1768	43	0
1	F	1726	0	1756	42	0
1	G	1739	0	1771	47	0
1	H	1738	0	1771	35	0
2	A	2	0	0	0	0
2	B	4	0	0	0	0
2	C	4	0	0	0	0
2	D	3	0	0	0	0
2	E	4	0	0	0	0
2	F	2	0	0	0	0
2	G	2	0	0	0	0
2	H	3	0	0	0	0
3	A	29	0	0	6	0
3	B	33	0	0	4	0
3	C	23	0	0	2	0
3	D	41	0	0	3	0
3	E	32	0	0	2	0
3	F	38	0	0	2	0
3	G	22	0	0	1	0
3	H	32	0	0	5	0
All	All	14138	0	14124	272	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 272 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:627:SER:HB2	1:D:629:GLN:N	1.15	1.45
1:G:448:LYS:HE3	1:G:452:TYR:CZ	1.62	1.32
1:C:627:SER:HA	1:C:629:GLN:N	1.46	1.29

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:591:HIS:NE2	3:H:801:HOH:O	1.69	1.23
1:C:627:SER:CA	1:C:629:GLN:N	2.05	1.20

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	219/227 (96%)	213 (97%)	6 (3%)	0	100	100
1	B	220/227 (97%)	215 (98%)	5 (2%)	0	100	100
1	C	221/227 (97%)	221 (100%)	0	0	100	100
1	D	221/227 (97%)	218 (99%)	2 (1%)	1 (0%)	29	52
1	E	223/227 (98%)	219 (98%)	4 (2%)	0	100	100
1	F	217/227 (96%)	214 (99%)	3 (1%)	0	100	100
1	G	221/227 (97%)	219 (99%)	2 (1%)	0	100	100
1	H	221/227 (97%)	219 (99%)	2 (1%)	0	100	100
All	All	1763/1816 (97%)	1738 (99%)	24 (1%)	1 (0%)	51	75

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	602	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	192/195 (98%)	181 (94%)	11 (6%)	20	41
1	B	193/195 (99%)	183 (95%)	10 (5%)	23	46
1	C	193/195 (99%)	184 (95%)	9 (5%)	26	50
1	D	193/195 (99%)	180 (93%)	13 (7%)	16	33
1	E	194/195 (100%)	188 (97%)	6 (3%)	40	66
1	F	192/195 (98%)	186 (97%)	6 (3%)	40	66
1	G	194/195 (100%)	186 (96%)	8 (4%)	30	56
1	H	193/195 (99%)	185 (96%)	8 (4%)	30	56
All	All	1544/1560 (99%)	1473 (95%)	71 (5%)	27	51

5 of 71 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	635	LEU
1	B	529	THR
1	H	508	GLN
1	B	458	SER
1	B	508	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	519	GLN
1	G	604	HIS
1	E	560	GLN
1	A	591	HIS
1	G	571	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 24 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	223/227 (98%)	0.16	10 (4%) 33 26	35, 58, 95, 111	0
1	B	224/227 (98%)	0.30	14 (6%) 20 15	37, 58, 93, 112	0
1	C	225/227 (99%)	0.17	6 (2%) 54 48	32, 56, 103, 126	0
1	D	225/227 (99%)	0.26	10 (4%) 34 27	34, 53, 90, 117	0
1	E	225/227 (99%)	0.12	3 (1%) 77 73	34, 56, 102, 123	0
1	F	223/227 (98%)	0.17	8 (3%) 42 35	36, 57, 93, 109	0
1	G	225/227 (99%)	0.16	5 (2%) 62 56	35, 55, 100, 119	0
1	H	225/227 (99%)	0.34	15 (6%) 17 13	33, 61, 100, 117	0
All	All	1795/1816 (98%)	0.21	71 (3%) 38 31	32, 57, 99, 126	0

The worst 5 of 71 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	507	GLN	5.3
1	F	509	ARG	4.7
1	H	601	LEU	4.3
1	B	601	LEU	4.2
1	H	625	SER	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CO	A	701	1/1	0.49	0.22	132,132,132,132	0
2	CO	E	704	1/1	0.74	0.11	109,109,109,109	0
2	CO	F	702	1/1	0.76	0.12	132,132,132,132	0
2	CO	H	701	1/1	0.79	0.09	106,106,106,106	0
2	CO	D	703	1/1	0.81	0.12	124,124,124,124	0
2	CO	C	704	1/1	0.82	0.21	117,117,117,117	0
2	CO	G	702	1/1	0.83	0.12	126,126,126,126	0
2	CO	H	702	1/1	0.86	0.07	123,123,123,123	0
2	CO	H	703	1/1	0.87	0.20	57,57,57,57	0
2	CO	F	701	1/1	0.88	0.12	97,97,97,97	0
2	CO	B	703	1/1	0.94	0.15	92,92,92,92	0
2	CO	E	701	1/1	0.95	0.22	56,56,56,56	0
2	CO	D	702	1/1	0.95	0.11	76,76,76,76	0
2	CO	G	701	1/1	0.96	0.20	72,72,72,72	0
2	CO	C	702	1/1	0.97	0.20	56,56,56,56	0
2	CO	B	701	1/1	0.97	0.18	56,56,56,56	0
2	CO	B	704	1/1	0.97	0.04	119,119,119,119	0
2	CO	A	702	1/1	0.97	0.27	89,89,89,89	0
2	CO	C	701	1/1	0.98	0.25	64,64,64,64	0
2	CO	B	702	1/1	0.98	0.28	56,56,56,56	0
2	CO	E	703	1/1	0.98	0.17	70,70,70,70	0
2	CO	C	703	1/1	0.99	0.18	63,63,63,63	0
2	CO	E	702	1/1	0.99	0.27	59,59,59,59	0
2	CO	D	701	1/1	1.00	0.20	34,34,34,34	0

6.5 Other polymers ⓘ

There are no such residues in this entry.