



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 26, 2021 – 12:10 PM JST

PDB ID : 7DKC
Title : Stenotrophomonas maltophilia DPP7 in complex with Tyr-Tyr
Authors : Sakamoto, Y.; Nakamura, A.; Suzuki, Y.; Honma, N.; Roppongi, S.; Kushibiki, C.; Yonezawa, N.; Takahashi, M.; Shida, Y.; Gouda, H.; Nonaka, T.; Ogasawara, W.; Tanaka, N.
Deposited on : 2020-11-23
Resolution : 1.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.23.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	720	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 98%);"></div> <div style="display: flex; justify-content: space-between; margin-top: 2px;"> % 90% 7% . </div> </div>
1	B	720	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 3%, orange 1%, yellow 1%, green 95%);"></div> <div style="display: flex; justify-content: space-between; margin-top: 2px;"> 3% 88% 9% . </div> </div>

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 12105 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dipeptidyl-peptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	697	Total	C	N	O	S	0	0	0
			5329	3375	927	1008	19			
1	B	697	Total	C	N	O	S	0	0	0
			5329	3375	927	1008	19			

- Molecule 2 is TYROSINE (three-letter code: TYR) (formula: C₉H₁₁NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			12	9	1	2		
2	A	1	Total	C	N	O	0	0
			13	9	1	3		
2	B	1	Total	C	N	O	0	0
			12	9	1	2		
2	B	1	Total	C	N	O	0	0
			13	9	1	3		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	866	Total	O	0	0
			866	866		
4	B	495	Total	O	0	0
			495	495		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Dipeptidyl-peptidase



• Molecule 1: Dipeptidyl-peptidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	67.85Å 74.32Å 153.32Å 90.00° 94.15° 90.00°	Depositor
Resolution (Å)	40.00 – 1.86 36.70 – 1.86	Depositor EDS
% Data completeness (in resolution range)	99.8 (40.00-1.86) 99.8 (36.70-1.86)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.83 (at 1.85Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.171 , 0.210 0.180 , 0.215	Depositor DCC
R_{free} test set	6217 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	22.9	Xtriage
Anisotropy	0.054	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 42.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12105	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	0/5442	0.86	2/7375 (0.0%)
1	B	0.73	0/5442	0.83	0/7375
All	All	0.74	0/10884	0.85	2/14750 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	513	ARG	CG-CD-NE	-7.16	96.77	111.80
1	A	579	ARG	NE-CZ-NH1	5.41	123.00	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5329	0	5271	36	0
1	B	5329	0	5271	41	0
2	A	25	0	17	1	0
2	B	25	0	17	2	0
3	A	24	0	32	3	0
3	B	12	0	16	0	0
4	A	866	0	0	12	2

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	495	0	0	4	0
All	All	12105	0	10624	77	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:805:GOL:H31	4:A:1327:HOH:O	1.50	1.09
3:A:805:GOL:C3	4:A:1327:HOH:O	2.14	0.87
1:A:334:LEU:HD12	4:A:1408:HOH:O	1.74	0.86
1:A:576:LEU:HD23	4:A:1034:HOH:O	1.80	0.80
1:A:576:LEU:CD2	4:A:1034:HOH:O	2.29	0.79
1:B:36:ALA:O	1:B:40:GLN:HG3	1.89	0.71
1:A:275:ASN:HD22	1:A:681:ASN:HD21	1.41	0.67
1:B:220:THR:H	1:B:643:ASN:HD21	1.42	0.67
1:B:696:SER:HA	1:B:699:MET:CE	2.27	0.64
1:A:272:GLY:HA2	1:A:576:LEU:HD23	1.79	0.64
1:A:700:GLN:HE21	1:A:704:GLN:NE2	1.95	0.63
1:A:220:THR:H	1:A:643:ASN:HD21	1.47	0.61
1:B:655:SER:OG	2:B:802:TYR:C	2.41	0.59
1:A:712:LEU:HD13	4:A:1494:HOH:O	2.04	0.57
1:A:275:ASN:HD22	1:A:681:ASN:ND2	2.03	0.56
1:B:696:SER:HA	1:B:699:MET:HE3	1.87	0.56
1:B:352:GLU:OE2	4:B:901:HOH:O	2.18	0.55
1:B:700:GLN:HE21	1:B:704:GLN:NE2	2.04	0.55
1:A:36:ALA:O	1:A:40:GLN:HG3	2.07	0.55
1:A:575:ASN:O	1:A:576:LEU:HB2	2.07	0.54
1:A:628:ARG:NH2	4:A:909:HOH:O	2.40	0.54
1:A:125:LEU:HA	1:A:188:ASN:HD22	1.73	0.53
1:B:696:SER:HA	1:B:699:MET:HE2	1.91	0.53
1:B:82:ASN:HD22	1:B:84:HIS:CE1	2.28	0.52
1:A:374:HIS:HD2	4:A:1589:HOH:O	1.92	0.51
1:A:144:ASN:HB2	4:A:1490:HOH:O	2.10	0.51
1:B:498:GLU:HB3	1:B:504:THR:HG21	1.93	0.50
1:A:593:VAL:CG2	1:B:217:PRO:HG3	2.42	0.50
1:B:410:TYR:CE2	1:B:528:VAL:HA	2.47	0.50
1:B:149:ARG:HH22	1:B:550:THR:HG22	1.78	0.49
1:A:655:SER:OG	2:A:802:TYR:C	2.51	0.49
1:B:655:SER:HG	2:B:802:TYR:C	2.15	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:53:ASN:ND2	1:B:56:GLY:HA3	2.29	0.48
1:B:98:LYS:HE2	4:B:915:HOH:O	2.13	0.48
1:B:620:LEU:C	1:B:620:LEU:HD23	2.34	0.48
1:B:47:SER:OG	1:B:50:GLN:HG3	2.13	0.47
1:B:552:ARG:HB3	1:B:553:PRO:HD3	1.96	0.47
1:A:368:LYS:N	1:A:369:PRO:CD	2.79	0.46
1:B:216:TRP:CG	1:B:217:PRO:HA	2.51	0.46
1:A:216:TRP:CG	1:A:217:PRO:HA	2.51	0.46
1:A:593:VAL:HG22	1:B:217:PRO:HG3	1.99	0.45
1:B:493:LYS:NZ	4:B:903:HOH:O	2.25	0.45
1:A:651:THR:H	1:A:654:ASN:HD22	1.65	0.45
1:A:275:ASN:ND2	1:A:681:ASN:HD21	2.13	0.45
1:A:36:ALA:O	1:A:40:GLN:CG	2.63	0.45
1:B:526:TYR:CZ	1:B:530:VAL:HG11	2.52	0.44
1:A:64:ALA:HB1	3:A:803:GOL:H32	2.00	0.44
1:B:600:THR:HB	1:B:639:SER:HB3	1.98	0.44
1:B:384:SER:O	1:B:387:THR:N	2.48	0.44
1:A:712:LEU:HA	1:A:712:LEU:HD12	1.71	0.44
1:A:267:VAL:HG13	1:A:657:SER:HB3	2.00	0.44
1:B:64:ALA:HB3	1:B:123:PHE:HB2	2.01	0.43
1:B:374:HIS:CE1	1:B:378:LEU:HD13	2.53	0.43
1:B:144:ASN:N	1:B:144:ASN:HD22	2.16	0.43
1:A:218:ARG:HB3	1:A:643:ASN:HD22	1.84	0.43
1:A:601:LEU:HB3	1:A:624:VAL:HG22	2.01	0.42
1:B:149:ARG:HH22	1:B:550:THR:CG2	2.31	0.42
1:B:513:ARG:NH2	4:B:932:HOH:O	2.49	0.42
1:B:267:VAL:HG13	1:B:657:SER:HB3	2.01	0.42
1:B:172:CYS:HA	1:B:186:PHE:O	2.19	0.42
1:A:700:GLN:HE21	1:A:704:GLN:HE21	1.66	0.42
1:B:368:LYS:N	1:B:369:PRO:CD	2.83	0.41
1:A:299:ASN:ND2	4:A:930:HOH:O	2.51	0.41
1:A:597:PRO:O	1:A:644:PHE:HA	2.20	0.41
1:B:475:LEU:HD12	1:B:475:LEU:HA	1.89	0.41
1:B:191:ILE:HG21	1:B:228:ALA:HB1	2.03	0.41
1:B:597:PRO:O	1:B:644:PHE:HA	2.20	0.41
1:A:406:ALA:HB2	1:A:499:LEU:CD1	2.51	0.41
1:B:384:SER:O	1:B:386:SER:N	2.54	0.41
1:A:219:HIS:HB3	1:A:604:VAL:HG22	2.02	0.41
1:B:49:GLU:O	1:B:52:ALA:HB3	2.21	0.40
1:B:29:PRO:HB3	1:B:54:LEU:HD21	2.04	0.40
1:B:259:LEU:HD22	1:B:667:VAL:HB	2.02	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:677:SER:HB2	1:A:689:THR:HG23	2.03	0.40
1:A:576:LEU:HD22	4:A:1034:HOH:O	2.12	0.40
1:A:719:ALA:HB2	4:A:1248:HOH:O	2.21	0.40
1:B:218:ARG:HB3	1:B:643:ASN:HD22	1.87	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1106:HOH:O	4:A:1238:HOH:O[2_655]	1.85	0.35
4:A:1238:HOH:O	4:A:1525:HOH:O[2_645]	2.16	0.04

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	695/720 (96%)	678 (98%)	17 (2%)	0	100	100
1	B	695/720 (96%)	672 (97%)	22 (3%)	1 (0%)	51	36
All	All	1390/1440 (96%)	1350 (97%)	39 (3%)	1 (0%)	51	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	385	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	540/557 (97%)	533 (99%)	7 (1%)	69	58
1	B	540/557 (97%)	536 (99%)	4 (1%)	84	79
All	All	1080/1114 (97%)	1069 (99%)	11 (1%)	76	69

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	GLU
1	A	40	GLN
1	A	247	GLN
1	A	381	LEU
1	A	382	ASP
1	A	553	PRO
1	A	714	LYS
1	B	376	GLN
1	B	382	ASP
1	B	507	LYS
1	B	635	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (21) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	76	GLN
1	A	188	ASN
1	A	275	ASN
1	A	299	ASN
1	A	455	GLN
1	A	584	ASN
1	A	643	ASN
1	A	654	ASN
1	A	704	GLN
1	B	82	ASN
1	B	144	ASN
1	B	188	ASN
1	B	213	ASN
1	B	328	ASN
1	B	376	GLN
1	B	441	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	584	ASN
1	B	643	ASN
1	B	654	ASN
1	B	704	GLN
1	B	717	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	TYR	B	801	2	11,12,13	0.46	0	12,15,17	0.77	0
3	GOL	A	804	-	5,5,5	0.18	0	5,5,5	0.89	0
3	GOL	A	806	-	5,5,5	0.19	0	5,5,5	0.36	0
2	TYR	A	802	2	10,13,13	0.41	0	12,17,17	0.83	0
3	GOL	B	804	-	5,5,5	0.12	0	5,5,5	0.36	0
3	GOL	A	803	-	5,5,5	0.23	0	5,5,5	0.70	0
2	TYR	A	801	2	11,12,13	0.54	0	12,15,17	1.09	0
2	TYR	B	802	2	10,13,13	0.49	0	12,17,17	0.84	1 (8%)
3	GOL	A	805	-	5,5,5	0.11	0	5,5,5	0.41	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	B	803	-	5,5,5	0.13	0	5,5,5	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TYR	B	801	2	-	0/5/6/8	0/1/1/1
3	GOL	A	804	-	-	4/4/4/4	-
3	GOL	A	806	-	-	2/4/4/4	-
2	TYR	A	802	2	-	2/4/8/8	0/1/1/1
3	GOL	B	804	-	-	2/4/4/4	-
3	GOL	A	803	-	-	2/4/4/4	-
2	TYR	A	801	2	-	0/5/6/8	0/1/1/1
2	TYR	B	802	2	-	2/4/8/8	0/1/1/1
3	GOL	A	805	-	-	0/4/4/4	-
3	GOL	B	803	-	-	0/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	802	TYR	CB-CA-C	2.63	114.95	110.69

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	803	GOL	O1-C1-C2-O2
3	A	803	GOL	O1-C1-C2-C3
3	A	804	GOL	C1-C2-C3-O3
3	A	806	GOL	C1-C2-C3-O3
3	A	806	GOL	O2-C2-C3-O3
3	B	804	GOL	O1-C1-C2-C3
3	B	804	GOL	O1-C1-C2-O2
3	A	804	GOL	O2-C2-C3-O3
3	A	804	GOL	O1-C1-C2-O2
2	B	802	TYR	CA-CB-CG-CD2
2	B	802	TYR	CA-CB-CG-CD1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	802	TYR	CA-CB-CG-CD1
2	A	802	TYR	CA-CB-CG-CD2
3	A	804	GOL	O1-C1-C2-C3

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	802	TYR	1	0
3	A	803	GOL	1	0
2	B	802	TYR	2	0
3	A	805	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	697/720 (96%)	-0.27	6 (0%) 84 84	12, 21, 38, 101	0
1	B	697/720 (96%)	0.20	24 (3%) 45 42	16, 33, 51, 105	0
All	All	1394/1440 (96%)	-0.04	30 (2%) 62 61	12, 26, 48, 105	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	383	THR	7.4
1	B	168	ALA	5.6
1	B	383	THR	5.1
1	B	381	LEU	3.8
1	B	719	ALA	3.8
1	A	381	LEU	3.6
1	B	382	ASP	3.6
1	B	231	GLY	3.2
1	B	68	CYS	3.2
1	B	385	LYS	3.2
1	B	378	LEU	3.0
1	B	376	GLN	2.8
1	A	364	GLY	2.7
1	B	69	THR	2.7
1	B	142	ALA	2.6
1	B	167	ASP	2.6
1	B	234	GLY	2.6
1	A	382	ASP	2.6
1	B	653	GLY	2.5
1	B	85	CYS	2.4
1	A	365	ALA	2.3
1	A	484	ALA	2.3
1	B	379	LYS	2.2
1	B	146	PRO	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	365	ALA	2.2
1	B	618	LYS	2.1
1	B	417	SER	2.1
1	B	67	GLY	2.1
1	B	162	ALA	2.1
1	B	511	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	B	803	6/6	0.81	0.26	42,47,52,55	0
3	GOL	B	804	6/6	0.89	0.21	52,56,57,59	0
3	GOL	A	806	6/6	0.91	0.19	26,38,43,57	0
3	GOL	A	805	6/6	0.93	0.16	27,44,54,54	0
3	GOL	A	803	6/6	0.93	0.14	23,30,36,37	0
2	TYR	B	801	12/13	0.94	0.15	19,25,30,41	0
3	GOL	A	804	6/6	0.94	0.13	24,29,34,35	0
2	TYR	B	802	13/13	0.94	0.16	20,22,27,29	0
2	TYR	A	801	12/13	0.97	0.12	14,19,23,30	0
2	TYR	A	802	13/13	0.97	0.11	15,18,21,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.