



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 01:25 am BST

PDB ID : 1DLF  
Title : HIGH RESOLUTION CRYSTAL STRUCTURE OF THE FV FRAGMENT  
FROM AN ANTI-DANSYL SWITCH VARIANT ANTIBODY IGG2A(S)  
CRYSTALLIZED AT PH 5.25  
Authors : Nakasako, M.; Takahashi, H.; Shimada, I.; Arata, Y.  
Deposited on : 1998-07-14  
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1202 (1.46-1.46)
Ramachandran outliers	138981	1178 (1.46-1.46)
Sidechain outliers	138945	1178 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	L	113	
2	H	124	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2394 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ANTI-DANSYL IMMUNOGLOBULIN IGG2A(S).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	113	Total	C	N	O	S	0	4	0
			890	560	153	174	3			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	17	ASN	ASP	CONFLICT	UNP Q9JL82
L	96	PHE	TYR	CONFLICT	UNP Q9JL82
L	100	SER	GLY	CONFLICT	UNP Q9JL82

- Molecule 2 is a protein called ANTI-DANSYL IMMUNOGLOBULIN IGG2A(S).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	120	Total	C	N	O	S	0	12	0
			1032	658	172	197	5			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	24	THR	ALA	CONFLICT	PIR PC1213
H	49	ALA	GLY	CONFLICT	PIR PC1213
H	52A	ASN	SER	CONFLICT	PIR PC1213
H	76	ARG	SER	CONFLICT	PIR PC1213
H	77	ARG	SER	CONFLICT	PIR PC1213
H	82B	THR	SER	CONFLICT	PIR PC1213
H	94	GLY	PRO	CONFLICT	PIR PC1213
H	96	TYR	-	INSERTION	PIR PC1213
H	97	TYR	THR	CONFLICT	PIR PC1213
H	98	HIS	THR	CONFLICT	PIR PC1213
H	99	TYR	GLY	CONFLICT	PIR PC1213
H	100	PRO	ALA	CONFLICT	PIR PC1213

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	H	1	Total	O	S	0	0
			5	4	1		
3	H	1	Total	O	S	0	0
			5	4	1		
3	H	1	Total	O	S	0	0
			5	4	1		
3	H	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	L	221	Total	O	0	0
			221	221		
4	H	231	Total	O	0	0
			231	231		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

- Molecule 1: ANTI-DANSYL IMMUNOGLOBULIN IGG2A(S)

Chain L:  92% 8%



- Molecule 2: ANTI-DANSYL IMMUNOGLOBULIN IGG2A(S)

Chain H:  82% 13% . .



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.10 Å 70.72 Å 53.46 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 1.45	Depositor
% Data completeness (in resolution range)	90.8 (8.00-1.45)	Depositor
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.07	Depositor
Refinement program	X-PLOR 3.0	Depositor
R, $R_{free}$	0.183 , 0.230	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2394	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	L	0.43	0/926	0.58	0/1252
2	H	0.42	0/1112	0.59	0/1504
All	All	0.42	0/2038	0.58	0/2756

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	H	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	H	58[A]	TYR	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	890	0	878	9	0
2	H	1032	0	976	20	0
3	H	20	0	0	0	0
4	H	231	0	0	9	0
4	L	221	0	0	7	0
All	All	2394	0	1854	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:42[B]:GLN:OE1	4:L:282:HOH:O	1.74	1.04
2:H:13[C]:GLN:NE2	4:H:691:HOH:O	1.97	0.96
2:H:13[B]:GLN:NE2	4:H:700:HOH:O	2.09	0.84
4:L:241:HOH:O	2:H:58[B]:TYR:CE2	2.30	0.84
2:H:13[B]:GLN:CD	4:H:700:HOH:O	2.16	0.82
2:H:52(A)[B]:ASN:HD21	2:H:53:ASN:HD22	1.26	0.82
2:H:13[B]:GLN:OE1	4:H:700:HOH:O	2.04	0.75
4:L:241:HOH:O	2:H:58[B]:TYR:HE2	1.68	0.71
1:L:74:LYS:HE2	4:L:182:HOH:O	1.94	0.66
2:H:3[B]:LYS:HE3	4:H:721:HOH:O	1.96	0.65
2:H:55:HIS:CD2	2:H:71:ARG:HD3	2.32	0.65
1:L:12:PRO:HB2	1:L:107:LYS:HE3	1.82	0.60
1:L:72[B]:THR:HG22	4:L:173:HOH:O	2.04	0.57
2:H:52(A)[B]:ASN:ND2	2:H:53:ASN:HD22	1.98	0.57
2:H:42[B]:GLU:OE2	4:H:633:HOH:O	2.17	0.54
2:H:107[B]:GLN:HG3	4:H:621:HOH:O	2.09	0.50
1:L:65:SER:OG	1:L:72[B]:THR:HG23	2.12	0.49
1:L:72[B]:THR:CG2	4:L:173:HOH:O	2.60	0.49
2:H:12:VAL:HG11	2:H:82(C):LEU:HD13	1.96	0.47
2:H:41:PRO:HG3	4:H:629:HOH:O	2.13	0.47
1:L:12:PRO:HB2	1:L:107:LYS:HG2	1.95	0.47
1:L:30:ASN:N	1:L:30:ASN:HD22	2.14	0.45
2:H:97:TYR:CG	2:H:98:HIS:N	2.85	0.45
2:H:83:ARG:NH1	4:H:604:HOH:O	2.50	0.44
2:H:52(A)[B]:ASN:HD21	2:H:53:ASN:ND2	2.06	0.43
2:H:85:GLU:HG2	2:H:85:GLU:O	2.20	0.42
1:L:1:ASP:HB2	4:L:114:HOH:O	2.20	0.41
2:H:75[A]:LYS:HB3	2:H:77:ARG:HG3	2.02	0.41



There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	116/113 (103%)	112 (97%)	4 (3%)	0	100	100
2	H	132/124 (106%)	130 (98%)	2 (2%)	0	100	100
All	All	248/237 (105%)	242 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	106/101 (105%)	105 (99%)	1 (1%)	78	57
2	H	114/103 (111%)	113 (99%)	1 (1%)	78	57
All	All	220/204 (108%)	218 (99%)	2 (1%)	76	57

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	11	LEU
2	H	85	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	H	54	ASN
2	H	55	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	H	503	-	4,4,4	0.61	0	6,6,6	0.41	0
3	SO4	H	501	-	4,4,4	0.58	0	6,6,6	0.27	0
3	SO4	H	502	-	4,4,4	0.76	0	6,6,6	0.42	0
3	SO4	H	500	-	4,4,4	0.42	0	6,6,6	0.24	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.