



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 21, 2020 – 02:46 PM BST

PDB ID : 6DLE  
Title : Crystal structure of IgLON5 homodimer  
Authors : Ranaivoson, F.M.; Turk, L.S.; Ozkan, E.; Montelione, G.T.; Comoletti, D.  
Deposited on : 2018-06-01  
Resolution : 3.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1087 (4.30-3.70)
Clashscore	141614	1148 (4.30-3.70)
Ramachandran outliers	138981	1108 (4.30-3.70)
Sidechain outliers	138945	1099 (4.30-3.70)
RSRZ outliers	127900	1028 (4.34-3.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	303	<div> <div>20%</div> <div>69%</div> <div>20%</div> <div>11%</div> </div>
1	B	303	<div> <div>5%</div> <div>73%</div> <div>17%</div> <div>11%</div> </div>
2	C	2	<div> <div>50%</div> <div>50%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	B	405	-	-	-	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4031 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IgLON family member 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	271	Total	C	N	O	S	0	0	0
			1956	1220	336	390	10			
1	B	271	Total	C	N	O	S	0	0	0
			1949	1208	340	391	10			

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	20	ASP	-	expression tag	UNP A6NGN9
A	21	TYR	-	expression tag	UNP A6NGN9
A	22	LYS	-	expression tag	UNP A6NGN9
A	23	ASP	-	expression tag	UNP A6NGN9
A	24	ASP	-	expression tag	UNP A6NGN9
A	25	ASP	-	expression tag	UNP A6NGN9
A	26	ASP	-	expression tag	UNP A6NGN9
A	27	LYS	-	expression tag	UNP A6NGN9
A	28	ALA	-	expression tag	UNP A6NGN9
A	29	ALA	-	expression tag	UNP A6NGN9
A	30	ALA	-	expression tag	UNP A6NGN9
A	317	LEU	-	expression tag	UNP A6NGN9
A	318	GLU	-	expression tag	UNP A6NGN9
A	319	VAL	-	expression tag	UNP A6NGN9
A	320	LEU	-	expression tag	UNP A6NGN9
A	321	PHE	-	expression tag	UNP A6NGN9
A	322	GLN	-	expression tag	UNP A6NGN9
B	20	ASP	-	expression tag	UNP A6NGN9
B	21	TYR	-	expression tag	UNP A6NGN9
B	22	LYS	-	expression tag	UNP A6NGN9
B	23	ASP	-	expression tag	UNP A6NGN9
B	24	ASP	-	expression tag	UNP A6NGN9
B	25	ASP	-	expression tag	UNP A6NGN9
B	26	ASP	-	expression tag	UNP A6NGN9
B	27	LYS	-	expression tag	UNP A6NGN9

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	28	ALA	-	expression tag	UNP A6NGN9
B	29	ALA	-	expression tag	UNP A6NGN9
B	30	ALA	-	expression tag	UNP A6NGN9
B	317	LEU	-	expression tag	UNP A6NGN9
B	318	GLU	-	expression tag	UNP A6NGN9
B	319	VAL	-	expression tag	UNP A6NGN9
B	320	LEU	-	expression tag	UNP A6NGN9
B	321	PHE	-	expression tag	UNP A6NGN9
B	322	GLN	-	expression tag	UNP A6NGN9

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: IgLON family member 5



- Molecule 1: IgLON family member 5



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	119.20Å 138.23Å 160.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.90 – 3.99 48.89 – 3.99	Depositor EDS
% Data completeness (in resolution range)	99.4 (48.90-3.99) 99.5 (48.89-3.99)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.01 (at 4.00Å)	Xtriage
Refinement program	PHENIX (1.13 _2998: ???)	Depositor
R, $R_{free}$	0.260 , 0.308 0.263 , 0.311	Depositor DCC
$R_{free}$ test set	1156 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	183.4	Xtriage
Anisotropy	0.268	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 188.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.43$ , $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	4031	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	226.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.59% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.26	0/1995	0.49	0/2742
1	B	0.25	0/1987	0.49	0/2730
All	All	0.26	0/3982	0.49	0/5472

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1956	0	1774	39	1
1	B	1949	0	1744	32	0
2	C	28	0	25	2	0
3	A	42	0	39	4	0
3	B	56	0	52	0	0
All	All	4031	0	3634	69	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:187:ARG:HA	1:A:214:VAL:HG21	1.80	0.64
1:B:263:GLU:O	1:B:285:HIS:NE2	2.31	0.62
1:B:129:VAL:O	1:B:159:ARG:NH2	2.31	0.61
1:A:239:GLU:HG2	1:A:273:ARG:HG2	1.82	0.61
1:B:107:GLU:OE1	1:B:159:ARG:NH2	2.31	0.61
1:A:76:ASN:OD1	1:A:90:ASN:ND2	2.34	0.61
1:B:56:ILE:HB	1:B:60:VAL:HG21	1.83	0.58
1:A:259:SER:HB2	1:A:267:VAL:HG22	1.86	0.58
1:A:236:LEU:HD12	1:A:276:LEU:HD23	1.86	0.57
1:B:259:SER:HB2	1:B:267:VAL:HG22	1.85	0.57
1:A:43:THR:HG22	1:A:128:ILE:HB	1.87	0.57
1:A:144:VAL:HG11	1:A:150:VAL:HG22	1.86	0.56
1:A:190:ALA:HB2	1:A:214:VAL:HG22	1.87	0.56
1:B:132:PRO:HA	1:B:199:ASN:HD21	1.69	0.56
1:A:154:CYS:HB2	1:A:166:TRP:CZ2	2.41	0.56
1:B:231:GLY:H	1:B:281:VAL:HG21	1.70	0.56
1:A:131:VAL:O	1:A:199:ASN:ND2	2.41	0.54
1:A:167:ARG:HG3	1:A:174:THR:HG22	1.88	0.54
1:B:54:CYS:HB2	1:B:65:TRP:CZ2	2.43	0.54
1:A:132:PRO:HA	1:A:199:ASN:HD21	1.74	0.53
1:A:62:ARG:HH21	1:B:77:ASP:HB3	1.74	0.53
1:B:144:VAL:HG11	1:B:150:VAL:HG22	1.90	0.53
1:B:226:ALA:O	1:B:306:LEU:HA	2.09	0.52
1:A:40:ASP:HA	3:A:403:NAG:H82	1.90	0.52
1:A:68:ARG:HE	3:A:401:NAG:H62	1.75	0.52
1:A:40:ASP:OD1	3:A:403:NAG:H82	2.11	0.51
1:A:135:ILE:HD12	1:A:208:ARG:HG3	1.93	0.51
1:B:251:TYR:HB2	1:B:290:THR:OG1	2.11	0.50
1:B:187:ARG:HA	1:B:214:VAL:HG21	1.94	0.50
1:A:61:THR:HG22	1:A:62:ARG:HH11	1.77	0.50
1:A:54:CYS:HB2	1:A:65:TRP:CZ2	2.47	0.50
1:B:43:THR:HG22	1:B:128:ILE:HB	1.94	0.50
1:B:249:GLN:O	1:B:292:ARG:N	2.38	0.49
1:A:218:PRO:HB3	1:A:242:ALA:HB2	1.93	0.49
1:A:230:LEU:HG	1:A:281:VAL:HG21	1.95	0.49
1:B:231:GLY:H	1:B:281:VAL:HG11	1.78	0.48
1:A:137:ASN:HB3	1:A:155:LEU:HB3	1.95	0.48
1:A:68:ARG:NH2	1:A:107:GLU:O	2.47	0.48
1:A:91:THR:HG22	1:A:93:GLU:H	1.79	0.47
1:A:45:CYS:HB3	1:A:48:ASP:OD2	2.14	0.47
1:B:61:THR:HG22	1:B:62:ARG:HG2	1.95	0.47
1:B:45:CYS:HB3	1:B:48:ASP:OD2	2.16	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:67:ASN:OD1	2:C:1:NAG:N2	2.50	0.45
1:B:110:TYR:CE1	2:C:1:NAG:H81	2.51	0.45
1:A:63:VAL:HG23	1:A:114:PHE:HB3	1.99	0.45
1:B:154:CYS:HB2	1:B:166:TRP:CZ2	2.52	0.45
1:A:251:TYR:HA	1:A:256:LEU:HA	1.99	0.45
1:B:230:LEU:HG	1:B:281:VAL:HG21	1.99	0.44
1:A:68:ARG:NE	3:A:401:NAG:H62	2.33	0.44
1:B:290:THR:HG22	1:B:303:SER:CB	2.48	0.44
1:A:91:THR:C	1:A:93:GLU:H	2.21	0.43
1:B:185:ILE:HG13	1:B:189:GLN:OE1	2.18	0.43
1:A:304:MET:N	1:A:304:MET:SD	2.92	0.43
1:A:61:THR:HG22	1:A:62:ARG:HG2	2.00	0.43
1:B:185:ILE:HG13	1:B:189:GLN:HB2	2.00	0.43
1:A:263:GLU:O	1:A:285:HIS:NE2	2.38	0.43
1:A:162:PRO:HB2	1:A:197:THR:OG1	2.19	0.43
1:A:224:THR:OG1	1:A:237:ARG:HG2	2.19	0.43
1:B:68:ARG:NH2	1:B:107:GLU:O	2.52	0.43
1:B:216:TYR:H	1:B:243:VAL:HG23	1.83	0.43
1:A:230:LEU:HA	1:A:281:VAL:HG11	2.01	0.42
1:B:290:THR:HG22	1:B:303:SER:HB3	2.01	0.42
1:B:215:ASN:HB3	1:B:297:LEU:HD11	2.00	0.42
1:A:285:HIS:O	1:A:289:TYR:OH	2.20	0.42
1:B:231:GLY:N	1:B:281:VAL:HG11	2.34	0.42
1:A:62:ARG:HB3	1:B:79:TRP:CZ2	2.55	0.42
1:A:56:ILE:HB	1:A:60:VAL:HG21	2.03	0.41
1:B:67:ASN:N	1:B:70:ASN:O	2.45	0.41
1:A:106:ASP:O	1:A:127:LEU:HD23	2.20	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:126:TYR:OH	1:A:299:ALA:O[8_455]	1.98	0.22

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	269/303 (89%)	259 (96%)	10 (4%)	0	100	100
1	B	269/303 (89%)	258 (96%)	11 (4%)	0	100	100
All	All	538/606 (89%)	517 (96%)	21 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/258 (75%)	191 (98%)	3 (2%)	65	80
1	B	191/258 (74%)	188 (98%)	3 (2%)	62	79
All	All	385/516 (75%)	379 (98%)	6 (2%)	62	79

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	73	TYR
1	A	122	THR
1	A	291	CYS
1	B	115	GLN
1	B	227	ARG
1	B	291	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	C	1	1,2	14,14,15	0.56	0	17,19,21	0.61	0
2	NAG	C	2	2	14,14,15	0.38	0	17,19,21	0.59	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	1/6/23/26	0/1/1/1
2	NAG	C	2	2	-	3/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	1	NAG	O5-C5-C6-O6
2	C	2	NAG	C4-C5-C6-O6
2	C	2	NAG	O5-C5-C6-O6
2	C	2	NAG	C3-C2-N2-C7

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	NAG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	A	401	1	14,14,15	0.37	0	17,19,21	0.71	0
3	NAG	B	406	1	14,14,15	0.25	0	17,19,21	0.49	0
3	NAG	B	405	1	14,14,15	0.32	0	17,19,21	0.36	0
3	NAG	A	403	1	14,14,15	0.66	1 (7%)	17,19,21	0.73	1 (5%)
3	NAG	B	403	1	14,14,15	0.24	0	17,19,21	0.65	1 (5%)
3	NAG	A	402	1	14,14,15	0.29	0	17,19,21	0.63	0
3	NAG	B	404	1	14,14,15	0.34	0	17,19,21	0.63	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	401	1	-	2/6/23/26	0/1/1/1
3	NAG	B	406	1	-	3/6/23/26	0/1/1/1
3	NAG	B	405	1	-	0/6/23/26	0/1/1/1
3	NAG	A	403	1	-	0/6/23/26	0/1/1/1
3	NAG	B	403	1	-	0/6/23/26	0/1/1/1
3	NAG	A	402	1	-	3/6/23/26	0/1/1/1
3	NAG	B	404	1	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	403	NAG	C1-C2	2.24	1.55	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	403	NAG	C1-O5-C5	2.25	115.25	112.19
3	A	403	NAG	C1-O5-C5	2.25	115.24	112.19
3	B	404	NAG	C1-O5-C5	2.16	115.12	112.19

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	402	NAG	O5-C5-C6-O6
3	B	406	NAG	C3-C2-N2-C7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	B	406	NAG	C4-C5-C6-O6
3	A	401	NAG	C1-C2-N2-C7
3	A	402	NAG	C1-C2-N2-C7
3	A	401	NAG	C3-C2-N2-C7
3	A	402	NAG	C3-C2-N2-C7
3	B	406	NAG	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	401	NAG	2	0
3	A	403	NAG	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	271/303 (89%)	1.53	60 (22%) 0 1	121, 219, 475, 592	0
1	B	271/303 (89%)	0.33	14 (5%) 27 24	129, 190, 249, 278	0
All	All	542/606 (89%)	0.93	74 (13%) 3 3	121, 200, 427, 592	0

All (74) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	226	ALA	21.1
1	A	234	ALA	16.7
1	A	236	LEU	13.3
1	A	225	SER	12.7
1	A	278	PHE	12.1
1	A	276	LEU	11.4
1	A	223	VAL	11.1
1	A	233	ALA	10.6
1	A	224	THR	8.5
1	A	235	LEU	8.2
1	A	289	TYR	7.3
1	A	238	CYS	6.1
1	A	280	ASN	6.1
1	A	279	ALA	5.9
1	A	240	ALA	5.8
1	A	227	ARG	5.7
1	A	250	TRP	5.6
1	A	290	THR	5.5
1	A	197	THR	5.4
1	B	214	VAL	5.3
1	A	237	ARG	5.2
1	A	277	LEU	5.2
1	A	267	VAL	5.0
1	A	164	VAL	4.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	274	SER	4.6
1	B	193	TYR	4.3
1	A	287	GLY	4.3
1	A	196	VAL	4.3
1	A	220	ILE	4.2
1	A	288	ASN	4.2
1	A	38	PRO	4.0
1	B	134	ARG	3.8
1	A	156	ALA	3.8
1	A	135	ILE	3.7
1	A	304	MET	3.7
1	B	167	ARG	3.7
1	A	266	LYS	3.5
1	A	230	LEU	3.5
1	A	264	GLY	3.5
1	B	215	ASN	3.4
1	A	138	ILE	3.3
1	A	302	ALA	3.3
1	A	222	ASP	3.3
1	B	156	ALA	3.1
1	A	144	VAL	3.1
1	A	228	THR	2.9
1	A	210	VAL	2.8
1	A	297	LEU	2.7
1	A	182	ILE	2.7
1	A	137	ASN	2.7
1	A	165	THR	2.7
1	A	265	LEU	2.6
1	A	98	LEU	2.6
1	A	221	THR	2.6
1	A	261	THR	2.6
1	A	303	SER	2.5
1	B	122	THR	2.5
1	B	114	PHE	2.5
1	A	306	LEU	2.5
1	B	133	ALA	2.5
1	A	97	ILE	2.4
1	A	166	TRP	2.4
1	A	69	SER	2.3
1	A	263	GLU	2.3
1	A	283	ALA	2.3
1	A	195	CYS	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	138	ILE	2.3
1	B	226	ALA	2.2
1	B	197	THR	2.2
1	B	276	LEU	2.1
1	A	285	HIS	2.1
1	A	232	ARG	2.1
1	B	223	VAL	2.1
1	A	231	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAG	C	1	14/15	0.67	0.39	179,219,290,295	0
2	NAG	C	2	14/15	0.86	0.32	210,249,299,307	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	B	405	14/15	0.41	0.72	184,278,300,315	0
3	NAG	A	402	14/15	0.68	0.35	171,209,247,270	0
3	NAG	B	403	14/15	0.77	0.31	214,254,295,318	0
3	NAG	A	403	14/15	0.80	0.29	239,246,263,275	0
3	NAG	A	401	14/15	0.81	0.34	61,197,238,258	0
3	NAG	B	404	14/15	0.84	0.18	114,240,255,264	0
3	NAG	B	406	14/15	0.89	0.25	186,221,255,255	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.