



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 10:23 am BST

PDB ID : 6DP8
Title : Crystal Structure of Bacillus Halodurans Ribonuclease H1 in Complex with an RNA/DNA Hybrid: Reaction in 5 mM Mg²⁺ and 200 mM Li⁺ for 240 s at 21 C
Authors : Samara, N.L.; Yang, W.
Deposited on : 2018-06-09
Resolution : 1.32 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1611 (1.34-1.30)
Clashscore	141614	1667 (1.34-1.30)
Ramachandran outliers	138981	1615 (1.34-1.30)
Sidechain outliers	138945	1615 (1.34-1.30)
RSRZ outliers	127900	1580 (1.34-1.30)
RNA backbone	3102	1000 (2.34-0.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	142	<div> <div>4%</div> <div>84%</div> <div>12%</div> <div>.</div> </div>
2	B	4	<div> <div>75%</div> <div>25%</div> </div>
3	b	2	<div> <div>50%</div> <div>50%</div> </div>
4	C	6	<div> <div>100%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	PEG	A	210	-	-	X	-
7	CL	A	203	-	-	X	-
8	GOL	A	212	-	-	X	-

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 2551 atoms, of which 895 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribonuclease H.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	136	Total	C	H	N	O	S	0	6	0
			2037	733	895	189	218	2			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	55	GLY	-	expression tag	UNP Q9KEI9
A	56	SER	-	expression tag	UNP Q9KEI9
A	57	HIS	-	expression tag	UNP Q9KEI9
A	58	MET	-	expression tag	UNP Q9KEI9

- Molecule 2 is a RNA chain called RNA (5'-R(*AP*CP*AP*U)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	4	Total	C	N	O	P	0	3	0
			80	38	15	24	3			

- Molecule 3 is a RNA chain called RNA (5'-R(P*CP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	b	2	Total	C	N	O	P	0	0	0
			43	19	8	14	2			

- Molecule 4 is a DNA chain called DNA (5'-D(*CP*GP*AP*TP*GP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	C	6	Total	C	N	O	P	0	0	0
			121	59	22	35	5			

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by author).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	b	1	Total Mg 1 1	0	0
5	A	1	Total Mg 1 1	0	0


- Molecule 6 is LITHIUM ION (three-letter code: LI) (formula: Li) (labeled as "Ligand of Interest" by author).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Li 1 1	0	0

- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	3	Total Cl 3 3	0	0

- Molecule 8 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total C O 6 3 3	0	0
8	A	1	Total C O 6 3 3	0	0
8	A	1	Total C O 6 3 3	0	0

- Molecule 9 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	C	O	0	0
			4	2	2		
9	A	1	Total	C	O	0	0
			4	2	2		
9	A	1	Total	C	O	0	0
			4	2	2		
9	A	1	Total	C	O	0	0
			4	2	2		
9	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 10 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	C	O	0	0
			7	4	3		
10	A	1	Total	C	O	0	0
			7	4	3		


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	164	Total	O	0	0
			164	164		
11	B	15	Total	O	0	0
			15	15		
11	b	8	Total	O	0	0
			8	8		
11	C	21	Total	O	0	0
			21	21		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ribonuclease H



- Molecule 2: RNA (5'-R(*AP*CP*AP*U)-3')



- Molecule 3: RNA (5'-R(P*CP*G)-3')



- Molecule 4: DNA (5'-D(*CP*GP*AP*TP*GP*T)-3')



There are no outlier residues recorded for this chain.

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	81.43Å 37.75Å 61.81Å 90.00° 95.45° 90.00°	Depositor
Resolution (Å)	23.45 – 1.32 23.46 – 1.32	Depositor EDS
% Data completeness (in resolution range)	95.6 (23.45-1.32) 95.3 (23.46-1.32)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.36 (at 1.32Å)	Xtriage
Refinement program	PHENIX 1.10_2155	Depositor
R, R_{free}	0.147 , 0.170 0.150 , 0.173	Depositor DCC
R_{free} test set	2081 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	16.8	Xtriage
Anisotropy	0.216	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 52.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	2551	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, CL, EDO, LI, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/1183	0.60	0/1599
2	B	0.70	0/89	1.23	1/136 (0.7%)
3	b	1.60	1/47 (2.1%)	1.03	0/69
4	C	0.92	0/135	1.02	0/207
All	All	0.55	1/1454 (0.1%)	0.73	1/2011 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	b	5	C	OP3-P	-9.56	1.49	1.61

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	3[C]	A	N7-C8-N9	-5.33	111.13	113.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1142	895	1145	18	0
2	B	80	0	44	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	b	43	0	23	0	0
4	C	121	0	70	0	0
5	A	1	0	0	0	0
5	b	1	0	0	0	0
6	A	1	0	0	0	0
7	A	3	0	0	3	0
8	A	18	0	24	12	0
9	A	20	0	30	6	0
9	B	4	0	6	0	0
10	A	14	0	20	6	0
11	A	164	0	0	7	1
11	B	15	0	0	0	0
11	C	21	0	0	0	0
11	b	8	0	0	0	0
All	All	1656	895	1362	32	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:203:CL:CL	8:A:212:GOL:O3	2.12	1.04
1:A:102:ILE:HD12	1:A:154:GLU:HB3	1.52	0.89
8:A:206:GOL:O3	8:A:206:GOL:O1	1.89	0.83
9:A:215:EDO:H12	11:A:420:HOH:O	1.84	0.78
1:A:96:GLU:OE2	11:A:301:HOH:O	2.04	0.75
10:A:210:PEG:H41	10:A:210:PEG:O1	1.89	0.73
8:A:212:GOL:H32	11:A:367:HOH:O	1.88	0.72
9:A:213:EDO:H22	11:A:371:HOH:O	1.88	0.71
8:A:212:GOL:H11	11:A:351:HOH:O	1.94	0.68
1:A:118:ARG:NH1	8:A:212:GOL:O1	2.31	0.63
8:A:212:GOL:C3	11:A:304:HOH:O	2.47	0.61
7:A:205:CL:CL	11:A:464:HOH:O	2.53	0.60
7:A:203:CL:CL	8:A:212:GOL:C3	2.86	0.59
1:A:150:VAL:CG2	10:A:210:PEG:H11	2.35	0.56
1:A:61:GLU:HG3	1:A:182:GLN:HE22	1.70	0.56
1:A:115:HIS:ND1	8:A:212:GOL:H12	2.23	0.53
1:A:151:ARG:O	10:A:210:PEG:H12	2.08	0.53
1:A:115:HIS:ND1	8:A:212:GOL:C1	2.75	0.49
1:A:118:ARG:CZ	8:A:212:GOL:O1	2.61	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:127:LYS:NZ	9:A:213:EDO:O2	2.33	0.49
1:A:157:LEU:O	1:A:157:LEU:HD12	2.14	0.47
10:A:210:PEG:H41	10:A:210:PEG:C1	2.43	0.46
10:A:209:PEG:H22	10:A:209:PEG:H42	1.72	0.46
1:A:89:LYS:HE2	1:A:89:LYS:HB3	1.64	0.45
1:A:118:ARG:NH1	8:A:212:GOL:HO1	2.14	0.44
8:A:206:GOL:HO1	8:A:206:GOL:HO3	1.57	0.44
1:A:97:ARG:HH22	9:A:211:EDO:H11	1.82	0.44
1:A:97:ARG:HH12	9:A:211:EDO:H11	1.84	0.42
1:A:182:GLN:OE1	1:A:185:LYS:HD2	2.20	0.42
1:A:150:VAL:HG12	10:A:209:PEG:H22	2.03	0.41
1:A:103:GLY:HA3	1:A:158:ILE:HD13	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
11:A:422:HOH:O	11:A:422:HOH:O[2_656]	1.97	0.23

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/142 (98%)	139 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	125/124 (101%)	124 (99%)	1 (1%)	81	57

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	62	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	2/4 (50%)	0	0
3	b	0/2	-	-
All	All	2/6 (33%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 17 ligands modelled in this entry, 6 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
9	EDO	A	213	-	3,3,3	0.48	0	2,2,2	0.33	0
8	GOL	A	207	-	5,5,5	0.34	0	5,5,5	0.63	0
9	EDO	B	101	-	3,3,3	0.47	0	2,2,2	0.38	0
9	EDO	A	208	-	3,3,3	0.42	0	2,2,2	0.42	0
9	EDO	A	211	-	3,3,3	0.50	0	2,2,2	0.26	0
9	EDO	A	215	-	3,3,3	0.46	0	2,2,2	0.24	0
10	PEG	A	209	-	6,6,6	0.50	0	5,5,5	0.31	0
8	GOL	A	212	-	5,5,5	0.41	0	5,5,5	1.42	1 (20%)
8	GOL	A	206	-	5,5,5	0.34	0	5,5,5	0.38	0
10	PEG	A	210	-	6,6,6	0.50	0	5,5,5	0.19	0
9	EDO	A	214	-	3,3,3	0.47	0	2,2,2	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	EDO	A	213	-	-	1/1/1/1	-
8	GOL	A	207	-	-	1/4/4/4	-
9	EDO	B	101	-	-	0/1/1/1	-
9	EDO	A	208	-	-	0/1/1/1	-
9	EDO	A	211	-	-	0/1/1/1	-
9	EDO	A	215	-	-	0/1/1/1	-
10	PEG	A	209	-	-	4/4/4/4	-
8	GOL	A	212	-	-	3/4/4/4	-
8	GOL	A	206	-	-	4/4/4/4	-
10	PEG	A	210	-	-	4/4/4/4	-
9	EDO	A	214	-	-	1/1/1/1	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	212	GOL	C3-C2-C1	-3.05	99.86	111.70

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	A	206	GOL	O1-C1-C2-C3
8	A	206	GOL	C1-C2-C3-O3
10	A	209	PEG	O1-C1-C2-O2
10	A	210	PEG	O2-C3-C4-O4
8	A	212	GOL	O1-C1-C2-C3
8	A	206	GOL	O1-C1-C2-O2
8	A	206	GOL	O2-C2-C3-O3
10	A	209	PEG	O2-C3-C4-O4
10	A	210	PEG	C4-C3-O2-C2
10	A	209	PEG	C4-C3-O2-C2
9	A	213	EDO	O1-C1-C2-O2
8	A	212	GOL	O1-C1-C2-O2
10	A	210	PEG	C1-C2-O2-C3
10	A	210	PEG	O1-C1-C2-O2
10	A	209	PEG	C1-C2-O2-C3
9	A	214	EDO	O1-C1-C2-O2
8	A	207	GOL	O2-C2-C3-O3
8	A	212	GOL	C1-C2-C3-O3

There are no ring outliers.

8 monomers are involved in 24 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	A	213	EDO	2	0
9	A	208	EDO	1	0
9	A	211	EDO	2	0
9	A	215	EDO	1	0
10	A	209	PEG	2	0
8	A	212	GOL	10	0
8	A	206	GOL	2	0
10	A	210	PEG	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	136/142 (95%)	-0.04	5 (3%) 41 42	13, 20, 39, 59	1 (0%)
2	B	4/4 (100%)	-0.34	0 100 100	17, 18, 23, 25	0
3	b	2/2 (100%)	-0.47	0 100 100	17, 17, 17, 21	0
4	C	6/6 (100%)	-0.32	0 100 100	17, 21, 26, 31	0
All	All	148/154 (96%)	-0.07	5 (3%) 45 46	13, 20, 38, 59	1 (0%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	196	LYS	5.2
1	A	61	GLU	3.9
1	A	194	GLY	2.4
1	A	195[A]	ARG	2.2
1	A	62	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
9	EDO	A	211	4/4	0.46	0.28	37,68,77,78	0
8	GOL	A	206	6/6	0.77	0.15	49,55,68,76	0
10	PEG	A	210	7/7	0.77	0.27	58,59,76,99	0
8	GOL	A	212	6/6	0.84	0.24	31,48,73,84	0
9	EDO	B	101	4/4	0.85	0.11	46,47,49,49	0
9	EDO	A	213	4/4	0.87	0.12	32,44,61,65	0
9	EDO	A	214	4/4	0.87	0.17	34,37,56,74	0
9	EDO	A	215	4/4	0.88	0.18	29,47,69,75	0
8	GOL	A	207	6/6	0.90	0.12	23,28,57,61	0
6	LI	A	202	1/1	0.91	0.66	17,17,17,17	0
9	EDO	A	208	4/4	0.92	0.10	48,49,50,52	0
7	CL	A	203	1/1	0.93	0.09	49,49,49,49	1
7	CL	A	205	1/1	0.95	0.05	50,50,50,50	1
10	PEG	A	209	7/7	0.96	0.18	29,42,102,110	0
5	MG	b	101	1/1	0.99	0.07	19,19,19,19	0
7	CL	A	204	1/1	0.99	0.04	32,32,32,32	1
5	MG	A	201	1/1	1.00	0.03	17,17,17,17	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around LI A 202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MG b 101:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MG A 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.