



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 9, 2020 – 09:17 PM BST

PDB ID : 2DTX  
Title : Structure of Thermoplasma acidophilum aldohexose dehydrogenase (AldT) in complex with D-mannose  
Authors : Yasutake, Y.; Nishiya, Y.; Tamura, N.; Tamura, T.  
Deposited on : 2006-07-18  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	264	<div> <div>6%</div> <div>87%</div> <div>9%</div> </div>
1	B	264	<div> <div>7%</div> <div>84%</div> <div>12%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4356 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glucose 1-dehydrogenase related protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	255	Total	C	N	O	S	0	2	0
			1964	1241	335	378	10			
1	B	255	Total	C	N	O	S	0	4	0
			1981	1250	340	381	10			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	expression tag	UNP Q9HK51
A	1	GLY	-	expression tag	UNP Q9HK51
A	256	LEU	-	expression tag	UNP Q9HK51
A	257	GLU	-	expression tag	UNP Q9HK51
A	258	HIS	-	expression tag	UNP Q9HK51
A	259	HIS	-	expression tag	UNP Q9HK51
A	260	HIS	-	expression tag	UNP Q9HK51
A	261	HIS	-	expression tag	UNP Q9HK51
A	262	HIS	-	expression tag	UNP Q9HK51
A	263	HIS	-	expression tag	UNP Q9HK51
B	0	MET	-	expression tag	UNP Q9HK51
B	1	GLY	-	expression tag	UNP Q9HK51
B	256	LEU	-	expression tag	UNP Q9HK51
B	257	GLU	-	expression tag	UNP Q9HK51
B	258	HIS	-	expression tag	UNP Q9HK51
B	259	HIS	-	expression tag	UNP Q9HK51
B	260	HIS	-	expression tag	UNP Q9HK51
B	261	HIS	-	expression tag	UNP Q9HK51
B	262	HIS	-	expression tag	UNP Q9HK51
B	263	HIS	-	expression tag	UNP Q9HK51

- Molecule 2 is beta-D-mannopyranose (three-letter code: BMA) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			10	6	4		
2	B	1	Total	C	O	0	0
			10	6	4		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	203	Total 203	O 203	0	0
4	B	178	Total 178	O 178	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.88Å 81.88Å 138.28Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.00 – 1.60 38.65 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.8 (40.00-1.60) 99.8 (38.65-1.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.63 (at 1.60Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.198 , 0.221 0.204 , 0.228	Depositor DCC
$R_{free}$ test set	3604 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.9	Xtriage
Anisotropy	0.400	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 47.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.017 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4356	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.41	0/2002	0.64	0/2712
1	B	0.41	0/2019	0.64	0/2734
All	All	0.41	0/4021	0.64	0/5446

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1964	0	1967	25	0
1	B	1981	0	1983	29	0
2	A	10	0	7	2	0
2	B	10	0	7	2	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	203	0	0	7	0
4	B	178	0	0	5	0
All	All	4356	0	3964	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.



All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:302:BMA:C1	4:B:772:HOH:O	1.89	1.19
2:A:301:BMA:C1	4:A:771:HOH:O	1.95	1.13
2:B:302:BMA:C3	4:B:774:HOH:O	2.24	0.84
1:B:46:LYS:HD3	1:B:46:LYS:H	1.43	0.83
1:B:194:PRO:HG2	1:B:195:MET:HE2	1.70	0.72
1:B:194:PRO:HG2	1:B:195:MET:CE	2.26	0.65
1:A:206:HIS:HD2	4:A:775:HOH:O	1.82	0.62
1:B:50:ILE:HD11	1:B:66:HIS:HD2	1.65	0.62
1:B:96:GLU:HG2	4:B:607:HOH:O	2.00	0.61
1:A:183:ARG:HD3	1:A:201:ILE:HD11	1.82	0.61
1:B:197:ILE:O	1:B:201:ILE:HG12	2.01	0.61
1:B:2:PHE:CD2	1:B:230:GLU:HG3	2.36	0.60
1:A:217:GLN:NE2	1:A:217:GLN:H	2.00	0.59
1:B:195:MET:H	1:B:195:MET:HE2	1.68	0.57
1:B:46:LYS:HD3	1:B:46:LYS:N	2.15	0.56
1:A:96:GLU:HG2	4:A:606:HOH:O	2.06	0.55
1:B:25:GLU:HG2	1:B:46:LYS:HE2	1.90	0.53
1:B:46:LYS:CD	1:B:46:LYS:H	2.19	0.52
1:B:181:LEU:HD12	1:B:181:LEU:N	2.24	0.52
1:B:48:ASP:OD1	1:B:71:TYR:OH	2.28	0.52
1:A:184:LYS:O	1:A:184:LYS:HD3	2.10	0.52
1:A:181:LEU:HD23	1:A:181:LEU:O	2.10	0.51
1:A:123:ARG:HA	1:A:123:ARG:HE	1.74	0.51
1:B:2:PHE:HD2	1:B:230:GLU:HG3	1.74	0.51
1:B:175:ALA:HB2	1:B:242:ASP:HB2	1.93	0.51
1:A:143:SER:HB2	4:A:750:HOH:O	2.10	0.51
1:A:88:LYS:HD2	1:B:162:ASP:HB3	1.92	0.51
1:A:193:ASP:OD1	1:A:195:MET:HB3	2.13	0.49
1:A:201:ILE:HG23	1:A:213:ILE:HD13	1.95	0.48
2:A:301:BMA:C3	4:A:773:HOH:O	2.60	0.48
1:A:41:ASP:OD2	1:A:42:PRO:HD2	2.13	0.48
1:B:50:ILE:CD1	1:B:66:HIS:HD2	2.26	0.48
1:A:175:ALA:HB2	1:A:242:ASP:HB2	1.95	0.48
1:A:181:LEU:HD23	1:A:181:LEU:C	2.34	0.48
1:A:123:ARG:HA	1:A:123:ARG:NE	2.29	0.46
1:B:5:LEU:HA	1:B:8:LYS:HD2	1.97	0.46
1:B:134:GLN:HA	1:B:137:ILE:O	2.16	0.46
1:A:216:PRO:HD2	1:A:217:GLN:NE2	2.31	0.46
1:A:215:LYS:HE2	1:A:217:GLN:OE1	2.16	0.45
1:B:166:LEU:HD13	1:B:166:LEU:C	2.38	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:184:LYS:O	1:B:188:LEU:HG	2.16	0.44
1:B:85:SER:OG	1:B:100:ILE:HD11	2.17	0.44
1:A:247:ILE:C	1:A:247:ILE:HD12	2.38	0.43
1:A:90:GLU:OE2	1:B:113:LYS:O	2.37	0.43
1:A:163:TYR:OH	1:B:90:GLU:HG2	2.18	0.43
1:B:201:ILE:HG23	1:B:213:ILE:HD13	2.01	0.42
1:B:247:ILE:HD12	1:B:247:ILE:C	2.40	0.42
1:A:123:ARG:NH2	4:A:526:HOH:O	2.52	0.42
1:A:134:GLN:HA	1:A:137:ILE:O	2.20	0.41
1:B:82:GLY:HA3	4:B:484:HOH:O	2.19	0.41
1:A:149:LYS:NZ	4:A:770:HOH:O	2.45	0.41
1:A:88:LYS:HG2	4:B:686:HOH:O	2.20	0.41
1:B:201:ILE:CG2	1:B:213:ILE:HD13	2.50	0.41
1:A:193:ASP:HA	1:A:194:PRO:HD2	1.95	0.41
1:B:166:LEU:HD13	1:B:166:LEU:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	255/264 (97%)	246 (96%)	9 (4%)	0	100	100
1	B	257/264 (97%)	247 (96%)	10 (4%)	0	100	100
All	All	512/528 (97%)	493 (96%)	19 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	215/222 (97%)	214 (100%)	1 (0%)	88	80
1	B	217/222 (98%)	214 (99%)	3 (1%)	67	47
All	All	432/444 (97%)	428 (99%)	4 (1%)	78	65

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	217	GLN
1	B	44	GLU
1	B	46	LYS
1	B	217	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	40	HIS
1	B	66	HIS
1	B	217	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	SO4	A	351	-	4,4,4	0.33	0	6,6,6	0.30	0
2	BMA	B	302	-	10,10,12	3.02	2 (20%)	12,13,17	2.39	4 (33%)
2	BMA	A	301	-	10,10,12	3.02	2 (20%)	12,13,17	2.39	4 (33%)
3	SO4	B	352	-	4,4,4	0.37	0	6,6,6	0.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BMA	B	302	-	-	2/2/15/22	0/1/1/1
2	BMA	A	301	-	-	2/2/15/22	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	302	BMA	O2-C2	-9.06	1.16	1.43
2	A	301	BMA	O2-C2	-9.05	1.16	1.43
2	A	301	BMA	C1-C2	2.06	1.53	1.51
2	B	302	BMA	C1-C2	2.04	1.53	1.51

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	BMA	O2-C2-C1	5.11	120.23	110.34
2	B	302	BMA	O2-C2-C1	5.10	120.22	110.34
2	A	301	BMA	C3-C4-C5	-3.88	105.74	110.77
2	B	302	BMA	C3-C4-C5	-3.86	105.76	110.77
2	B	302	BMA	O5-C1-C2	-3.81	106.12	110.66
2	A	301	BMA	O5-C1-C2	-3.78	106.15	110.66
2	B	302	BMA	O4-C4-C5	2.37	115.03	110.01
2	A	301	BMA	O4-C4-C5	2.35	115.00	110.01

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	302	BMA	O5-C5-C6-O6
2	A	301	BMA	O5-C5-C6-O6
2	B	302	BMA	C4-C5-C6-O6
2	A	301	BMA	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	302	BMA	2	0
2	A	301	BMA	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	255/264 (96%)	0.20	16 (6%) 20 18	9, 16, 35, 50	0
1	B	255/264 (96%)	0.38	19 (7%) 14 12	10, 19, 39, 52	0
All	All	510/528 (96%)	0.29	35 (6%) 16 15	9, 17, 37, 52	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	194	PRO	7.3
1	A	44	GLU	5.6
1	A	190	VAL	5.5
1	A	195	MET	5.0
1	A	43	GLY	4.9
1	B	44	GLU	4.8
1	A	255	GLU	4.6
1	B	82	GLY	4.2
1	A	123	ARG	4.1
1	A	191	GLY	4.0
1	B	123	ARG	3.8
1	B	192	SER	3.6
1	B	187	GLU	3.5
1	A	181	LEU	3.5
1	B	196	ARG	3.5
1	A	46	LYS	3.5
1	B	255	GLU	3.4
1	B	166	LEU	3.3
1	B	49	HIS	3.3
1	A	194	PRO	3.1
1	B	195	MET	3.0
1	A	192	SER	2.9
1	A	196	ARG	2.9
1	B	69	LYS	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	41	ASP	2.7
1	B	181	LEU	2.7
1	A	42	PRO	2.7
1	B	184	LYS	2.6
1	A	197	ILE	2.6
1	B	197	ILE	2.4
1	B	46	LYS	2.3
1	B	51	GLU	2.3
1	A	91	SER	2.2
1	B	188	LEU	2.1
1	B	191	GLY	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	BMA	A	301	10/12	0.67	0.22	28,33,36,37	10
2	BMA	B	302	10/12	0.73	0.19	27,32,36,36	0
3	SO4	B	352	5/5	0.84	0.17	26,28,32,32	0
3	SO4	A	351	5/5	0.88	0.17	23,27,30,31	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.