



# Full wwPDB X-ray Structure Validation Report ⓘ

May 17, 2020 – 11:12 pm BST

PDB ID : 6DVX  
Title : Citrobacter freundii tyrosine phenol-lyase F448A mutant complexed with L-phenylalanine  
Authors : Phillips, R.S.  
Deposited on : 2018-06-25  
Resolution : 2.27 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6980 (2.30-2.26)
Clashscore	141614	7711 (2.30-2.26)
Ramachandran outliers	138981	7597 (2.30-2.26)
Sidechain outliers	138945	7598 (2.30-2.26)
RSRZ outliers	127900	6849 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	456	<div> <div>13%</div> <div> <div></div> <div>85%</div> <div>13%</div> <div>..</div> </div> </div>
1	B	456	<div> <div>5%</div> <div> <div></div> <div>92%</div> <div>7%</div> </div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7794 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine phenol-lyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	451	Total	C	N	O	S	0	4	0
			3596	2276	621	673	26			
1	B	455	Total	C	N	O	S	0	3	0
			3627	2292	632	677	26			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	205	ALA	GLU	conflict	UNP P31013
A	448	ALA	PHE	engineered mutation	UNP P31013
B	205	ALA	GLU	conflict	UNP P31013
B	448	ALA	PHE	engineered mutation	UNP P31013

- Molecule 2 is POTASSIUM ION (three-letter code: K) (formula: K).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	K	0	0
			2	2		

- Molecule 3 is (2E)-2-{[(Z)-{3-hydroxy-2-methyl-5-[(phosphonoxy)methyl]pyridin-4(1H)-ylidene}methyl]imino}-3-phenylpropanoic acid (three-letter code: P71) (formula: C<sub>17</sub>H<sub>19</sub>N<sub>2</sub>O<sub>7</sub>P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			27	17	2	7	1		

- Molecule 4 is 3,6,9,12,15,18-HEXAOXAICOSANE-1,20-DIOL (three-letter code: P33) (formula:  $C_{14}H_{30}O_8$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			22	14	8		

- Molecule 5 is (E)-N-({3-hydroxy-2-methyl-5-[(phosphonoxy)methyl]pyridin-4-yl}methylidene)-L-phenylalanine (three-letter code: P70) (formula:  $C_{17}H_{19}N_2O_7P$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	P	0	0
			27	17	2	7	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	231	Total	O	0	0
			231	231		
6	B	262	Total	O	0	0
			262	262		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine phenol-lyase



- Molecule 1: Tyrosine phenol-lyase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.78Å 132.67Å 144.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.75 – 2.27 37.75 – 2.27	Depositor EDS
% Data completeness (in resolution range)	99.8 (37.75-2.27) 99.8 (37.75-2.27)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.24 (at 2.27Å)	Xtriage
Refinement program	PHENIX 1.13 _2998	Depositor
R, $R_{free}$	0.186 , 0.219 0.186 , 0.219	Depositor DCC
$R_{free}$ test set	1997 reflections (3.70%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	54.6	Xtriage
Anisotropy	0.457	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 71.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	7794	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	79.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.80% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: P33, K, P70, P71

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.24	0/3667	0.41	0/4937
1	B	0.24	0/3699	0.41	0/4981
All	All	0.24	0/7366	0.41	0/9918

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3596	0	3538	37	0
1	B	3627	0	3571	21	0
2	A	2	0	0	0	0
3	A	27	0	0	2	0
4	A	22	0	30	0	0
5	B	27	0	0	1	0
6	A	231	0	0	1	0
6	B	262	0	0	1	0
All	All	7794	0	7139	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.



All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:361:PHE:HD1	1:A:383:ILE:HD11	1.63	0.64
1:A:43:ILE:HD13	1:A:377:ARG:HD2	1.82	0.62
1:A:347:LEU:HD11	1:A:424:ILE:HD13	1.84	0.58
1:A:379:MET:HE3	1:A:381:ARG:HG2	1.85	0.58
1:A:112:LYS:O	1:A:114:GLY:N	2.37	0.58
1:A:392:THR:OG1	1:A:393:GLY:N	2.36	0.57
1:B:392:THR:HB	1:B:394:GLU:OE1	2.08	0.54
1:A:148:ALA:HB1	1:A:337:VAL:HG22	1.90	0.54
1:A:374:THR:HG21	1:A:423:GLY:HA3	1.90	0.53
1:A:394:GLU:OE2	1:A:394:GLU:N	2.38	0.52
1:B:347:LEU:HD11	1:B:424:ILE:HD13	1.90	0.52
1:B:45:ILE:HB	1:B:376:VAL:HG22	1.91	0.51
1:B:142:ARG:NH2	1:B:154:PHE:O	2.32	0.51
1:A:361:PHE:CD1	1:A:383:ILE:HD11	2.45	0.51
1:A:97:HIS:CD2	1:A:98:GLN:HG2	2.47	0.50
1:B:35:THR:HB	1:B:377:ARG:HG3	1.93	0.50
1:B:46:ASP:OD2	1:B:377:ARG:NH2	2.44	0.49
1:A:355:HIS:CD2	1:A:432:GLU:HA	2.47	0.49
1:B:394:GLU:N	1:B:394:GLU:OE1	2.45	0.48
1:B:97:HIS:CD2	1:B:98:GLN:HG2	2.47	0.48
1:A:188:GLY:HA2	1:A:346:PHE:CE1	2.49	0.48
1:B:210:LYS:NZ	6:B:611:HOH:O	2.46	0.47
1:A:79[B]:HIS:HD2	1:A:82:ARG:HH22	1.61	0.47
1:A:2:ASN:HB3	1:A:3:TYR:H	1.59	0.47
1:A:359:ASP:OD1	1:A:397:ARG:NH2	2.47	0.47
1:B:201:ARG:HB2	1:B:247:TYR:HB3	1.96	0.46
1:A:318:ARG:NH1	1:A:407:ILE:O	2.34	0.46
1:A:81:GLU:HA	1:A:93[B]:ILE:HG21	1.99	0.46
1:B:352:PHE:O	1:B:431:LYS:HD2	2.16	0.45
1:B:362:PRO:HG2	1:B:401:GLU:HG2	1.99	0.45
1:A:355:HIS:CE1	1:A:356:LEU:HG	2.52	0.45
3:A:502:P71:N11	3:A:502:P71:O15	2.47	0.45
1:A:364:GLN:HB2	1:A:384:ILE:HD12	1.99	0.45
1:A:48:LEU:HB2	1:A:377:ARG:HD3	1.99	0.44
1:A:122:TYR:HB3	1:A:137:PHE:CZ	2.52	0.44
1:A:404:ARG:NE	1:A:406:THR:OG1	2.50	0.44
1:A:257:LYS:HZ1	3:A:502:P71:C12	2.31	0.44
1:A:444:LYS:O	1:A:446:LEU:N	2.47	0.44
1:A:17:SER:OG	1:A:41:LYS:O	2.22	0.43
1:B:92:HIS:HB3	1:B:275:PHE:CD1	2.53	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:336:ILE:HG22	1:A:347:LEU:HD23	2.01	0.43
1:A:362:PRO:HG2	1:A:401:GLU:HG2	2.01	0.42
1:A:361:PHE:HD2	1:A:437:LEU:HB2	1.85	0.42
1:B:201:ARG:HD2	1:B:247:TYR:O	2.19	0.42
1:A:92:HIS:HB3	1:A:275:PHE:CD1	2.55	0.41
1:B:122:TYR:HB3	1:B:137:PHE:CZ	2.55	0.41
1:A:447:ARG:H	1:A:447:ARG:HD2	1.84	0.41
1:A:52:GLY:N	1:A:256:LYS:HB3	2.35	0.41
1:B:120:ASN:OD1	1:B:120:ASN:N	2.53	0.41
1:B:351:ARG:HE	1:B:351:ARG:HB2	1.75	0.41
1:B:344:ALA:HB2	1:B:406:THR:HG23	2.01	0.41
1:A:374:THR:CG2	1:A:423:GLY:HA3	2.50	0.41
1:B:257:LYS:HZ1	5:B:501:P70:C05	2.33	0.41
1:B:52:GLY:N	1:B:256:LYS:HB3	2.36	0.41
1:B:216:THR:HG22	1:B:254:SER:H	1.85	0.41
1:A:256:LYS:NZ	6:A:616:HOH:O	2.51	0.41
1:A:58:ASP:N	1:A:58:ASP:OD1	2.54	0.41
1:A:216:THR:HG22	1:A:254:SER:H	1.87	0.40
1:A:381:ARG:HA	1:A:381:ARG:HD3	1.87	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	451/456 (99%)	434 (96%)	14 (3%)	3 (1%)	22	25
1	B	456/456 (100%)	441 (97%)	14 (3%)	1 (0%)	47	57
All	All	907/912 (100%)	875 (96%)	28 (3%)	4 (0%)	34	40

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	446	LEU
1	A	441	TYR
1	A	113	PRO
1	A	114	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	376/377 (100%)	370 (98%)	6 (2%)	62	76
1	B	379/377 (100%)	378 (100%)	1 (0%)	92	96
All	All	755/754 (100%)	748 (99%)	7 (1%)	78	88

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	355	HIS
1	A	392	THR
1	A	395	HIS
1	A	406	THR
1	A	447	ARG
1	A	450	THR
1	B	406	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	P33	A	503	-	21,21,21	0.54	0	20,20,20	0.20	0
5	P70	B	501	-	25,28,28	3.06	7 (28%)	31,39,39	2.65	7 (22%)
3	P71	A	502	-	24,28,28	1.93	5 (20%)	29,39,39	1.92	8 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	P33	A	503	-	-	8/19/19/19	-
5	P70	B	501	-	-	1/15/19/19	0/2/2/2
3	P71	A	502	-	-	1/11/19/19	0/2/2/2

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	501	P70	C07-C08	9.81	1.50	1.40
3	A	502	P71	P23-O22	6.63	1.81	1.60
5	B	501	P70	C03-N04	6.43	1.53	1.46
5	B	501	P70	O19-C07	-5.50	1.24	1.37
5	B	501	P70	C06-C07	4.71	1.47	1.40
3	A	502	P71	C16-N18	3.69	1.40	1.33
5	B	501	P70	C06-C05	2.97	1.52	1.46
3	A	502	P71	C19-N18	2.92	1.40	1.34
3	A	502	P71	O15-C14	-2.58	1.27	1.35

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	501	P70	C05-N04	2.23	1.31	1.27
5	B	501	P70	C11-N10	2.16	1.39	1.34
3	A	502	P71	O22-C21	-2.03	1.37	1.45

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	501	P70	C03-N04-C05	12.41	134.23	117.40
3	A	502	P71	C13-C14-C16	-5.98	116.48	120.19
3	A	502	P71	C14-C13-C20	4.14	121.44	118.26
5	B	501	P70	C06-C07-C08	-3.38	118.09	120.19
5	B	501	P70	C12-C06-C05	3.34	127.06	121.56
3	A	502	P71	O25-P23-O22	-3.00	98.75	106.73
5	B	501	P70	C07-C06-C05	-2.85	115.10	120.41
3	A	502	P71	O22-C21-C20	2.76	114.61	109.35
3	A	502	P71	O26-P23-O25	2.68	117.89	107.64
3	A	502	P71	O26-P23-O22	-2.41	100.33	106.73
3	A	502	P71	O15-C14-C16	2.37	122.66	117.49
5	B	501	P70	C21-C20-C03	2.35	117.84	113.23
5	B	501	P70	C06-C05-N04	2.21	127.91	123.01
3	A	502	P71	C21-C20-C19	2.05	122.74	119.37
5	B	501	P70	O17-P15-O16	2.02	115.38	107.64

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	B	501	P70	C20-C03-N04-C05
4	A	503	P33	O4-C5-C6-O7
4	A	503	P33	O7-C8-C9-O10
4	A	503	P33	O1-C2-C3-O4
4	A	503	P33	O13-C14-C15-O16
4	A	503	P33	C2-C3-O4-C5
3	A	502	P71	C13-C20-C21-O22
4	A	503	P33	C12-C11-O10-C9
4	A	503	P33	C9-C8-O7-C6
4	A	503	P33	O10-C11-C12-O13

There are no ring outliers.


2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	501	P70	1	0
3	A	502	P71	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## Ligand P70 B 501





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	451/456 (98%)	0.49	58 (12%) <b>3</b> <b>4</b>	48, 74, 134, 188	0
1	B	455/456 (99%)	0.19	23 (5%) 28 33	48, 71, 108, 179	0
All	All	906/912 (99%)	0.34	81 (8%) <b>9</b> <b>12</b>	48, 73, 128, 188	0

All (81) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	391	VAL	11.8
1	B	390	ASN	6.8
1	B	392	THR	4.9
1	A	441	TYR	4.8
1	A	455	TYR	4.7
1	A	439	PHE	4.6
1	A	453	PHE	4.4
1	A	168	ASP	4.3
1	A	33	TYR	4.3
1	A	456	ILE	4.1
1	A	22	ASP	4.1
1	A	394	GLU	3.9
1	A	392	THR	3.7
1	A	51	SER	3.7
1	A	167	ILE	3.6
1	B	116	TYR	3.6
1	B	134	GLY	3.4
1	B	394	GLU	3.3
1	A	393	GLY	3.3
1	A	408	PRO	3.2
1	A	445	GLN	3.2
1	B	173	GLU	3.2
1	A	52	GLY	3.1
1	A	53	THR	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	166	LEU	3.1
1	A	447	ARG	3.1
1	A	254	SER	3.0
1	A	23	GLU	3.0
1	A	48	LEU	3.0
1	A	216	THR	3.0
1	B	171	GLY	2.9
1	A	256	LYS	2.9
1	A	411	VAL	2.9
1	B	112	LYS	2.9
1	A	163	LEU	2.8
1	A	49	THR	2.8
1	A	395	HIS	2.8
1	A	173	GLU	2.8
1	A	397	ARG	2.7
1	B	389	ASN	2.7
1	A	409	ARG	2.7
1	A	361	PHE	2.7
1	A	387	GLY	2.7
1	B	49	THR	2.6
1	B	433	ASP	2.6
1	A	261	VAL	2.5
1	B	456	ILE	2.5
1	B	71	TYR	2.5
1	A	113	PRO	2.5
1	B	445	GLN	2.4
1	A	175	ILE	2.4
1	A	172	ALA	2.4
1	B	113	PRO	2.4
1	B	97	HIS	2.4
1	B	294	LEU	2.3
1	A	96	THR	2.3
1	A	116	TYR	2.3
1	B	174	ASN	2.3
1	A	263	ILE	2.3
1	A	452	ARG	2.2
1	A	253	MET	2.2
1	A	20	PRO	2.2
1	A	99	GLY	2.2
1	B	234	ASN	2.2
1	A	207	HIS	2.2
1	A	50	ASP	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	25	LEU	2.1
1	B	20	PRO	2.1
1	A	258	ASP	2.1
1	A	359	ASP	2.1
1	B	393	GLY	2.1
1	A	54	ASN	2.1
1	A	443	PRO	2.1
1	A	383	ILE	2.1
1	A	203	LEU	2.1
1	A	97	HIS	2.1
1	A	183	THR	2.1
1	A	442	GLU	2.0
1	A	19	ILE	2.0
1	A	438	LYS	2.0
1	A	134	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	P33	A	503	22/22	0.82	0.17	89,103,124,132	0
2	K	A	504	1/1	0.96	0.08	63,63,63,63	0
5	P70	B	501	27/27	0.96	0.22	51,85,94,105	0
2	K	A	501	1/1	0.97	0.14	60,60,60,60	0
3	P71	A	502	27/27	0.97	0.27	56,69,94,97	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around P70 B 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around P71 A 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers ⓘ

There are no such residues in this entry.