



Full wwPDB X-ray Structure Validation Report ⓘ

May 24, 2020 – 10:44 pm BST

PDB ID : 4DWH
Title : Structure of the major type 1 pilus subunit FIMA bound to the FIMC (2.5 Å resolution)
Authors : Scharer, M.A.; Puorger, C.; Crespo, M.; Glockshuber, R.; Capitani, G.
Deposited on : 2012-02-24
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	205	<div> <div>2%</div> <div> <div></div> <div>78%</div> <div>17%</div> <div>• •</div> </div> </div>
1	C	205	<div> <div>%</div> <div> <div></div> <div>83%</div> <div>14%</div> <div>• •</div> </div> </div>
2	B	143	<div> <div>3%</div> <div> <div></div> <div>81%</div> <div>12%</div> <div>7%</div> </div> </div>
2	D	143	<div> <div>6%</div> <div> <div></div> <div>82%</div> <div>10%</div> <div>• 6%</div> </div> </div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 5253 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chaperone protein fimC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	197	Total	C	N	O	S	17	0	0
			1536	973	266	291	6			
1	C	201	Total	C	N	O	S	12	0	0
			1567	991	271	299	6			

- Molecule 2 is a protein called Type-1 fimbrial protein, A chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	133	Total	C	N	O	S	1	0	0
			936	573	164	196	3			
2	D	134	Total	C	N	O	S	4	1	0
			953	583	166	200	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	17	MET	-	EXPRESSION TAG	UNP P04128
D	17	MET	-	EXPRESSION TAG	UNP P04128

- Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	3	0
			7	4	3		
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			7	4	3		
3	B	1	Total	C	O	0	0
			7	4	3		
3	C	1	Total	C	O	0	0
			7	4	3		
3	C	1	Total	C	O	3	0
			7	4	3		
3	D	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: C₈H₁₈O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			13	8	5		
4	D	1	Total	C	O	0	0
			13	8	5		

- Molecule 5 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	P	0	0
			5	4	1		
5	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 6 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	C	1	Total	C	O	0	0
			10	6	4		
6	C	1	Total	C	O	0	0
			10	6	4		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	54	Total	O	0	0
			54	54		
7	B	22	Total	O	0	0
			22	22		
7	C	33	Total	O	0	0
			33	33		
7	D	26	Total	O	0	0
			26	26		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Chaperone protein fimC




- Molecule 1: Chaperone protein fimC



- Molecule 2: Type-1 fimbrial protein, A chain



- Molecule 2: Type-1 fimbrial protein, A chain



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	122.37Å 48.81Å 113.48Å 90.00° 98.90° 90.00°	Depositor
Resolution (Å)	45.26 – 2.50 45.26 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.9 (45.26-2.50) 100.0 (45.26-2.50)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.89 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1.7.3 _928	Depositor
R, R_{free}	0.233 , 0.294 0.227 , 0.290	Depositor DCC
R_{free} test set	792 reflections (3.40%)	wwPDB-VP
Wilson B-factor (Å ²)	34.7	Xtriage
Anisotropy	0.399	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 63.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5253	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 18.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PG4, PO4, PEG, PGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.28	0/1563	0.48	0/2124
1	C	0.28	0/1595	0.50	0/2168
2	B	0.28	0/944	0.55	0/1284
2	D	0.27	0/961	0.48	0/1306
All	All	0.28	0/5063	0.50	0/6882

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1536	0	1570	25	0
1	C	1567	0	1599	17	0
2	B	936	0	901	8	0
2	D	953	0	914	8	0
3	A	42	0	60	6	0
3	B	7	0	10	0	0
3	C	14	0	20	0	0
3	D	7	0	10	0	0
4	A	13	0	18	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	13	0	18	0	0
5	A	5	0	0	0	0
5	B	5	0	0	0	0
6	C	20	0	28	2	0
7	A	54	0	0	2	0
7	B	22	0	0	0	0
7	C	33	0	0	0	0
7	D	26	0	0	0	0
All	All	5253	0	5148	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:118:PHE:HB3	2:B:119:SER:HA	1.52	0.89
1:A:61:LYS:HG2	1:C:61:LYS:HG2	1.71	0.72
2:D:72:ALA:HB2	2:D:117:THR:HA	1.75	0.69
1:A:200:MET:HG2	3:A:305:PEG:H42	1.77	0.67
1:C:132:ARG:HB2	1:C:143:ILE:HB	1.76	0.67
1:A:38:GLU:OE1	1:A:110:ARG:NH1	2.28	0.65
2:D:97:THR:HG1	2:D:144:THR:HG1	1.45	0.64
1:C:117:PRO:HG2	1:C:120:LEU:HD21	1.82	0.61
1:C:47:ARG:HB3	1:C:71:THR:HG22	1.83	0.61
1:A:191:ASN:HA	3:A:301:PEG:H22	1.86	0.57
1:A:125:ASP:OD2	1:A:126:GLN:NE2	2.35	0.57
2:B:104:LEU:HD23	2:B:108:GLY:HA2	1.87	0.56
1:C:178:PRO:HG2	1:C:181:ALA:HB2	1.87	0.55
1:A:112:LYS:NZ	1:A:151:THR:OG1	2.36	0.55
2:B:85:VAL:HG22	2:B:102:GLN:HE21	1.73	0.53
1:A:201:THR:H	3:A:305:PEG:H21	1.75	0.51
1:A:66:ARG:NH2	7:A:434:HOH:O	2.42	0.51
2:B:79:ASP:OD1	2:B:82:HIS:ND1	2.33	0.51
1:C:133:PHE:HB3	1:C:140:LEU:HD11	1.93	0.50
1:A:57:MET:HA	1:A:61:LYS:HD3	1.94	0.50
1:A:73:ASN:ND2	3:A:306:PEG:O2	2.44	0.49
2:B:42:LEU:HD12	2:B:139:ALA:HB2	1.95	0.48
1:A:110:ARG:NH2	7:A:418:HOH:O	2.46	0.48
1:A:183:SER:HA	1:A:204:MET:SD	2.54	0.47
1:A:61:LYS:HE2	1:A:61:LYS:HB3	1.70	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:25:ASN:ND2	2:B:60:ASP:OD2	2.42	0.47
2:D:106:ARG:HB3	2:D:132:PRO:HG2	1.96	0.46
1:A:32:LEU:HB2	3:A:302:PEG:H12	1.97	0.46
2:D:97:THR:OG1	2:D:144:THR:OG1	2.25	0.46
2:B:118:PHE:CB	2:B:119:SER:HA	2.32	0.45
6:C:303:PGE:H1	6:C:303:PGE:H3	1.67	0.45
1:C:45:ASP:OD2	1:C:47:ARG:NE	2.49	0.45
1:C:136:SER:OG	1:C:137:ALA:N	2.50	0.44
1:C:39:ASN:HD21	1:C:43:VAL:HB	1.82	0.44
1:A:49:ILE:HB	1:A:68:LEU:HB2	1.98	0.44
1:C:38:GLU:OE1	1:C:110:ARG:NH1	2.48	0.44
1:A:191:ASN:HB2	3:A:301:PEG:H41	2.00	0.44
1:C:170:MET:HE3	1:C:170:MET:HB2	1.93	0.44
2:D:62:ASP:HB3	2:D:65:VAL:HG23	2.00	0.44
4:A:307:PG4:H11	4:A:307:PG4:H32	1.79	0.43
1:A:2:VAL:HA	1:A:23:THR:O	2.18	0.43
1:C:11:TYR:HA	1:C:12:PRO:HD3	1.87	0.43
1:C:60:LYS:HE3	6:C:302:PGE:H12	2.00	0.43
1:A:167:VAL:HA	1:A:168:PRO:HD2	1.94	0.42
1:C:45:ASP:HB3	1:C:46:GLY:H	1.59	0.42
1:C:55:PHE:HE1	1:C:65:LEU:HD21	1.85	0.42
2:D:104:LEU:HD23	2:D:108:GLY:HA2	2.02	0.42
1:A:126:GLN:O	1:A:129:GLU:HG2	2.20	0.41
1:A:177:LEU:HA	1:A:178:PRO:HD3	1.78	0.41
1:C:79:ARG:HB3	1:C:170:MET:HE1	2.02	0.41
1:C:116:ARG:HA	1:C:117:PRO:HD2	1.90	0.41
2:D:71:VAL:HG13	2:D:154:PHE:CE1	2.56	0.41
2:D:62:ASP:OD2	2:D:64:ASN:HB2	2.21	0.41
1:A:160:ARG:HG2	1:A:178:PRO:HG2	2.03	0.41
2:B:106:ARG:HE	2:B:106:ARG:HB3	1.71	0.41
1:A:32:LEU:HG	1:A:54:LEU:HD11	2.04	0.40
1:A:78:ASP:N	1:A:78:ASP:OD1	2.54	0.40
1:A:4:LEU:HD23	1:A:111:ILE:HD13	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	191/205 (93%)	181 (95%)	9 (5%)	1 (0%)	29	48
1	C	197/205 (96%)	184 (93%)	9 (5%)	4 (2%)	7	12
2	B	127/143 (89%)	118 (93%)	9 (7%)	0	100	100
2	D	128/143 (90%)	121 (94%)	6 (5%)	1 (1%)	19	35
All	All	643/696 (92%)	604 (94%)	33 (5%)	6 (1%)	17	31

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	45	ASP
1	C	44	LYS
1	C	137	ALA
1	A	101	ASN
1	C	101	ASN
2	D	105	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	169/176 (96%)	160 (95%)	9 (5%)	22	43
1	C	172/176 (98%)	171 (99%)	1 (1%)	86	95
2	B	94/101 (93%)	89 (95%)	5 (5%)	22	43
2	D	96/101 (95%)	91 (95%)	5 (5%)	23	44

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	531/554 (96%)	511 (96%)	20 (4%)	33 58

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	ARG
1	A	16	LYS
1	A	45	ASP
1	A	110	ARG
1	A	125	ASP
1	A	153	THR
1	A	160	ARG
1	A	172	GLU
1	A	184	ASN
2	B	17	MET
2	B	28	VAL
2	B	57	GLN
2	B	88	LEU
2	B	112	THR
1	C	125	ASP
2	D	24	ASP
2	D	71	VAL
2	D	114	ASP
2	D	117	THR
2	D	159	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PEG	A	304	-	6,6,6	0.63	0	5,5,5	1.43	0
3	PEG	C	301	-	6,6,6	0.61	0	5,5,5	1.55	0
3	PEG	D	202	-	6,6,6	0.64	0	5,5,5	1.47	0
5	PO4	A	308	-	4,4,4	0.86	0	6,6,6	0.50	0
3	PEG	A	305	-	6,6,6	0.65	0	5,5,5	1.40	0
3	PEG	A	301	-	6,6,6	0.61	0	5,5,5	1.53	1 (20%)
6	PGE	C	302	-	9,9,9	0.48	0	8,8,8	0.47	0
3	PEG	A	306	-	6,6,6	0.58	0	5,5,5	1.58	0
3	PEG	B	201	-	6,6,6	0.60	0	5,5,5	1.55	0
3	PEG	A	302	-	6,6,6	0.62	0	5,5,5	1.48	0
6	PGE	C	303	-	9,9,9	0.49	0	8,8,8	0.70	0
3	PEG	A	303	-	6,6,6	0.62	0	5,5,5	1.48	0
4	PG4	D	201	-	12,12,12	0.63	0	11,11,11	1.65	1 (9%)
4	PG4	A	307	-	12,12,12	0.68	0	11,11,11	1.45	0
5	PO4	B	202	-	4,4,4	0.90	0	6,6,6	0.46	0
3	PEG	C	304	-	6,6,6	0.66	0	5,5,5	1.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEG	A	304	-	-	2/4/4/4	-
3	PEG	C	301	-	-	0/4/4/4	-
3	PEG	D	202	-	-	2/4/4/4	-
3	PEG	A	305	-	-	2/4/4/4	-
3	PEG	A	301	-	-	2/4/4/4	-
6	PGE	C	302	-	-	2/7/7/7	-
3	PEG	A	306	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEG	B	201	-	-	0/4/4/4	-
3	PEG	A	302	-	-	0/4/4/4	-
6	PGE	C	303	-	-	1/7/7/7	-
3	PEG	A	303	-	-	1/4/4/4	-
4	PG4	D	201	-	-	4/10/10/10	-
4	PG4	A	307	-	-	7/10/10/10	-
3	PEG	C	304	-	-	1/4/4/4	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	301	PEG	O2-C3-C4	2.04	119.02	110.07
4	D	201	PG4	O3-C5-C6	2.00	119.42	110.39

There are no chirality outliers.

All (26) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	307	PG4	C1-C2-O2-C3
3	A	305	PEG	C4-C3-O2-C2
4	D	201	PG4	O3-C5-C6-O4
4	A	307	PG4	O3-C5-C6-O4
3	A	304	PEG	O2-C3-C4-O4
4	D	201	PG4	O2-C3-C4-O3
3	A	306	PEG	O1-C1-C2-O2
4	D	201	PG4	O4-C7-C8-O5
4	A	307	PG4	O4-C7-C8-O5
3	D	202	PEG	C4-C3-O2-C2
3	A	301	PEG	O2-C3-C4-O4
3	A	303	PEG	O1-C1-C2-O2
3	C	304	PEG	O1-C1-C2-O2
4	A	307	PG4	C6-C5-O3-C4
3	A	301	PEG	C4-C3-O2-C2
4	D	201	PG4	C3-C4-O3-C5
6	C	303	PGE	C1-C2-O2-C3
3	A	305	PEG	O1-C1-C2-O2
4	A	307	PG4	C3-C4-O3-C5
6	C	302	PGE	C4-C3-O2-C2
3	D	202	PEG	C1-C2-O2-C3
4	A	307	PG4	O2-C3-C4-O3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	304	PEG	C1-C2-O2-C3
6	C	302	PGE	O2-C3-C4-O3
3	A	306	PEG	O2-C3-C4-O4
4	A	307	PG4	C4-C3-O2-C2

There are no ring outliers.

7 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	305	PEG	2	0
3	A	301	PEG	2	0
6	C	302	PGE	1	0
3	A	306	PEG	1	0
3	A	302	PEG	1	0
6	C	303	PGE	1	0
4	A	307	PG4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	197/205 (96%)	0.12	4 (2%) 65 68	17, 32, 66, 90	4 (2%)
1	C	201/205 (98%)	0.14	3 (1%) 73 75	18, 35, 64, 97	4 (1%)
2	B	133/143 (93%)	0.37	5 (3%) 40 43	17, 41, 68, 73	1 (0%)
2	D	134/143 (93%)	0.41	8 (5%) 21 22	20, 43, 69, 88	1 (0%)
All	All	665/696 (95%)	0.23	20 (3%) 50 53	17, 36, 67, 97	10 (1%)

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	94	GLY	4.6
2	D	128	THR	3.9
2	B	120	SER	3.6
1	C	138	ASN	3.3
1	A	136	SER	3.2
2	D	109	ALA	2.9
2	D	147	ALA	2.5
1	C	118	ALA	2.4
2	D	81	GLY	2.3
2	B	17	MET	2.3
1	A	203	VAL	2.3
1	C	136	SER	2.3
2	B	81	GLY	2.3
2	B	142	ALA	2.1
2	B	118	PHE	2.1
2	D	144	THR	2.1
2	D	110	ALA	2.1
1	A	94	ASP	2.1
1	A	141	THR	2.0
2	D	95	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	PEG	A	304	7/7	0.69	0.24	42,47,51,51	0
5	PO4	B	202	5/5	0.76	0.40	104,105,109,110	0
3	PEG	A	306	7/7	0.79	0.23	41,48,53,55	0
3	PEG	C	301	7/7	0.79	0.25	42,48,50,51	0
3	PEG	A	303	7/7	0.81	0.23	44,44,47,48	3
3	PEG	B	201	7/7	0.86	0.16	33,34,39,40	0
5	PO4	A	308	5/5	0.88	0.21	57,61,67,68	0
4	PG4	D	201	13/13	0.89	0.15	29,36,48,53	0
3	PEG	D	202	7/7	0.89	0.19	50,51,59,61	0
3	PEG	A	302	7/7	0.90	0.16	36,37,47,51	0
3	PEG	C	304	7/7	0.90	0.16	56,59,59,59	3
6	PGE	C	303	10/10	0.92	0.20	43,47,49,51	0
3	PEG	A	301	7/7	0.92	0.17	20,28,33,36	0
3	PEG	A	305	7/7	0.93	0.14	46,47,51,53	0
4	PG4	A	307	13/13	0.93	0.17	29,32,54,55	0
6	PGE	C	302	10/10	0.96	0.13	27,35,36,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.