



Full wwPDB NMR Structure Validation Report ⓘ

Jun 15, 2020 – 11:50 pm BST

PDB ID : 6DZB
Title : Solution structure of Rbfox2 RRM mimetic peptide CPfox5
Authors : Sun, Y.-T.; Varani, G.
Deposited on : 2018-07-03

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 50%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	16	

2 Ensemble composition and analysis ⓘ

This entry contains 10 models.

Cyrange was unable to find well-defined residues.

Error message: Only domains with < 8 residues could be identified.

NmrClust was unable to cluster the ensemble.

Error message: Wrapper check: not enough residues in core to run NmrClust

3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 274 atoms, of which 138 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called CPfox5.


Mol	Chain	Residues	Atoms					Trace
1	A	16	Total	C	H	N	O	0
			274	89	138	27	20	

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: CPfox5




4.2 Scores per residue for each member of the ensemble

Colouring as in section [4.1](#) above.

4.2.1 Score per residue for model 1

- Molecule 1: CPfox5



4.2.2 Score per residue for model 2

- Molecule 1: CPfox5



4.2.3 Score per residue for model 3

- Molecule 1: CPfox5

Chain A:  50% 44% 6%



4.2.4 Score per residue for model 4

- Molecule 1: CPfox5

Chain A:  50% 38% 13%



4.2.5 Score per residue for model 5

- Molecule 1: CPfox5


Chain A:  63% 38%



4.2.6 Score per residue for model 6


- Molecule 1: CPfox5


Chain A:  56% 25% 13% 6%



4.2.7 Score per residue for model 7

- Molecule 1: CPfox5

Chain A:  38% 50% 13%



4.2.8 Score per residue for model 8

- Molecule 1: CPfox5


Chain A: 56% 31% 13%



4.2.9 Score per residue for model 9

- Molecule 1: CPfox5

Chain A: 50% 38% 13%



4.2.10 Score per residue for model 10

- Molecule 1: CPfox5

Chain A: 38% 50% 6% 6%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 200 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	refinement	
X-PLOR NIH	structure calculation	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	input_cs.cif
Number of chemical shift lists	1
Total number of shifts	127
Number of shifts mapped to atoms	127
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	50%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality ⓘ

6.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: DPR

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	136	138	138	4±1
All	All	1360	1380	1379	36

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:3:ARG:CG	1:A:4:GLY:H	0.54	2.15	7	1
1:A:2:SER:OG	1:A:3:ARG:N	0.53	2.42	10	8
1:A:14:ASN:OD1	1:A:15:GLU:N	0.53	2.42	1	2
1:A:5:PHE:N	1:A:12:ILE:O	0.52	2.42	1	3
1:A:3:ARG:O	1:A:5:PHE:CZ	0.51	2.63	5	1
1:A:2:SER:CB	1:A:14:ASN:H	0.51	2.19	6	1
1:A:3:ARG:NH1	1:A:14:ASN:OD1	0.49	2.45	9	1
1:A:3:ARG:NH2	1:A:14:ASN:HD21	0.49	2.06	9	1
1:A:3:ARG:O	1:A:5:PHE:CD1	0.49	2.66	7	1
1:A:14:ASN:CG	1:A:15:GLU:N	0.47	2.67	3	2
1:A:13:PHE:CD1	1:A:13:PHE:N	0.45	2.83	1	1
1:A:14:ASN:O	1:A:16:ARG:N	0.45	2.48	3	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:14:ASN:ND2	1:A:14:ASN:C	0.45	2.68	10	1
1:A:3:ARG:CG	1:A:4:GLY:N	0.44	2.80	7	3
1:A:13:PHE:C	1:A:13:PHE:CD2	0.43	2.89	4	2
1:A:3:ARG:O	1:A:5:PHE:CE1	0.43	2.71	7	1
1:A:3:ARG:O	1:A:4:GLY:O	0.42	2.37	4	1
1:A:2:SER:N	1:A:16:ARG:C	0.42	2.72	9	1
1:A:15:GLU:O	1:A:16:ARG:O	0.42	2.37	1	1
1:A:8:DPR:C	1:A:10:LYS:H	0.41	2.29	10	1
1:A:2:SER:H	1:A:16:ARG:C	0.41	2.18	9	1
1:A:1:GLY:O	1:A:2:SER:O	0.41	2.37	10	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	13/16 (81%)	9±1 (66±9%)	3±1 (25±7%)	1±1 (8±5%)	2	13
All	All	130/160 (81%)	86 (66%)	33 (25%)	11 (8%)	2	13

All 4 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	14	ASN	4
1	A	4	GLY	3
1	A	2	SER	3
1	A	15	GLU	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	13/13 (100%)	9±1 (65±8%)	5±1 (35±8%)	1	10
All	All	130/130 (100%)	85 (65%)	45 (35%)	1	10

All 11 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	7	PHE	10
1	A	12	ILE	9
1	A	13	PHE	9
1	A	14	ASN	6
1	A	16	ARG	3
1	A	2	SER	2
1	A	3	ARG	2
1	A	6	ARG	1
1	A	15	GLU	1
1	A	11	ILE	1
1	A	10	LYS	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.6 Ligand geometry

There are no ligands in this entry.

6.7 Other polymers

There are no such molecules in this entry.

6.8 Polymer linkage issues

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 50% for the well-defined parts and 50% for the entire structure.

7.1 Chemical shift list 1

File name: input_cs.cif

Chemical shift list name: *cpfox5_assign*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	127
Number of shifts mapped to atoms	127
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 50%, i.e. 111 atoms were assigned a chemical shift out of a possible 220. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	42/73 (58%)	28/29 (97%)	0/30 (0%)	14/14 (100%)
Sidechain	61/120 (51%)	61/73 (84%)	0/36 (0%)	0/11 (0%)
Aromatic	8/27 (30%)	8/15 (53%)	0/12 (0%)	0/0 (—%)
Overall	111/220 (50%)	97/117 (83%)	0/78 (0%)	14/25 (56%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 50%, i.e. 111 atoms were assigned a chemical shift out of a possible 220. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	42/73 (58%)	28/29 (97%)	0/30 (0%)	14/14 (100%)
Sidechain	61/120 (51%)	61/73 (84%)	0/36 (0%)	0/11 (0%)
Aromatic	8/27 (30%)	8/15 (53%)	0/12 (0%)	0/0 (—%)
Overall	111/220 (50%)	97/117 (83%)	0/78 (0%)	14/25 (56%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

