



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 17, 2022 – 10:08 AM JST

PDB ID : 7DZN
Title : Crystal Structure of the cross-restricted T18A TCR and HLAB4201 bound to HIV-1 Gag TL9 peptide
Authors : Liu, Y.; Yin, L.
Deposited on : 2021-01-25
Resolution : 2.63 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.25
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.25


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.63 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1426 (2.66-2.62)
Clashscore	141614	1472 (2.66-2.62)
Ramachandran outliers	138981	1446 (2.66-2.62)
Sidechain outliers	138945	1446 (2.66-2.62)
RSRZ outliers	127900	1408 (2.66-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	279	<div> <div>9%</div> <div>80%</div> <div>17%</div> <div>..</div> </div>
2	B	100	<div> <div>%</div> <div>80%</div> <div>19%</div> <div>.</div> </div>
3	C	9	<div> <div>78%</div> <div>22%</div> </div>
4	D	244	<div> <div>86%</div> <div>13%</div> <div>.</div> </div>
5	E	204	<div> <div>2%</div> <div>76%</div> <div>21%</div> <div>..</div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6806 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MHC class I antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	277	Total	C	N	O	S	0	6	0
			2292	1423	414	449	6			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP V6E0U6
A	2	GLY	-	expression tag	UNP V6E0U6

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	expression tag	UNP P61769

- Molecule 3 is a protein called Gag-Pol polyprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	9	Total	C	N	O	S	0	0	0
			70	43	11	15	1			

- Molecule 4 is a protein called beta chain T18A TCR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	241	Total	C	N	O	S	0	0	0
			1919	1214	330	365	10			

- Molecule 5 is a protein called alpha chain T18A TCR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	202	Total	C	N	O	S	0	0	0
			1572	974	270	316	12			


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	39	Total	O	0	0
			39	39		
6	B	10	Total	O	0	0
			10	10		
6	C	1	Total	O	0	0
			1	1		
6	D	46	Total	O	0	0
			46	46		
6	E	20	Total	O	0	0
			20	20		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MHC class I antigen




- Molecule 2: Beta-2-microglobulin





- Molecule 3: Gag-Pol polyprotein



- Molecule 4: beta chain T18A TCR



- Molecule 5: alpha chain T18A TCR



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	96.99Å 96.99Å 263.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.70 – 2.63 47.70 – 2.63	Depositor EDS
% Data completeness (in resolution range)	99.8 (47.70-2.63) 99.8 (47.70-2.63)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.52 (at 2.61Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.207 , 0.261 0.204 , 0.257	Depositor DCC
R_{free} test set	1998 reflections (5.20%)	wwPDB-VP
Wilson B-factor (Å ²)	48.5	Xtriage
Anisotropy	0.695	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 39.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6806	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/2367	0.56	0/3218
2	B	0.26	0/860	0.52	0/1162
3	C	0.26	0/70	0.67	0/95
4	D	0.26	0/1973	0.52	0/2687
5	E	0.27	0/1605	0.53	1/2179 (0.0%)
All	All	0.27	0/6875	0.54	1/9341 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
5	E	164	LEU	CA-CB-CG	5.37	127.65	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2292	0	2134	40	0
2	B	837	0	803	12	1
3	C	70	0	72	2	0
4	D	1919	0	1825	16	0
5	E	1572	0	1497	28	1
6	A	39	0	0	2	0
6	B	10	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	C	1	0	0	0	0
6	D	46	0	0	0	0
6	E	20	0	0	3	0
All	All	6806	0	6331	87	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:191:VAL:HG11	1:A:276:TRP:HA	1.65	0.78
5:E:150:GLN:NE2	6:E:302:HOH:O	2.24	0.70
5:E:3:ASP:N	6:E:303:HOH:O	2.25	0.68
5:E:39:ARG:HB3	5:E:49:ILE:HD11	1.76	0.66
5:E:53:LEU:O	5:E:70:ARG:NH1	2.27	0.65
2:B:17:ASN:HD21	2:B:97:ARG:HH12	1.46	0.64
1:A:272:LEU:HD12	1:A:273:THR:H	1.62	0.64
4:D:139:LYS:HE3	5:E:169:MET:HE1	1.80	0.64
5:E:128:ASP:OD1	5:E:130:LYS:N	2.31	0.63
5:E:143:ASP:OD2	5:E:145:GLN:HB2	1.99	0.63
1:A:46:ARG:NH2	1:A:63:ASP:OD1	2.30	0.61
1:A:221:ARG:HD3	1:A:258:ARG:HH22	1.64	0.61
5:E:127:ARG:HH22	5:E:133:ASP:H	1.49	0.61
1:A:255:GLU:HB2	1:A:258:ARG:HD3	1.83	0.60
4:D:194:ARG:HB2	5:E:164:LEU:HD11	1.84	0.60
4:D:35:TRP:HB2	4:D:48:ILE:HG22	1.84	0.59
2:B:17:ASN:ND2	2:B:97:ARG:HH12	2.01	0.59
5:E:18:ARG:NH1	6:E:305:HOH:O	2.36	0.58
5:E:125:GLN:HG3	5:E:187:CYS:SG	2.43	0.58
1:A:221:ARG:O	1:A:223:GLY:N	2.34	0.57
1:A:199:HIS:HA	1:A:253:SER:HB2	1.86	0.57
1:A:221:ARG:HB2	1:A:226:GLN:NE2	2.20	0.55
1:A:171:ARG:NH2	6:A:305:HOH:O	2.40	0.54
1:A:100:MET:HG3	2:B:58:LYS:HZ1	1.73	0.54
1:A:119:ALA:HB2	2:B:60:TRP:CE2	2.42	0.54
1:A:100:MET:HG3	2:B:58:LYS:NZ	2.23	0.54
4:D:179:GLN:HE21	4:D:182:LEU:HD12	1.75	0.52
1:A:192:THR:HG23	1:A:194:HIS:HE1	1.75	0.52
1:A:221:ARG:HD3	1:A:258:ARG:NH2	2.24	0.52
1:A:192:THR:HG23	1:A:194:HIS:CE1	2.46	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:255:GLU:HB2	1:A:258:ARG:CD	2.40	0.51
5:E:111:MET:HE2	5:E:113:LYS:HD3	1.93	0.50
1:A:217:LEU:HD22	1:A:263:VAL:HG22	1.94	0.50
1:A:221:ARG:HG3	1:A:259:TYR:CZ	2.46	0.50
2:B:5:PRO:HB3	2:B:30:PHE:HB3	1.93	0.50
4:D:123:GLU:O	4:D:146:ALA:HA	2.11	0.50
1:A:276:TRP:O	1:A:278:PRO:HD3	2.12	0.49
1:A:192:THR:HG22	1:A:204:ARG:HB3	1.92	0.49
1:A:203:LEU:O	1:A:248:ALA:HA	2.13	0.49
1:A:251:VAL:HG22	1:A:259:TYR:CZ	2.47	0.49
5:E:30:SER:O	5:E:70:ARG:NH2	2.46	0.49
1:A:26[A]:SER:OG	1:A:38:PHE:HB3	2.12	0.49
5:E:7:THR:OG1	5:E:25:ASN:HB2	2.13	0.48
1:A:15:SER:HA	1:A:22:PRO:HB3	1.96	0.48
1:A:11:TYR:HB2	1:A:99:SER:HB2	1.96	0.48
1:A:121:ASP:HB3	2:B:0:MET:HG3	1.94	0.48
1:A:163:GLU:OE1	6:A:301:HOH:O	2.20	0.48
1:A:191:VAL:CG1	1:A:276:TRP:HA	2.39	0.47
1:A:66:THR:HG22	1:A:70:LYS:HE3	1.95	0.47
4:D:143:VAL:HG11	5:E:136:VAL:HG11	1.95	0.47
2:B:17:ASN:HD21	2:B:97:ARG:NH1	2.10	0.47
4:D:126:VAL:HG23	4:D:236:ALA:HB3	1.96	0.47
5:E:83:ARG:HB2	5:E:83:ARG:CZ	2.45	0.46
4:D:30:HIS:CD2	4:D:97:LEU:HB2	2.50	0.46
5:E:157:TYR:O	5:E:178:ALA:HA	2.16	0.46
1:A:229:ASP:N	1:A:229:ASP:OD1	2.48	0.45
5:E:132:SER:OG	5:E:134:LYS:HE3	2.15	0.45
1:A:192:THR:CG2	1:A:194:HIS:HE1	2.29	0.45
1:A:226:GLN:HB2	1:A:230:THR:OG1	2.16	0.45
5:E:130:LYS:HE3	5:E:130:LYS:HB3	1.69	0.45
5:E:187:CYS:C	5:E:189:ASN:H	2.20	0.45
1:A:3:GLY:HA2	1:A:107:PRO:HB3	2.00	0.44
1:A:171:ARG:O	1:A:171:ARG:HD3	2.17	0.44
4:D:222:TRP:CE2	4:D:224:GLN:HB2	2.52	0.44
2:B:49:VAL:HG22	2:B:68:THR:HB	2.00	0.44
4:D:25:LYS:HG2	4:D:74:SER:O	2.17	0.43
2:B:87:LEU:HD22	2:B:91:LYS:HE3	1.99	0.43
4:D:16:MET:HE3	4:D:85:PRO:HD3	2.00	0.43
1:A:60[B]:GLU:H	1:A:60[B]:GLU:HG3	1.53	0.42
1:A:222:ASP:OD1	1:A:258:ARG:HB3	2.19	0.42
1:A:108:ASP:OD1	1:A:109:GLY:N	2.52	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:E:171:PHE:CE2	5:E:173:SER:HB3	2.55	0.42
2:B:23:LEU:O	2:B:67:TYR:HA	2.20	0.42
4:D:48:ILE:HD13	4:D:68:ALA:HB3	2.01	0.42
4:D:139:LYS:HE3	5:E:169:MET:CE	2.47	0.42
5:E:111:MET:HE2	5:E:111:MET:HB3	1.98	0.42
1:A:78[A]:GLU:HB3	3:C:8:MET:HE1	2.02	0.42
5:E:128:ASP:HB3	5:E:131:SER:HA	2.01	0.42
4:D:46:LEU:HD22	5:E:101:MET:HG3	2.02	0.41
2:B:21:ASN:OD1	2:B:22:PHE:N	2.44	0.41
3:C:8:MET:HE3	4:D:52:ASN:HB3	2.02	0.41
4:D:129:PRO:HD2	4:D:200:TRP:CZ2	2.56	0.41
1:A:192:THR:HG21	1:A:206:TRP:HE1	1.85	0.41
5:E:161:LYS:HB2	5:E:161:LYS:HE3	1.70	0.41
5:E:8:GLN:HB3	5:E:108:THR:OG1	2.22	0.40
5:E:131:SER:O	5:E:131:SER:OG	2.27	0.40
1:A:226:GLN:OE1	1:A:226:GLN:HA	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:48:LYS:NZ	5:E:151:SER:O[3_544]	2.19	0.01

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	281/279 (101%)	266 (95%)	13 (5%)	2 (1%)	22	32
2	B	98/100 (98%)	95 (97%)	3 (3%)	0	100	100
3	C	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
4	D	239/244 (98%)	232 (97%)	6 (2%)	1 (0%)	34	48

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
5	E	200/204 (98%)	180 (90%)	17 (8%)	3 (2%)	10	14
All	All	825/836 (99%)	779 (94%)	40 (5%)	6 (1%)	22	32

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	D	44	LEU
1	A	223	GLY
5	E	188	ALA
1	A	4	SER
5	E	78	PRO
5	E	203	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	241/236 (102%)	229 (95%)	12 (5%)	24	38
2	B	95/95 (100%)	90 (95%)	5 (5%)	22	36
3	C	9/9 (100%)	8 (89%)	1 (11%)	6	8
4	D	211/213 (99%)	205 (97%)	6 (3%)	43	61
5	E	179/180 (99%)	170 (95%)	9 (5%)	24	38
All	All	735/733 (100%)	702 (96%)	33 (4%)	28	42

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	GLN
1	A	60[A]	GLU
1	A	60[B]	GLU
1	A	136	THR
1	A	190	HIS
1	A	192	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	194	HIS
1	A	196	ILE
1	A	198	ASP
1	A	199	HIS
1	A	218	THR
1	A	229	ASP
2	B	0	MET
2	B	36	GLU
2	B	70	PHE
2	B	75	LYS
2	B	92	ILE
3	C	9	LEU
4	D	40	MET
4	D	81	GLN
4	D	130	SER
4	D	163	LYS
4	D	192	ARG
4	D	241	ARG
5	E	5	LYS
5	E	32	ASN
5	E	79	HIS
5	E	83	ARG
5	E	131	SER
5	E	143	ASP
5	E	152	LYS
5	E	155	ASP
5	E	202	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	190	HIS
1	A	226	GLN
2	B	2	GLN
2	B	17	ASN
4	D	179	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	277/279 (99%)	0.38	26 (9%) 8 6	31, 55, 121, 147	0
2	B	100/100 (100%)	-0.01	1 (1%) 82 81	37, 49, 74, 102	0
3	C	9/9 (100%)	-0.12	0 100 100	36, 38, 42, 43	0
4	D	241/244 (98%)	-0.12	1 (0%) 92 93	32, 47, 73, 114	0
5	E	202/204 (99%)	0.20	4 (1%) 65 61	33, 55, 91, 166	0
All	All	829/836 (99%)	0.14	32 (3%) 39 36	31, 52, 104, 166	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
5	E	132	SER	9.6
1	A	254	GLY	5.3
1	A	251	VAL	4.4
1	A	201	ALA	4.1
1	A	253	SER	3.9
1	A	193	HIS	3.8
2	B	0	MET	3.7
1	A	196	ILE	3.6
4	D	243	ASP	3.5
1	A	250	VAL	3.5
1	A	203	LEU	3.3
1	A	199	HIS	3.3
1	A	230	THR	3.3
1	A	229	ASP	3.3
5	E	131	SER	3.2
5	E	194	SER	3.2
1	A	278	PRO	3.0
1	A	256	GLU	3.0
1	A	275	ARG	2.6
1	A	226	GLN	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
5	E	133	ASP	2.4
1	A	221	ARG	2.4
1	A	259	TYR	2.3
1	A	276	TRP	2.3
1	A	219	TRP	2.3
1	A	200	GLU	2.2
1	A	262	HIS	2.2
1	A	228	GLN	2.2
1	A	248	ALA	2.2
1	A	252	PRO	2.1
1	A	279	SER	2.0
1	A	249	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.