



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2021 – 02:31 PM EDT

PDB ID : 3E5V  
Title : Crystal Structure Analysis of eqFP611 Double Mutant T122R, N143S  
Authors : Nar, H.; Nienhaus, K.; Nienhaus, U.; Wiedenmann, J.  
Deposited on : 2008-08-14  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	242	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1924 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Red fluorescent protein eqFP611.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	229	Total	C	N	O	S	0	1	0
			1849	1178	310	342	19			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	63	NRQ	MET	chromophore	UNP Q8ISF8
A	63	NRQ	TYR	chromophore	UNP Q8ISF8
A	63	NRQ	GLY	chromophore	UNP Q8ISF8
A	122	ARG	THR	engineered mutation	UNP Q8ISF8
A	143	SER	ASN	engineered mutation	UNP Q8ISF8

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	75	Total	O	0	0
			75	75		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Red fluorescent protein eqFP611

Chain A: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	62.54Å 62.54Å 208.02Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	13.89 – 2.10 13.84 – 2.10	Depositor EDS
% Data completeness (in resolution range)	93.3 (13.89-2.10) 93.4 (13.84-2.10)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.71 (at 2.10Å)	Xtriage
Refinement program	TNT, BUSTER-TNT 2.1.1	Depositor
R, $R_{free}$	0.226 , 0.278 0.241 , 0.289	Depositor DCC
$R_{free}$ test set	695 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	45.7	Xtriage
Anisotropy	0.102	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 66.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.39$ , $\langle L^2 \rangle = 0.22$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	1924	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.14% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NRQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.28	0/1846	0.50	0/2477

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1849	0	1783	59	0
2	A	75	0	0	3	0
All	All	1924	0	1783	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:13:VAL:HG23	1:A:117:TYR:HB2	1.44	0.96
1:A:150:ALA:HB3	1:A:155:ARG:HG3	1.46	0.95

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:73:THR:HG22	1:A:74:LYS:H	1.41	0.84
1:A:11:MET:HE3	1:A:39:GLN:HE21	1.45	0.80
1:A:183:THR:OG1	1:A:186:ASN:HB2	1.81	0.80
1:A:11:MET:CE	1:A:39:GLN:HE21	2.01	0.74
1:A:63[B]:NRQ:SD	1:A:215:GLU:HB2	2.28	0.72
1:A:25:LYS:HB3	1:A:47:GLU:HB2	1.72	0.71
1:A:13:VAL:HG23	1:A:117:TYR:CB	2.22	0.66
1:A:73:THR:HG22	1:A:74:LYS:N	2.09	0.66
1:A:1:MET:O	1:A:6:LYS:HD3	1.96	0.65
1:A:63[B]:NRQ:O2	1:A:63[B]:NRQ:HD1	1.98	0.64
1:A:150:ALA:CB	1:A:155:ARG:HG3	2.24	0.62
1:A:1:MET:HB3	1:A:7:GLU:H	1.65	0.61
1:A:39:GLN:HE22	1:A:66:SER:CB	2.14	0.60
1:A:39:GLN:HE22	1:A:66:SER:HB3	1.67	0.59
1:A:92:ARG:HD3	1:A:174:PHE:CD2	2.39	0.57
1:A:140:TRP:CE2	1:A:162:LEU:HD13	2.40	0.57
1:A:186:ASN:HA	2:A:359:HOH:O	2.05	0.57
1:A:77:PRO:HG2	1:A:190:PRO:HA	1.86	0.56
1:A:134:GLN:HB3	1:A:136:LYS:HZ2	1.71	0.55
1:A:94:THR:HA	1:A:174:PHE:HB3	1.89	0.55
1:A:2:ASN:O	1:A:3:SER:HB2	2.07	0.53
1:A:150:ALA:HB3	1:A:155:ARG:CG	2.30	0.53
1:A:228:LEU:HB2	1:A:230:ARG:HG3	1.92	0.51
1:A:73:THR:CG2	1:A:74:LYS:H	2.20	0.50
1:A:143:SER:HB2	2:A:325:HOH:O	2.12	0.50
1:A:2:ASN:HD22	1:A:2:ASN:N	2.10	0.48
1:A:176:THR:HG21	1:A:178:TYR:CZ	2.48	0.48
1:A:155:ARG:NH2	1:A:157:TYR:OH	2.46	0.48
1:A:1:MET:HA	1:A:6:LYS:HA	1.96	0.48
1:A:53:PHE:CD1	1:A:57:ILE:HD11	2.48	0.48
1:A:1:MET:HB2	1:A:6:LYS:HB3	1.95	0.47
1:A:1:MET:CB	1:A:7:GLU:H	2.27	0.47
1:A:27:THR:HG23	1:A:46:VAL:CG1	2.45	0.46
1:A:39:GLN:NE2	1:A:66:SER:CB	2.78	0.46
1:A:34:PRO:HA	1:A:69:PHE:HA	1.98	0.46
1:A:63[A]:NRQ:HE1	1:A:199:LEU:HB2	1.96	0.46
1:A:54:ALA:O	1:A:57:ILE:HG12	2.17	0.45
1:A:42:ARG:HG3	1:A:212:VAL:HG22	1.99	0.45
1:A:95:ARG:HG3	2:A:348:HOH:O	2.17	0.44
1:A:196:ASP:O	1:A:217:ALA:HA	2.18	0.44
1:A:50:PRO:HD3	1:A:209:MET:SD	2.58	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:41:MET:HB2	1:A:63[A]:NRQ:HE2A	1.99	0.43
1:A:41:MET:HB2	1:A:63[A]:NRQ:CE	2.48	0.43
1:A:90:TRP:HA	1:A:177:THR:O	2.18	0.43
1:A:66:SER:HG	1:A:69:PHE:HD1	1.67	0.43
1:A:110:LEU:HD12	1:A:114:CYS:O	2.19	0.43
1:A:158:SER:HB2	1:A:174:PHE:CE2	2.54	0.42
1:A:185:GLU:H	1:A:185:GLU:HG3	1.53	0.42
1:A:120:LYS:HB2	1:A:120:LYS:HE3	1.53	0.42
1:A:206:ASP:HB3	1:A:209:MET:HB2	2.01	0.42
1:A:1:MET:HB2	1:A:6:LYS:HD3	2.01	0.42
1:A:35:TYR:CE2	1:A:81:LYS:HE2	2.55	0.42
1:A:1:MET:HA	1:A:7:GLU:H	1.85	0.41
1:A:219:ALA:C	1:A:220:LYS:HG3	2.41	0.41
1:A:22:TYR:HE2	1:A:48:GLY:O	2.04	0.41
1:A:23:GLN:HE21	1:A:23:GLN:HB3	1.64	0.41
1:A:149:PRO:HG3	1:A:190:PRO:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	224/242 (93%)	219 (98%)	4 (2%)	1 (0%)	34 32

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	3	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	196/207 (95%)	177 (90%)	19 (10%)	8 5

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	11	MET
1	A	13	VAL
1	A	16	GLU
1	A	27	THR
1	A	31	ASP
1	A	42	ARG
1	A	71	LYS
1	A	120	LYS
1	A	128	SER
1	A	136	LYS
1	A	172	CYS
1	A	177	THR
1	A	185	GLU
1	A	201	ARG
1	A	206	ASP
1	A	215	GLU
1	A	220	LYS
1	A	226	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2	ASN
1	A	23	GLN
1	A	39	GLN
1	A	159	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	NRQ	A	63[B]	1	23,24,25	1.92	3 (13%)	23,32,34	3.84	9 (39%)
1	NRQ	A	63[A]	1	23,24,25	1.99	3 (13%)	23,32,34	4.21	9 (39%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	NRQ	A	63[B]	1	-	3/9/31/32	0/2/2/2
1	NRQ	A	63[A]	1	-	2/9/31/32	0/2/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	63[B]	NRQ	CA1-N1	6.77	1.43	1.27
1	A	63[A]	NRQ	CA1-N1	6.75	1.43	1.27
1	A	63[B]	NRQ	CB2-CA2	4.93	1.39	1.35
1	A	63[A]	NRQ	CB2-CA2	4.67	1.39	1.35
1	A	63[A]	NRQ	CA2-C2	-3.04	1.45	1.48
1	A	63[B]	NRQ	C1-N2	-2.05	1.28	1.33

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	63[A]	NRQ	O2-C2-CA2	-15.82	122.08	130.96
1	A	63[B]	NRQ	O2-C2-CA2	-12.10	124.17	130.96
1	A	63[A]	NRQ	CA2-C2-N3	7.65	106.99	103.37
1	A	63[B]	NRQ	C2-CA2-N2	-6.73	104.22	108.93
1	A	63[B]	NRQ	CA2-C2-N3	6.60	106.49	103.37
1	A	63[B]	NRQ	CB2-CA2-C2	6.43	129.95	122.28
1	A	63[A]	NRQ	C2-CA2-N2	-5.42	105.13	108.93
1	A	63[B]	NRQ	CA2-N2-C1	4.29	112.18	104.33
1	A	63[A]	NRQ	CA2-N2-C1	3.57	110.87	104.33
1	A	63[A]	NRQ	CG2-CB2-CA2	-3.35	125.84	129.94
1	A	63[A]	NRQ	O2-C2-N3	3.30	130.91	124.35
1	A	63[B]	NRQ	CG2-CB2-CA2	2.97	133.59	129.94
1	A	63[A]	NRQ	CB2-CA2-N2	2.93	132.89	128.83
1	A	63[A]	NRQ	N3-C1-N2	-2.74	109.67	113.28
1	A	63[B]	NRQ	O2-C2-N3	2.51	129.34	124.35
1	A	63[B]	NRQ	O3-C3-CA3	-2.39	119.16	126.39
1	A	63[A]	NRQ	O3-C3-CA3	-2.32	119.38	126.39
1	A	63[B]	NRQ	CB2-CA2-N2	-2.15	125.84	128.83

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	63[A]	NRQ	C1-CA1-CB1-CG1
1	A	63[B]	NRQ	C2-CA2-CB2-CG2
1	A	63[A]	NRQ	CB1-CG1-SD-CE
1	A	63[B]	NRQ	N2-CA2-CB2-CG2
1	A	63[B]	NRQ	C3-CA3-N3-C1

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	63[B]	NRQ	2	0
1	A	63[A]	NRQ	3	0

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	228/242 (94%)	0.30	14 (6%) 21 26	35, 47, 67, 82	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	184	VAL	8.4
1	A	22	TYR	5.4
1	A	3	SER	5.3
1	A	2	ASN	5.2
1	A	1	MET	4.0
1	A	167	GLY	3.6
1	A	185	GLU	2.8
1	A	188	LYS	2.8
1	A	229	GLY	2.6
1	A	187	PHE	2.6
1	A	129	ASN	2.3
1	A	151	ASP	2.1
1	A	166	GLY	2.1
1	A	111	GLU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	NRQ	A	63[A]	23/24	0.90	0.17	44,47,49,50	23
1	NRQ	A	63[B]	23/24	0.90	0.17	34,40,46,48	23

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.