



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 25, 2020 – 03:18 PM BST

PDB ID : 4E5G
Title : Crystal structure of avian influenza virus PAN bound to compound 2
Authors : DuBois, R.M.; Slavish, P.J.; Webb, T.R.; White, S.W.
Deposited on : 2012-03-14
Resolution : 2.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1426 (2.66-2.62)
Clashscore	141614	1472 (2.66-2.62)
Ramachandran outliers	138981	1446 (2.66-2.62)
Sidechain outliers	138945	1446 (2.66-2.62)
RSRZ outliers	127900	1408 (2.66-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	187	<div> <div>2%</div> <div> <div></div> <div>83%</div> <div>13%</div> <div>..</div> </div> </div>
1	B	187	<div> <div>2%</div> <div> <div></div> <div>81%</div> <div>14%</div> <div>..</div> </div> </div>
1	C	187	<div> <div>0%</div> <div> <div></div> <div>79%</div> <div>16%</div> <div>..</div> </div> </div>
1	D	187	<div> <div>2%</div> <div> <div></div> <div>81%</div> <div>14%</div> <div>..</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	XI7	D	306	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6111 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polymerase protein PA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	181	Total	C	N	O	S	0	0	0
			1489	937	257	284	11			
1	B	181	Total	C	N	O	S	0	0	0
			1489	937	257	284	11			
1	C	181	Total	C	N	O	S	0	0	0
			1489	937	257	284	11			
1	D	179	Total	C	N	O	S	0	0	0
			1475	929	255	280	11			

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	50A	GLY	-	LINKER	UNP Q5EP34
A	50B	GLY	-	LINKER	UNP Q5EP34
A	50C	SER	-	LINKER	UNP Q5EP34
A	196A	ALA	-	EXPRESSION TAG	UNP Q5EP34
A	196B	ALA	-	EXPRESSION TAG	UNP Q5EP34
A	196C	ALA	-	EXPRESSION TAG	UNP Q5EP34
A	196D	GLU	-	EXPRESSION TAG	UNP Q5EP34
A	197	LEU	-	EXPRESSION TAG	UNP Q5EP34
A	198	ALA	-	EXPRESSION TAG	UNP Q5EP34
A	199	LEU	-	EXPRESSION TAG	UNP Q5EP34
A	200	VAL	-	EXPRESSION TAG	UNP Q5EP34
A	201	PRO	-	EXPRESSION TAG	UNP Q5EP34
A	202	ARG	-	EXPRESSION TAG	UNP Q5EP34
B	50A	GLY	-	LINKER	UNP Q5EP34
B	50B	GLY	-	LINKER	UNP Q5EP34
B	50C	SER	-	LINKER	UNP Q5EP34
B	196A	ALA	-	EXPRESSION TAG	UNP Q5EP34
B	196B	ALA	-	EXPRESSION TAG	UNP Q5EP34
B	196C	ALA	-	EXPRESSION TAG	UNP Q5EP34
B	196D	GLU	-	EXPRESSION TAG	UNP Q5EP34
B	197	LEU	-	EXPRESSION TAG	UNP Q5EP34

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	198	ALA	-	EXPRESSION TAG	UNP Q5EP34
B	199	LEU	-	EXPRESSION TAG	UNP Q5EP34
B	200	VAL	-	EXPRESSION TAG	UNP Q5EP34
B	201	PRO	-	EXPRESSION TAG	UNP Q5EP34
B	202	ARG	-	EXPRESSION TAG	UNP Q5EP34
C	50A	GLY	-	LINKER	UNP Q5EP34
C	50B	GLY	-	LINKER	UNP Q5EP34
C	50C	SER	-	LINKER	UNP Q5EP34
C	196A	ALA	-	EXPRESSION TAG	UNP Q5EP34
C	196B	ALA	-	EXPRESSION TAG	UNP Q5EP34
C	196C	ALA	-	EXPRESSION TAG	UNP Q5EP34
C	196D	GLU	-	EXPRESSION TAG	UNP Q5EP34
C	197	LEU	-	EXPRESSION TAG	UNP Q5EP34
C	198	ALA	-	EXPRESSION TAG	UNP Q5EP34
C	199	LEU	-	EXPRESSION TAG	UNP Q5EP34
C	200	VAL	-	EXPRESSION TAG	UNP Q5EP34
C	201	PRO	-	EXPRESSION TAG	UNP Q5EP34
C	202	ARG	-	EXPRESSION TAG	UNP Q5EP34
D	50A	GLY	-	LINKER	UNP Q5EP34
D	50B	GLY	-	LINKER	UNP Q5EP34
D	50C	SER	-	LINKER	UNP Q5EP34
D	196A	ALA	-	EXPRESSION TAG	UNP Q5EP34
D	196B	ALA	-	EXPRESSION TAG	UNP Q5EP34
D	197	ALA	-	EXPRESSION TAG	UNP Q5EP34
D	198	GLU	-	EXPRESSION TAG	UNP Q5EP34
D	199	LEU	-	EXPRESSION TAG	UNP Q5EP34
D	200	ALA	-	EXPRESSION TAG	UNP Q5EP34
D	201	LEU	-	EXPRESSION TAG	UNP Q5EP34
D	202	VAL	-	EXPRESSION TAG	UNP Q5EP34
D	203	PRO	-	EXPRESSION TAG	UNP Q5EP34
D	204	ARG	-	EXPRESSION TAG	UNP Q5EP34

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Mn	0	0
			2	2		
3	A	2	Total	Mn	0	0
			2	2		
3	D	2	Total	Mn	0	0
			2	2		
3	C	2	Total	Mn	0	0
			2	2		

- Molecule 4 is 2-4-DIOXO-4-PHENYLBUTANOIC ACID (three-letter code: XI7) (formula: $C_{10}H_8O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			14	10	4		
4	A	1	Total	C	O	0	0
			14	10	4		
4	B	1	Total	C	O	0	0
			14	10	4		
4	B	1	Total	C	O	0	0
			14	10	4		
4	C	1	Total	C	O	0	0
			14	10	4		
4	C	1	Total	C	O	0	0
			14	10	4		
4	D	1	Total	C	O	0	0
			14	10	4		
4	D	1	Total	C	O	0	0
			14	10	4		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	4	Total	O	0	0
			4	4		
5	B	1	Total	O	0	0
			1	1		

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	1	Total	O	0	0
			1	1		
5	D	3	Total	O	0	0
			3	3		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Polymerase protein PA





- Molecule 1: Polymerase protein PA



- Molecule 1: Polymerase protein PA



- Molecule 1: Polymerase protein PA



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	126.89Å 133.66Å 126.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.84 – 2.65 44.84 – 2.65	Depositor EDS
% Data completeness (in resolution range)	99.7 (44.84-2.65) 99.8 (44.84-2.65)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.34 (at 2.65Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.243 , 0.285 0.244 , 0.284	Depositor DCC
R_{free} test set	1599 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	50.9	Xtriage
Anisotropy	0.072	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 34.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.003 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6111	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 28.08 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.9629e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN, SO4, XI7

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.45	0/1519	0.58	0/2040
1	B	0.43	0/1519	0.59	0/2040
1	C	0.44	0/1519	0.56	0/2040
1	D	0.43	0/1505	0.57	0/2021
All	All	0.44	0/6062	0.58	0/8141

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1489	0	1450	24	0
1	B	1489	0	1450	30	0
1	C	1489	0	1450	33	0
1	D	1475	0	1439	32	0
2	A	10	0	0	0	0
2	B	10	0	0	0	0
2	C	10	0	0	0	0
2	D	10	0	0	0	0
3	A	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
4	A	28	0	14	1	0
4	B	28	0	14	2	0
4	C	28	0	14	2	0
4	D	28	0	14	2	0
5	A	4	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	3	0	0	0	0
All	All	6111	0	5845	110	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 110 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:73:LYS:H	1:D:73:LYS:CD	1.57	1.17
1:B:73:LYS:H	1:B:73:LYS:HE3	1.13	1.12
1:C:73:LYS:CE	1:C:73:LYS:H	1.64	1.09
1:A:73:LYS:HE3	1:A:73:LYS:H	0.90	1.06
1:A:73:LYS:HE3	1:A:73:LYS:N	1.73	1.03

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	179/187 (96%)	175 (98%)	4 (2%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	179/187 (96%)	176 (98%)	3 (2%)	0	100	100
1	C	179/187 (96%)	174 (97%)	5 (3%)	0	100	100
1	D	177/187 (95%)	174 (98%)	3 (2%)	0	100	100
All	All	714/748 (96%)	699 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	160/165 (97%)	157 (98%)	3 (2%)	57	74
1	B	160/165 (97%)	155 (97%)	5 (3%)	40	58
1	C	160/165 (97%)	156 (98%)	4 (2%)	47	66
1	D	159/165 (96%)	157 (99%)	2 (1%)	69	82
All	All	639/660 (97%)	625 (98%)	14 (2%)	52	70

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	104	LYS
1	B	124	ARG
1	C	153	GLU
1	B	96	ASN
1	C	96	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	96	ASN
1	D	92	ASN
1	C	92	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	92	ASN
1	C	41	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 24 ligands modelled in this entry, 8 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	302	-	4,4,4	0.15	0	6,6,6	0.11	0
4	XI7	A	306	-	11,14,14	2.49	2 (18%)	13,18,18	0.95	1 (7%)
4	XI7	A	305	3	11,14,14	2.48	2 (18%)	13,18,18	0.78	0
4	XI7	C	306	-	11,14,14	2.47	2 (18%)	13,18,18	1.01	1 (7%)
2	SO4	B	301	-	4,4,4	0.14	0	6,6,6	0.16	0
2	SO4	A	301	-	4,4,4	0.13	0	6,6,6	0.15	0
2	SO4	D	301	-	4,4,4	0.14	0	6,6,6	0.13	0
2	SO4	C	302	-	4,4,4	0.13	0	6,6,6	0.09	0
2	SO4	A	302	-	4,4,4	0.14	0	6,6,6	0.12	0
2	SO4	C	301	-	4,4,4	0.15	0	6,6,6	0.12	0
4	XI7	D	305	3	11,14,14	2.48	2 (18%)	13,18,18	0.82	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	XI7	C	305	3	11,14,14	2.45	2 (18%)	13,18,18	0.63	0
4	XI7	B	305	3	11,14,14	2.47	2 (18%)	13,18,18	0.89	0
4	XI7	B	306	-	11,14,14	2.41	2 (18%)	13,18,18	0.84	0
4	XI7	D	306	-	11,14,14	2.54	2 (18%)	13,18,18	1.38	2 (15%)
2	SO4	D	302	-	4,4,4	0.14	0	6,6,6	0.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	XI7	A	306	-	-	0/8/12/12	0/1/1/1
4	XI7	C	306	-	-	0/8/12/12	0/1/1/1
4	XI7	B	306	-	-	0/8/12/12	0/1/1/1
4	XI7	D	305	3	-	0/8/12/12	0/1/1/1
4	XI7	C	305	3	-	0/8/12/12	0/1/1/1
4	XI7	B	305	3	-	0/8/12/12	0/1/1/1
4	XI7	A	305	3	-	0/8/12/12	0/1/1/1
4	XI7	D	306	-	-	0/8/12/12	0/1/1/1

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	306	XI7	O8-C9	7.19	1.33	1.22
4	A	306	XI7	O8-C9	7.05	1.33	1.22
4	C	306	XI7	O8-C9	7.03	1.33	1.22
4	D	305	XI7	O8-C9	7.02	1.33	1.22
4	B	305	XI7	O8-C9	7.01	1.33	1.22

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	306	XI7	C7-C8-C9	2.47	126.42	114.90
4	C	306	XI7	C7-C8-C9	2.34	125.79	114.90
4	D	306	XI7	O14-C7-C8	2.09	122.91	120.56
4	A	306	XI7	C7-C8-C9	2.08	124.59	114.90

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	306	XI7	1	0
4	C	306	XI7	2	0
4	B	306	XI7	2	0
4	D	306	XI7	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	181/187 (96%)	0.17	3 (1%) 70 67	29, 46, 68, 84	0
1	B	181/187 (96%)	0.18	3 (1%) 70 67	31, 47, 73, 86	0
1	C	181/187 (96%)	0.14	1 (0%) 89 88	29, 46, 71, 76	0
1	D	179/187 (95%)	0.19	4 (2%) 62 58	32, 46, 68, 73	0
All	All	722/748 (96%)	0.17	11 (1%) 73 71	29, 46, 70, 86	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	124	ARG	3.6
1	D	141	GLU	3.5
1	D	125	ARG	2.9
1	D	124	ARG	2.7
1	C	74	HIS	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	XI7	D	306	14/14	0.76	0.42	62,65,66,66	0
2	SO4	B	301	5/5	0.82	0.23	101,101,102,102	0
4	XI7	C	306	14/14	0.83	0.42	67,69,73,74	0
2	SO4	A	302	5/5	0.84	0.32	108,108,108,108	0
2	SO4	D	302	5/5	0.85	0.23	116,116,116,116	0
4	XI7	A	306	14/14	0.85	0.35	64,67,67,67	0
2	SO4	B	302	5/5	0.87	0.31	99,99,99,100	0
2	SO4	C	302	5/5	0.87	0.24	90,90,91,91	0
4	XI7	B	306	14/14	0.88	0.31	61,66,69,69	0
2	SO4	D	301	5/5	0.89	0.24	100,100,101,101	0
4	XI7	D	305	14/14	0.89	0.27	53,57,62,62	0
2	SO4	C	301	5/5	0.91	0.21	87,88,88,88	0
4	XI7	A	305	14/14	0.93	0.26	53,54,55,55	0
2	SO4	A	301	5/5	0.93	0.12	82,83,83,83	0
4	XI7	B	305	14/14	0.94	0.20	50,54,57,57	0
4	XI7	C	305	14/14	0.95	0.24	42,48,52,52	0
3	MN	D	304	1/1	0.95	0.16	56,56,56,56	0
3	MN	C	304	1/1	0.95	0.20	53,53,53,53	0
3	MN	D	303	1/1	0.96	0.18	44,44,44,44	0
3	MN	C	303	1/1	0.97	0.18	42,42,42,42	0
3	MN	B	304	1/1	0.97	0.18	38,38,38,38	0
3	MN	B	303	1/1	0.98	0.17	59,59,59,59	0
3	MN	A	303	1/1	0.98	0.23	57,57,57,57	0
3	MN	A	304	1/1	0.99	0.17	37,37,37,37	0

6.5 Other polymers ⓘ

There are no such residues in this entry.