



wwPDB X-ray Structure Validation Summary Report ⓘ

May 22, 2020 – 01:00 am BST

PDB ID : 4E9M
Title : NOD1 card domain with three disulfide-clinched, domain-swapped dimers in the asymmetric unit
Authors : Berry, E.A.; Huang, L.S.; Walker-Kopp, N.
Deposited on : 2012-03-21
Resolution : 2.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	144	<div> <div>4%</div> <div> <div>47%</div> <div>19%</div> <div>•</div> <div>33%</div> </div> </div>
1	B	144	<div> <div>%</div> <div> <div>49%</div> <div>15%</div> <div>•</div> <div>34%</div> </div> </div>
1	C	144	<div> <div>6%</div> <div> <div>44%</div> <div>19%</div> <div>•</div> <div>37%</div> </div> </div>
1	D	144	<div> <div>5%</div> <div> <div>45%</div> <div>19%</div> <div>•</div> <div>34%</div> </div> </div>
1	E	144	<div> <div>13%</div> <div> <div>40%</div> <div>20%</div> <div>•</div> <div>40%</div> </div> </div>
1	F	144	<div> <div>40%</div> <div> <div>30%</div> <div>33%</div> <div>•</div> <div>35%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FLC	D	1101	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4641 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nucleotide-binding oligomerization domain-containing protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	97	Total	C	N	O	S	0	0	0
			786	502	131	150	3			
1	B	95	Total	C	N	O	S	0	0	0
			771	494	129	145	3			
1	C	91	Total	C	N	O	S	0	0	0
			742	474	125	140	3			
1	D	95	Total	C	N	O	S	0	0	0
			771	494	129	145	3			
1	E	87	Total	C	N	O	S	0	0	0
			705	446	120	136	3			
1	F	94	Total	C	N	O	S	0	0	0
			765	491	128	143	3			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	MET	-	INITIATING METHIONINE	UNP Q9Y239
A	-4	HIS	-	EXPRESSION TAG	UNP Q9Y239
A	-3	HIS	-	EXPRESSION TAG	UNP Q9Y239
A	-2	HIS	-	EXPRESSION TAG	UNP Q9Y239
A	-1	HIS	-	EXPRESSION TAG	UNP Q9Y239
A	0	HIS	-	EXPRESSION TAG	UNP Q9Y239
A	1	HIS	-	EXPRESSION TAG	UNP Q9Y239
B	-5	MET	-	INITIATING METHIONINE	UNP Q9Y239
B	-4	HIS	-	EXPRESSION TAG	UNP Q9Y239
B	-3	HIS	-	EXPRESSION TAG	UNP Q9Y239
B	-2	HIS	-	EXPRESSION TAG	UNP Q9Y239
B	-1	HIS	-	EXPRESSION TAG	UNP Q9Y239
B	0	HIS	-	EXPRESSION TAG	UNP Q9Y239
B	1	HIS	-	EXPRESSION TAG	UNP Q9Y239
C	-5	MET	-	INITIATING METHIONINE	UNP Q9Y239
C	-4	HIS	-	EXPRESSION TAG	UNP Q9Y239

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-3	HIS	-	EXPRESSION TAG	UNP Q9Y239
C	-2	HIS	-	EXPRESSION TAG	UNP Q9Y239
C	-1	HIS	-	EXPRESSION TAG	UNP Q9Y239
C	0	HIS	-	EXPRESSION TAG	UNP Q9Y239
C	1	HIS	-	EXPRESSION TAG	UNP Q9Y239
D	-5	MET	-	INITIATING METHIONINE	UNP Q9Y239
D	-4	HIS	-	EXPRESSION TAG	UNP Q9Y239
D	-3	HIS	-	EXPRESSION TAG	UNP Q9Y239
D	-2	HIS	-	EXPRESSION TAG	UNP Q9Y239
D	-1	HIS	-	EXPRESSION TAG	UNP Q9Y239
D	0	HIS	-	EXPRESSION TAG	UNP Q9Y239
D	1	HIS	-	EXPRESSION TAG	UNP Q9Y239
E	-5	MET	-	INITIATING METHIONINE	UNP Q9Y239
E	-4	HIS	-	EXPRESSION TAG	UNP Q9Y239
E	-3	HIS	-	EXPRESSION TAG	UNP Q9Y239
E	-2	HIS	-	EXPRESSION TAG	UNP Q9Y239
E	-1	HIS	-	EXPRESSION TAG	UNP Q9Y239
E	0	HIS	-	EXPRESSION TAG	UNP Q9Y239
E	1	HIS	-	EXPRESSION TAG	UNP Q9Y239
F	-5	MET	-	INITIATING METHIONINE	UNP Q9Y239
F	-4	HIS	-	EXPRESSION TAG	UNP Q9Y239
F	-3	HIS	-	EXPRESSION TAG	UNP Q9Y239
F	-2	HIS	-	EXPRESSION TAG	UNP Q9Y239
F	-1	HIS	-	EXPRESSION TAG	UNP Q9Y239
F	0	HIS	-	EXPRESSION TAG	UNP Q9Y239
F	1	HIS	-	EXPRESSION TAG	UNP Q9Y239

- Molecule 2 is CITRATE ANION (three-letter code: FLC) (formula: C₆H₅O₇).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	D	1	Total	C	O	0	0
			13	6	7		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	21	Total	O	0	0
			21	21		
3	B	15	Total	O	0	0
			15	15		
3	C	28	Total	O	0	0
			28	28		
3	D	17	Total	O	0	0
			17	17		
3	E	5	Total	O	0	0
			5	5		
3	F	2	Total	O	0	0
			2	2		

- Molecule 1: Nucleotide-binding oligomerization domain-containing protein 1







● Molecule 1: Nucleotide-binding oligomerization domain-containing protein 1



● Molecule 1: Nucleotide-binding oligomerization domain-containing protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	82.85Å 85.33Å 113.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.84 – 2.15 24.84 – 2.15	Depositor EDS
% Data completeness (in resolution range)	76.8 (24.84-2.15) 76.9 (24.84-2.15)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.30 (at 2.15Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.245 , 0.275 0.235 , 0.264	Depositor DCC
R_{free} test set	1701 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	37.1	Xtriage
Anisotropy	0.056	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 50.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.012 for k,h,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4641	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/801	0.62	1/1087 (0.1%)
1	B	0.39	0/786	0.62	1/1067 (0.1%)
1	C	0.37	0/756	0.59	0/1027
1	D	0.36	0/786	0.56	0/1067
1	E	0.33	0/716	0.54	0/970
1	F	0.28	0/780	0.48	0/1059
All	All	0.35	0/4625	0.57	2/6277 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	39	CYS	CA-CB-SG	7.12	126.81	114.00
1	A	39	CYS	CA-CB-SG	6.93	126.47	114.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	786	0	781	22	0
1	B	771	0	770	16	0
1	C	742	0	742	33	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	771	0	770	30	0
1	E	705	0	703	23	0
1	F	765	0	765	47	0
2	D	13	0	5	2	0
3	A	21	0	0	0	0
3	B	15	0	0	0	0
3	C	28	0	0	0	0
3	D	17	0	0	0	0
3	E	5	0	0	0	0
3	F	2	0	0	1	0
All	All	4641	0	4536	142	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 142 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:27:ARG:HH11	1:A:27:ARG:HG2	1.41	0.86
1:C:74:LEU:O	1:C:78:LYS:HG2	1.83	0.77
1:E:97:TYR:HB3	1:E:100:LEU:HD12	1.69	0.75
1:D:31:VAL:HA	1:D:68:VAL:HG11	1.68	0.74
1:A:74:LEU:O	1:A:78:LYS:HG2	1.87	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	95/144 (66%)	91 (96%)	3 (3%)	1 (1%)	14	8
1	B	93/144 (65%)	90 (97%)	3 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	89/144 (62%)	85 (96%)	3 (3%)	1 (1%)	14	8
1	D	93/144 (65%)	93 (100%)	0	0	100	100
1	E	85/144 (59%)	79 (93%)	4 (5%)	2 (2%)	6	1
1	F	92/144 (64%)	77 (84%)	13 (14%)	2 (2%)	6	2
All	All	547/864 (63%)	515 (94%)	26 (5%)	6 (1%)	14	8

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	77	SER
1	E	25	SER
1	F	62	PRO
1	A	107	ILE
1	E	62	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	90/135 (67%)	88 (98%)	2 (2%)	52	55
1	B	88/135 (65%)	85 (97%)	3 (3%)	37	35
1	C	85/135 (63%)	84 (99%)	1 (1%)	71	76
1	D	88/135 (65%)	83 (94%)	5 (6%)	20	16
1	E	81/135 (60%)	80 (99%)	1 (1%)	71	76
1	F	87/135 (64%)	84 (97%)	3 (3%)	37	35
All	All	519/810 (64%)	504 (97%)	15 (3%)	42	42

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	38	GLN
1	D	64	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	61	CYS
1	C	33	HIS
1	E	33	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	91	GLN
1	D	92	GLN
1	F	43	ASN
1	D	19	HIS
1	D	38	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FLC	D	1101	-	3,12,12	1.20	0	3,17,17	1.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FLC	D	1101	-	-	1/6/16/16	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	1101	FLC	CAC-CA-CB-OHB

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1101	FLC	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	97/144 (67%)	0.55	6 (6%) 20 27	25, 36, 52, 71	0
1	B	95/144 (65%)	0.20	2 (2%) 63 71	18, 33, 47, 68	0
1	C	91/144 (63%)	0.51	9 (9%) 7 11	19, 30, 89, 103	0
1	D	95/144 (65%)	0.61	7 (7%) 14 20	25, 37, 65, 77	0
1	E	87/144 (60%)	1.32	19 (21%) 0 0	33, 58, 78, 89	0
1	F	94/144 (65%)	2.78	57 (60%) 0 0	46, 112, 132, 141	0
All	All	559/864 (64%)	0.99	100 (17%) 1 1	18, 41, 117, 141	0

The worst 5 of 100 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	33	HIS	9.1
1	F	17	HIS	9.0
1	F	18	PRO	8.8
1	A	14	SER	8.1
1	F	57	ILE	7.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	FLC	D	1101	13/13	0.47	0.51	148,148,148,148	0

6.5 Other polymers [i](#)

There are no such residues in this entry.