



wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 07:03 am BST

PDB ID : 5E9F
Title : Structural insights of isocitrate lyases from *Magnaporthe oryzae*
Authors : Park, Y.; Cho, Y.; Lee, Y.-H.; Lee, Y.-W.; Rhee, S.
Deposited on : 2015-10-15
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	560	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 12%, green 78%, grey 9%);"></div> <div style="display: flex; justify-content: space-between; font-size: 0.8em;"> % 78% 12% 9% </div> </div>
1	B	560	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 11%, green 77%, grey 11%);"></div> <div style="display: flex; justify-content: space-between; font-size: 0.8em;"> % 77% 11% 11% </div> </div>
1	C	560	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, orange 1%, yellow 16%, green 73%, grey 11%);"></div> <div style="display: flex; justify-content: space-between; font-size: 0.8em;"> 2% 73% 16% 11% </div> </div>
1	D	560	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 3%, orange 1%, yellow 15%, green 65%, grey 19%);"></div> <div style="display: flex; justify-content: space-between; font-size: 0.8em;"> 3% 65% 15% 19% </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 14983 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Isocitrate lyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	507	Total	C	N	O	S	0	0	0
			3851	2440	662	731	18			
1	B	501	Total	C	N	O	S	0	0	0
			3853	2444	662	729	18			
1	C	496	Total	C	N	O	S	0	0	0
			3736	2368	642	709	17			
1	D	453	Total	C	N	O	S	0	0	0
			3474	2200	597	662	15			

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	548	ALA	-	expression tag	UNP P0CT06
A	549	ALA	-	expression tag	UNP P0CT06
A	550	ALA	-	expression tag	UNP P0CT06
A	551	LEU	-	expression tag	UNP P0CT06
A	552	GLU	-	expression tag	UNP P0CT06
A	553	HIS	-	expression tag	UNP P0CT06
A	554	HIS	-	expression tag	UNP P0CT06
A	555	HIS	-	expression tag	UNP P0CT06
A	556	HIS	-	expression tag	UNP P0CT06
A	557	HIS	-	expression tag	UNP P0CT06
A	558	HIS	-	expression tag	UNP P0CT06
A	559	HIS	-	expression tag	UNP P0CT06
A	560	HIS	-	expression tag	UNP P0CT06
B	548	ALA	-	expression tag	UNP P0CT06
B	549	ALA	-	expression tag	UNP P0CT06
B	550	ALA	-	expression tag	UNP P0CT06
B	551	LEU	-	expression tag	UNP P0CT06
B	552	GLU	-	expression tag	UNP P0CT06
B	553	HIS	-	expression tag	UNP P0CT06
B	554	HIS	-	expression tag	UNP P0CT06
B	555	HIS	-	expression tag	UNP P0CT06

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	556	HIS	-	expression tag	UNP P0CT06
B	557	HIS	-	expression tag	UNP P0CT06
B	558	HIS	-	expression tag	UNP P0CT06
B	559	HIS	-	expression tag	UNP P0CT06
B	560	HIS	-	expression tag	UNP P0CT06
C	548	ALA	-	expression tag	UNP P0CT06
C	549	ALA	-	expression tag	UNP P0CT06
C	550	ALA	-	expression tag	UNP P0CT06
C	551	LEU	-	expression tag	UNP P0CT06
C	552	GLU	-	expression tag	UNP P0CT06
C	553	HIS	-	expression tag	UNP P0CT06
C	554	HIS	-	expression tag	UNP P0CT06
C	555	HIS	-	expression tag	UNP P0CT06
C	556	HIS	-	expression tag	UNP P0CT06
C	557	HIS	-	expression tag	UNP P0CT06
C	558	HIS	-	expression tag	UNP P0CT06
C	559	HIS	-	expression tag	UNP P0CT06
C	560	HIS	-	expression tag	UNP P0CT06
D	548	ALA	-	expression tag	UNP P0CT06
D	549	ALA	-	expression tag	UNP P0CT06
D	550	ALA	-	expression tag	UNP P0CT06
D	551	LEU	-	expression tag	UNP P0CT06
D	552	GLU	-	expression tag	UNP P0CT06
D	553	HIS	-	expression tag	UNP P0CT06
D	554	HIS	-	expression tag	UNP P0CT06
D	555	HIS	-	expression tag	UNP P0CT06
D	556	HIS	-	expression tag	UNP P0CT06
D	557	HIS	-	expression tag	UNP P0CT06
D	558	HIS	-	expression tag	UNP P0CT06
D	559	HIS	-	expression tag	UNP P0CT06
D	560	HIS	-	expression tag	UNP P0CT06




- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Mg 1 1	0	0
2	A	1	Total Mg 1 1	0	0
2	D	1	Total Mg 1 1	0	0
2	C	1	Total Mg 1 1	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	20	Total 20	O 20	0	0
3	B	16	Total 16	O 16	0	0
3	C	17	Total 17	O 17	0	0
3	D	12	Total 12	O 12	0	0

- Molecule 1: Isocitrate lyase





• Molecule 1: Isocitrate lyase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	121.58Å 135.31Å 158.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	32.40 – 2.80 43.39 – 2.81	Depositor EDS
% Data completeness (in resolution range)	99.9 (32.40-2.80) 95.3 (43.39-2.81)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.25 (at 2.81Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.186 , 0.226 0.208 , 0.261	Depositor DCC
R_{free} test set	2000 reflections (3.12%)	wwPDB-VP
Wilson B-factor (Å ²)	46.6	Xtriage
Anisotropy	0.572	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 44.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	14983	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	1/3942 (0.0%)	0.59	0/5371
1	B	0.43	1/3942 (0.0%)	0.59	2/5361 (0.0%)
1	C	0.43	0/3822	0.57	0/5213
1	D	0.44	0/3556	0.57	0/4840
All	All	0.44	2/15262 (0.0%)	0.58	2/20785 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	C	0	1
1	D	0	1
All	All	0	3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	127	TYR	CD1-CE1	-6.59	1.29	1.39
1	B	87	CYS	CB-SG	-5.38	1.73	1.81

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	292	GLU	N-CA-C	5.50	125.85	111.00
1	B	403	TRP	CA-CB-CG	5.21	123.60	113.70

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	293	GLN	Peptide
1	C	443	ALA	Peptide
1	D	88	LEU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3851	0	3626	53	0
1	B	3853	0	3687	47	0
1	C	3736	0	3502	58	0
1	D	3474	0	3268	61	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	20	0	0	0	0
3	B	16	0	0	0	0
3	C	17	0	0	1	0
3	D	12	0	0	0	0
All	All	14983	0	14083	184	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 184 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:159:LYS:HA	1:D:162:ARG:HG3	1.58	0.84
1:A:127:TYR:OH	1:A:179:ASP:OD2	1.96	0.82
1:C:444:MET:HG2	1:C:448:GLU:HB2	1.63	0.81
1:D:64:GLN:HE21	1:D:64:GLN:N	1.79	0.80
1:D:389:ILE:HD13	1:D:420:GLY:HA3	1.68	0.76

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	501/560 (90%)	478 (95%)	18 (4%)	5 (1%)	15	44
1	B	493/560 (88%)	472 (96%)	17 (3%)	4 (1%)	19	49
1	C	488/560 (87%)	463 (95%)	21 (4%)	4 (1%)	19	49
1	D	443/560 (79%)	415 (94%)	25 (6%)	3 (1%)	22	53
All	All	1925/2240 (86%)	1828 (95%)	81 (4%)	16 (1%)	19	49

5 of 16 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	9	PRO
1	A	242	ILE
1	B	292	GLU
1	C	350	ASN
1	A	10	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	381/462 (82%)	376 (99%)	5 (1%)	69	91
1	B	389/462 (84%)	382 (98%)	7 (2%)	59	86
1	C	364/462 (79%)	361 (99%)	3 (1%)	81	94
1	D	347/462 (75%)	338 (97%)	9 (3%)	46	79
All	All	1481/1848 (80%)	1457 (98%)	24 (2%)	62	88

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	381	ILE
1	C	106	SER
1	D	405	GLU
1	B	447	ASP
1	C	79	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	313	ASN
1	D	64	GLN
1	C	413	GLN
1	B	524	GLN
1	C	524	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	507/560 (90%)	-0.17	4 (0%) 86 81	20, 34, 53, 68	0
1	B	501/560 (89%)	-0.08	7 (1%) 75 70	22, 39, 57, 82	0
1	C	496/560 (88%)	-0.08	13 (2%) 56 46	22, 37, 67, 85	0
1	D	453/560 (80%)	0.00	17 (3%) 40 30	23, 38, 73, 93	0
All	All	1957/2240 (87%)	-0.08	41 (2%) 63 54	20, 37, 62, 93	0

The worst 5 of 41 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	277	THR	4.7
1	D	276	CYS	4.6
1	D	367	PHE	4.3
1	D	275	GLY	4.2
1	D	360	LEU	3.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MG	A	601	1/1	0.82	0.38	41,41,41,41	0
2	MG	D	601	1/1	0.86	0.53	36,36,36,36	0
2	MG	B	601	1/1	0.91	0.27	30,30,30,30	0
2	MG	C	601	1/1	0.92	0.26	28,28,28,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.