



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 09:48 am BST

PDB ID : 6E90
Title : Ternary complex of human glycerol 3-phosphate dehydrogenase
Authors : Mydy, L.S.; Gulick, A.M.
Deposited on : 2018-07-31
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	349	<div> <div>86%</div> <div>12%</div> <div>..</div> </div>
1	B	349	<div> <div>91%</div> <div>7%</div> <div>..</div> </div>
1	C	349	<div> <div>93%</div> <div>6%</div> <div>.</div> </div>
1	D	349	<div> <div>%</div> <div>88%</div> <div>9%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MPD	D	402	-	-	X	-

2 Entry composition ⓘ


There are 7 unique types of molecules in this entry. The entry contains 20645 atoms, of which 9887 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycerol-3-phosphate dehydrogenase [NAD(+)], cytoplasmic.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	344	Total	C	H	N	O	S	0	1	0
			5063	1623	2520	436	465	19			
1	B	345	Total	C	H	N	O	S	0	0	0
			4954	1599	2448	422	467	18			
1	C	344	Total	C	H	N	O	S	0	1	0
			4869	1574	2404	417	455	19			
1	D	336	Total	C	H	N	O	S	0	0	0
			4829	1554	2399	407	452	17			

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	P	0	0
			70	21	26	7	14	2		
2	B	1	Total	C	H	N	O	P	0	0
			70	21	26	7	14	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	C	1	Total	C	H	N	O	P	0	0
			70	21	26	7	14	2		

- Molecule 3 is 1,3-DIHYDROXYACETONEPHOSPHATE (three-letter code: 13P) (formula: $C_3H_7O_6P$) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	O	P	0	0
			15	3	5	6	1		
3	B	1	Total	C	H	O	P	0	0
			15	3	5	6	1		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Ca	0	0
			1	1		
4	A	1	Total	Ca	0	0
			1	1		

- Molecule 5 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	0	0
			22	6	14	2		
5	D	1	Total	C	H	O	0	0
			22	6	14	2		

- Molecule 6 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	C	1	Total	O	P	0	0
			5	4	1		
6	D	1	Total	O	P	0	0
			5	4	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	155	Total 155	O 155	0	0
7	B	191	Total 191	O 191	0	0
7	C	161	Total 161	O 161	0	0
7	D	127	Total 127	O 127	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glycerol-3-phosphate dehydrogenase [NAD(+)], cytoplasmic




- Molecule 1: Glycerol-3-phosphate dehydrogenase [NAD(+)], cytoplasmic



- Molecule 1: Glycerol-3-phosphate dehydrogenase [NAD(+)], cytoplasmic



- Molecule 1: Glycerol-3-phosphate dehydrogenase [NAD(+)], cytoplasmic



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	58.94Å 79.56Å 85.06Å 63.87° 86.60° 83.61°	Depositor
Resolution (Å)	71.11 – 2.05 76.36 – 2.05	Depositor EDS
% Data completeness (in resolution range)	92.0 (71.11-2.05) 93.4 (76.36-2.05)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.49 (at 2.05Å)	Xtriage
Refinement program	PHENIX (dev_2650)	Depositor
R, R_{free}	0.177 , 0.226 0.176 , 0.220	Depositor DCC
R_{free} test set	3907 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å ²)	20.8	Xtriage
Anisotropy	0.058	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 49.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.015 for -h,-k,-k+l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	20645	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, MPD, 13P, CA, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	0/2588	0.69	0/3508
1	B	0.61	0/2549	0.66	0/3463
1	C	0.56	0/2508	0.68	0/3412
1	D	0.55	0/2469	0.64	0/3352
All	All	0.59	0/10114	0.67	0/13735

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2543	2520	2528	27	0
1	B	2506	2448	2459	14	0
1	C	2465	2404	2403	12	0
1	D	2430	2399	2398	24	0
2	A	44	26	24	0	0
2	B	44	26	25	1	0
2	C	44	26	24	0	0
3	A	10	5	5	0	0
3	B	10	5	5	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	8	14	14	3	0
5	D	8	14	14	7	0
6	C	5	0	0	0	0
6	D	5	0	0	0	0
7	A	155	0	0	4	0
7	B	191	0	0	2	0
7	C	161	0	0	1	0
7	D	127	0	0	3	0
All	All	10758	9887	9899	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:345:HIS:CE1	5:D:402:MPD:H11	2.24	0.73
1:D:347:GLU:O	5:D:402:MPD:H53	1.98	0.63
1:D:241:LEU:HD11	1:D:315:LEU:HD13	1.81	0.62
1:D:240:LYS:HA	1:D:247:VAL:HG21	1.82	0.61
5:D:402:MPD:O4	5:D:402:MPD:HM1	2.00	0.60
1:B:244:SER:C	7:B:502:HOH:O	2.41	0.59
1:A:281:GLY:HA2	7:A:544:HOH:O	2.04	0.57
1:D:247:VAL:O	1:D:247:VAL:HG23	2.04	0.57
1:D:170:ASP:HB3	1:D:173:GLN:HB3	1.86	0.57
1:C:224:LYS:HD2	1:C:266:TYR:CZ	2.39	0.56
1:A:204:LYS:HE2	1:A:260:ASP:OD2	2.06	0.56
1:C:315:LEU:HD22	1:C:318:LYS:HD2	1.89	0.55
1:D:54:ILE:O	1:D:58:HIS:HA	2.06	0.55
1:A:129:LEU:HD21	1:A:199:ILE:HD11	1.90	0.54
1:B:196:THR:CG2	1:B:251:THR:HG21	2.38	0.54
1:D:27:ALA:HB2	1:D:72:ASN:ND2	2.23	0.54
1:B:204:LYS:HE2	1:B:260:ASP:OD2	2.08	0.53
1:C:129:LEU:HD13	1:C:309:ILE:HD11	1.90	0.53
1:A:181:MET:HB2	1:A:188:ILE:CD1	2.39	0.52
1:C:132:ILE:O	1:C:136:ILE:HD12	2.10	0.52
1:B:100:LYS:O	1:B:104:GLN:OE1	2.27	0.52
1:C:275:GLU:OE2	1:C:279:ARG:NH1	2.43	0.52
1:C:246:PRO:HG3	7:D:550:HOH:O	2.12	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:345:HIS:HE1	5:D:402:MPD:H11	1.72	0.49
1:A:103:ASP:OD1	1:A:139:ARG:NH1	2.45	0.49
1:D:147:LEU:HD23	1:D:147:LEU:C	2.33	0.49
1:A:316:VAL:HG12	1:A:323:MET:SD	2.53	0.49
1:D:254:GLU:HB3	7:D:522:HOH:O	2.12	0.49
1:A:183:THR:HB	1:A:184:PRO:HD2	1.95	0.48
1:D:61:VAL:O	1:D:62:LYS:C	2.50	0.48
1:C:98:ILE:HG12	1:C:136:ILE:HD11	1.96	0.48
1:D:347:GLU:O	5:D:402:MPD:C5	2.62	0.47
1:D:321:LEU:HD12	1:D:321:LEU:O	2.14	0.47
1:A:203:LEU:HD22	1:A:306:LEU:HD11	1.95	0.47
5:A:404:MPD:C1	1:C:346:PRO:HA	2.44	0.47
1:D:234:GLU:HG2	5:D:402:MPD:H13	1.96	0.47
1:B:45:ILE:HG13	1:B:45:ILE:O	2.13	0.47
1:B:114:THR:HG22	1:B:143:PRO:HG2	1.97	0.47
1:D:116:ILE:HD11	1:D:147:LEU:HB2	1.97	0.47
1:D:52:GLU:CB	7:D:515:HOH:O	2.63	0.47
1:B:252:PHE:O	1:B:258:VAL:HG23	2.16	0.46
1:A:241:LEU:HD21	1:A:315:LEU:HD13	1.98	0.46
1:C:241:LEU:CD1	1:D:245:GLY:HA2	2.46	0.46
1:D:116:ILE:HD12	1:D:177:LEU:HD13	1.98	0.46
1:A:114:THR:HG22	1:A:143:PRO:HG2	1.98	0.46
1:A:116:ILE:HD11	1:A:147:LEU:HB2	1.99	0.45
1:A:163:GLU:OE2	5:A:404:MPD:H51	2.16	0.44
1:D:125:GLY:HA3	1:D:130:LYS:HB2	1.99	0.44
1:D:27:ALA:HB2	1:D:72:ASN:HD21	1.82	0.44
1:B:248:SER:N	7:B:517:HOH:O	2.51	0.44
1:A:240:LYS:NZ	7:A:511:HOH:O	2.51	0.43
1:C:345:HIS:ND1	1:C:346:PRO:N	2.67	0.43
1:A:337:GLU:HA	1:A:340:HIS:HD2	1.83	0.43
1:B:311:GLN:HG2	1:B:316:VAL:HG21	2.01	0.43
1:D:21:ILE:HD12	1:D:183:THR:HG21	2.00	0.43
1:A:45:ILE:HD12	1:A:50:LEU:HB2	2.01	0.42
1:D:92:VAL:O	1:D:92:VAL:HG13	2.19	0.42
1:A:20:LYS:NZ	7:A:504:HOH:O	2.45	0.42
1:A:234:GLU:OE2	1:A:345:HIS:NE2	2.47	0.42
1:A:5:LYS:NZ	1:A:33:ASP:OD2	2.47	0.42
1:A:162:CYS:HB2	7:C:599:HOH:O	2.19	0.42
1:A:241:LEU:HB3	1:A:242:PHE:CD1	2.55	0.42
1:D:55:ASN:O	1:D:58:HIS:CB	2.68	0.42
1:B:89:LEU:HD12	1:B:89:LEU:N	2.35	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:212:GLY:HA3	1:A:274:ALA:HB3	2.02	0.41
1:A:187:ARG:HB2	5:A:404:MPD:H53	2.02	0.41
1:B:40:VAL:HB	1:B:51:THR:HG22	2.02	0.41
1:A:59:GLU:HG3	1:A:61:VAL:CG1	2.50	0.41
1:A:199:ILE:HA	1:A:199:ILE:HD13	1.88	0.41
5:D:402:MPD:O4	5:D:402:MPD:CM	2.68	0.41
1:A:224:LYS:HE3	7:A:611:HOH:O	2.21	0.41
1:A:346:PRO:HA	1:A:349:MET:HE2	2.02	0.41
1:B:203:LEU:HD22	1:B:306:LEU:HD22	2.03	0.41
2:B:401:NAD:H2N	2:B:401:NAD:H2D	1.96	0.40
1:C:292:LEU:O	1:C:295:GLN:HG3	2.22	0.40
1:B:224:LYS:HG2	1:B:228:ILE:CD1	2.51	0.40
1:B:54:ILE:O	1:B:58:HIS:HA	2.22	0.40
1:A:343:GLN:HG2	1:C:184:PRO:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	341/349 (98%)	335 (98%)	5 (2%)	1 (0%)	41	31
1	B	341/349 (98%)	334 (98%)	7 (2%)	0	100	100
1	C	343/349 (98%)	332 (97%)	10 (3%)	1 (0%)	41	31
1	D	328/349 (94%)	319 (97%)	9 (3%)	0	100	100
All	All	1353/1396 (97%)	1320 (98%)	31 (2%)	2 (0%)	51	45

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	92	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	92	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/283 (92%)	258 (99%)	3 (1%)	73	73
1	B	255/283 (90%)	249 (98%)	6 (2%)	49	42
1	C	246/283 (87%)	244 (99%)	2 (1%)	81	82
1	D	247/283 (87%)	245 (99%)	2 (1%)	81	82
All	All	1009/1132 (89%)	996 (99%)	13 (1%)	69	67

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	139	ARG
1	A	204	LYS
1	A	344	ASN
1	B	36	VAL
1	B	104	GLN
1	B	192	GLN
1	B	204	LYS
1	B	251	THR
1	B	333	GLN
1	C	283	SER
1	C	341	CYS
1	D	129	LEU
1	D	341	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	340	HIS
1	C	344	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 11 ligands modelled in this entry, 2 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAD	A	401	-	42,48,48	5.06	14 (33%)	50,73,73	1.76	8 (16%)
2	NAD	B	401	-	42,48,48	4.94	16 (38%)	50,73,73	1.75	10 (20%)
6	PO4	C	402	-	4,4,4	0.88	0	6,6,6	1.10	0
5	MPD	D	402	-	7,7,7	1.31	1 (14%)	9,10,10	0.97	1 (11%)
6	PO4	D	401	-	4,4,4	0.81	0	6,6,6	0.63	0
3	13P	B	402	-	9,9,9	1.06	1 (11%)	10,12,12	1.08	1 (10%)
2	NAD	C	401	-	42,48,48	5.43	17 (40%)	50,73,73	1.83	12 (24%)
3	13P	A	402	-	9,9,9	1.32	1 (11%)	10,12,12	0.77	0
5	MPD	A	404	-	7,7,7	0.41	0	9,10,10	1.79	3 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	401	-	-	6/26/62/62	0/5/5/5
2	NAD	B	401	-	-	8/26/62/62	0/5/5/5
5	MPD	D	402	-	-	2/5/5/5	-
3	13P	B	402	-	-	1/7/8/8	-
2	NAD	C	401	-	-	4/26/62/62	0/5/5/5
3	13P	A	402	-	-	1/7/8/8	-
5	MPD	A	404	-	-	1/5/5/5	-

All (50) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	401	NAD	C2B-C1B	-16.66	1.28	1.53
2	A	401	NAD	C2B-C1B	-15.84	1.29	1.53
2	B	401	NAD	C2B-C1B	-15.20	1.30	1.53
2	C	401	NAD	O4D-C1D	14.82	1.61	1.41
2	C	401	NAD	C2D-C1D	-14.73	1.31	1.53
2	B	401	NAD	O4B-C1B	14.67	1.61	1.41
2	C	401	NAD	O4B-C1B	14.55	1.61	1.41
2	A	401	NAD	O4B-C1B	14.31	1.61	1.41
2	A	401	NAD	O4D-C1D	14.18	1.60	1.41
2	B	401	NAD	C2D-C1D	-13.52	1.33	1.53
2	B	401	NAD	O4D-C1D	13.36	1.59	1.41
2	A	401	NAD	C2D-C1D	-12.84	1.34	1.53
2	C	401	NAD	C7N-N7N	8.97	1.50	1.33
2	A	401	NAD	C7N-N7N	8.41	1.49	1.33
2	C	401	NAD	O4B-C4B	-7.89	1.27	1.45
2	B	401	NAD	C7N-N7N	7.77	1.47	1.33
2	B	401	NAD	O4B-C4B	-6.59	1.30	1.45
2	A	401	NAD	O4B-C4B	-6.35	1.30	1.45
2	C	401	NAD	O4D-C4D	-6.19	1.31	1.45
2	A	401	NAD	O4D-C4D	-5.59	1.32	1.45
2	C	401	NAD	C2N-N1N	5.12	1.41	1.35
2	C	401	NAD	O2B-C2B	4.40	1.53	1.43
2	A	401	NAD	C2N-N1N	4.12	1.40	1.35
2	B	401	NAD	O4D-C4D	-4.08	1.35	1.45
2	B	401	NAD	O2B-C2B	3.88	1.52	1.43
2	A	401	NAD	O2B-C2B	3.87	1.52	1.43
2	C	401	NAD	O3D-C3D	-3.80	1.34	1.43
2	A	401	NAD	O2D-C2D	3.45	1.51	1.43
2	B	401	NAD	C6A-N6A	3.41	1.46	1.34
2	C	401	NAD	C6A-N6A	3.41	1.46	1.34
2	A	401	NAD	C6A-N6A	3.18	1.45	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	401	NAD	O2D-C2D	3.01	1.50	1.43
2	A	401	NAD	O3D-C3D	-2.98	1.36	1.43
5	D	402	MPD	O2-C2	-2.97	1.37	1.44
2	A	401	NAD	O3B-C3B	-2.89	1.36	1.43
2	B	401	NAD	O3B-C3B	-2.89	1.36	1.43
2	B	401	NAD	O7N-C7N	-2.85	1.18	1.24
2	B	401	NAD	C2A-N3A	2.77	1.36	1.32
2	C	401	NAD	O3B-C3B	-2.75	1.36	1.43
3	A	402	13P	O1-C1	-2.74	1.41	1.43
2	B	401	NAD	O2D-C2D	2.70	1.49	1.43
2	C	401	NAD	C2N-C3N	2.68	1.43	1.39
3	B	402	13P	O1-C1	-2.40	1.41	1.43
2	C	401	NAD	C5A-C4A	-2.34	1.34	1.40
2	B	401	NAD	C6N-C5N	2.31	1.43	1.38
2	C	401	NAD	C2A-N3A	2.28	1.35	1.32
2	B	401	NAD	C5A-C4A	-2.27	1.34	1.40
2	C	401	NAD	C6N-C5N	2.25	1.43	1.38
2	A	401	NAD	C5A-C4A	-2.22	1.35	1.40
2	B	401	NAD	O3D-C3D	-2.20	1.37	1.43

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	NAD	C5A-C6A-N6A	6.51	130.24	120.35
2	C	401	NAD	C5A-C6A-N6A	6.38	130.05	120.35
2	B	401	NAD	C5A-C6A-N6A	5.92	129.35	120.35
2	C	401	NAD	N3A-C2A-N1A	-5.21	120.54	128.68
2	A	401	NAD	N3A-C2A-N1A	-4.99	120.88	128.68
2	B	401	NAD	N3A-C2A-N1A	-4.58	121.51	128.68
2	C	401	NAD	N6A-C6A-N1A	-4.25	109.75	118.57
2	A	401	NAD	N6A-C6A-N1A	-4.01	110.24	118.57
5	A	404	MPD	CM-C2-C1	-3.92	102.41	110.57
2	C	401	NAD	C6N-N1N-C2N	-3.61	118.68	121.97
2	B	401	NAD	N6A-C6A-N1A	-3.55	111.20	118.57
2	B	401	NAD	C3D-C2D-C1D	3.35	106.02	100.98
2	B	401	NAD	O7N-C7N-N7N	-3.32	117.86	122.58
2	A	401	NAD	C3B-C2B-C1B	3.17	105.75	100.98
2	B	401	NAD	C3B-C2B-C1B	3.15	105.72	100.98
2	C	401	NAD	C3D-C2D-C1D	3.15	105.71	100.98
2	A	401	NAD	C2N-C3N-C4N	2.81	121.45	118.26
2	C	401	NAD	O5B-C5B-C4B	-2.53	100.28	108.99
2	B	401	NAD	C5D-C4D-C3D	-2.50	105.81	115.18

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	NAD	C6N-N1N-C2N	-2.46	119.73	121.97
2	C	401	NAD	PN-O3-PA	-2.38	124.66	132.83
2	A	401	NAD	O4D-C1D-C2D	-2.35	103.49	106.93
2	B	401	NAD	C5B-C4B-C3B	-2.28	106.64	115.18
2	C	401	NAD	O7N-C7N-C3N	2.27	122.35	119.63
2	C	401	NAD	O2D-C2D-C3D	-2.25	104.53	111.82
5	A	404	MPD	C5-C4-C3	-2.24	101.12	111.69
5	D	402	MPD	CM-C2-C1	-2.24	105.92	110.57
2	C	401	NAD	C2D-C3D-C4D	2.21	106.94	102.64
2	B	401	NAD	O4D-C4D-C3D	2.19	109.45	105.11
2	C	401	NAD	C2N-N1N-C1D	2.16	123.94	119.14
2	A	401	NAD	PN-O3-PA	-2.15	125.44	132.83
2	C	401	NAD	C1B-N9A-C4A	-2.13	122.89	126.64
2	B	401	NAD	O7N-C7N-C3N	2.13	122.18	119.63
3	B	402	13P	O2-C2-C3	2.07	123.92	120.77
5	A	404	MPD	O2-C2-CM	2.04	114.61	108.08

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	NAD	O4D-C1D-N1N-C2N
2	A	401	NAD	O4D-C1D-N1N-C6N
2	A	401	NAD	C2D-C1D-N1N-C2N
2	A	401	NAD	C2D-C1D-N1N-C6N
2	B	401	NAD	C5B-O5B-PA-O1A
2	B	401	NAD	O4D-C1D-N1N-C2N
2	B	401	NAD	O4D-C1D-N1N-C6N
2	B	401	NAD	C2D-C1D-N1N-C2N
2	B	401	NAD	C2D-C1D-N1N-C6N
2	C	401	NAD	O4B-C4B-C5B-O5B
2	C	401	NAD	C3B-C4B-C5B-O5B
2	C	401	NAD	O4D-C1D-N1N-C2N
2	C	401	NAD	O4D-C1D-N1N-C6N
3	B	402	13P	O1-C1-C2-O2
3	A	402	13P	O1-C1-C2-O2
2	B	401	NAD	C5B-O5B-PA-O3
2	B	401	NAD	PA-O3-PN-O1N
2	B	401	NAD	C5B-O5B-PA-O2A
5	A	404	MPD	CM-C2-C3-C4
2	A	401	NAD	C4B-C5B-O5B-PA
5	D	402	MPD	C2-C3-C4-C5

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	401	NAD	C5D-O5D-PN-O1N
5	D	402	MPD	C2-C3-C4-O4


There are no ring outliers.


3 monomers are involved in 11 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	NAD	1	0
5	D	402	MPD	7	0
5	A	404	MPD	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	344/349 (98%)	-0.55	0	100 100	9, 21, 38, 49	0
1	B	345/349 (98%)	-0.57	0	100 100	11, 22, 38, 50	0
1	C	344/349 (98%)	-0.54	0	100 100	11, 23, 42, 53	0
1	D	336/349 (96%)	-0.37	2 (0%)	89 91	12, 26, 49, 57	0
All	All	1369/1396 (98%)	-0.51	2 (0%)	95 95	9, 23, 42, 57	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	122	VAL	3.0
1	D	2	ALA	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	MPD	A	404	8/8	0.81	0.20	31,40,51,52	0
6	PO4	C	402	5/5	0.82	0.10	27,31,41,71	0
6	PO4	D	401	5/5	0.87	0.22	48,51,66,79	0
5	MPD	D	402	8/8	0.87	0.20	20,20,20,20	0
2	NAD	C	401	44/44	0.91	0.17	15,30,52,61	70
3	13P	B	402	10/10	0.98	0.10	10,14,20,20	0
2	NAD	B	401	44/44	0.98	0.09	7,16,27,30	0
2	NAD	A	401	44/44	0.98	0.09	9,17,35,41	0
4	CA	A	403	1/1	0.99	0.04	31,31,31,31	0
3	13P	A	402	10/10	0.99	0.09	11,15,21,21	0
4	CA	B	403	1/1	0.99	0.05	25,25,25,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around 13P B 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAD B 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAD A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 13P A 402:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.