



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 18, 2021 – 12:59 AM JST

PDB ID : 7E9W
Title : The Crystal Structure of D-psicose-3-epimerase from Biortus.
Authors : Wang, F.; Xu, C.; Qi, J.; Zhang, M.; Tian, F.; Wang, M.
Deposited on : 2021-03-05
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.18
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.18

i

X-RAY DIFFRACTION

A.

Metric	Percentile Banks	Value
--------	------------------	-------



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

[illegible]

Mol	Chain	Length	Quality of chain
1	A	316	<div> <div></div> <div>85%</div> <div>7%</div> <div>8%</div> </div>
1	B	316	<div> <div></div> <div>86%</div> <div>5%</div> <div>8%</div> </div>
1	C	316	<div> <div></div> <div>87%</div> <div>5%</div> <div>8%</div> </div>
1	D	316	<div> <div></div> <div>84%</div> <div>7%</div> <div>8%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9872 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-psicose 3-epimerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	290	Total	C	N	O	S	0	0	0
			2303	1465	391	436	11			
1	B	290	Total	C	N	O	S	0	1	0
			2308	1468	391	438	11			
1	C	290	Total	C	N	O	S	0	1	0
			2308	1468	392	437	11			
1	D	290	Total	C	N	O	S	0	0	0
			2303	1465	391	436	11			

There are 92 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-22	MET	-	initiating methionine	UNP B8I944
A	-21	ASP	-	expression tag	UNP B8I944
A	-20	TYR	-	expression tag	UNP B8I944
A	-19	LYS	-	expression tag	UNP B8I944
A	-18	ASP	-	expression tag	UNP B8I944
A	-17	ASP	-	expression tag	UNP B8I944
A	-16	ASP	-	expression tag	UNP B8I944
A	-15	ASP	-	expression tag	UNP B8I944
A	-14	LYS	-	expression tag	UNP B8I944
A	-13	HIS	-	expression tag	UNP B8I944
A	-12	HIS	-	expression tag	UNP B8I944
A	-11	HIS	-	expression tag	UNP B8I944
A	-10	HIS	-	expression tag	UNP B8I944
A	-9	HIS	-	expression tag	UNP B8I944
A	-8	HIS	-	expression tag	UNP B8I944
A	-7	GLU	-	expression tag	UNP B8I944
A	-6	ASN	-	expression tag	UNP B8I944
A	-5	LEU	-	expression tag	UNP B8I944
A	-4	TYR	-	expression tag	UNP B8I944
A	-3	PHE	-	expression tag	UNP B8I944
A	-2	GLN	-	expression tag	UNP B8I944

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP B8I944
A	0	GLY	-	expression tag	UNP B8I944
B	-22	MET	-	initiating methionine	UNP B8I944
B	-21	ASP	-	expression tag	UNP B8I944
B	-20	TYR	-	expression tag	UNP B8I944
B	-19	LYS	-	expression tag	UNP B8I944
B	-18	ASP	-	expression tag	UNP B8I944
B	-17	ASP	-	expression tag	UNP B8I944
B	-16	ASP	-	expression tag	UNP B8I944
B	-15	ASP	-	expression tag	UNP B8I944
B	-14	LYS	-	expression tag	UNP B8I944
B	-13	HIS	-	expression tag	UNP B8I944
B	-12	HIS	-	expression tag	UNP B8I944
B	-11	HIS	-	expression tag	UNP B8I944
B	-10	HIS	-	expression tag	UNP B8I944
B	-9	HIS	-	expression tag	UNP B8I944
B	-8	HIS	-	expression tag	UNP B8I944
B	-7	GLU	-	expression tag	UNP B8I944
B	-6	ASN	-	expression tag	UNP B8I944
B	-5	LEU	-	expression tag	UNP B8I944
B	-4	TYR	-	expression tag	UNP B8I944
B	-3	PHE	-	expression tag	UNP B8I944
B	-2	GLN	-	expression tag	UNP B8I944
B	-1	GLY	-	expression tag	UNP B8I944
B	0	GLY	-	expression tag	UNP B8I944
C	-22	MET	-	initiating methionine	UNP B8I944
C	-21	ASP	-	expression tag	UNP B8I944
C	-20	TYR	-	expression tag	UNP B8I944
C	-19	LYS	-	expression tag	UNP B8I944
C	-18	ASP	-	expression tag	UNP B8I944
C	-17	ASP	-	expression tag	UNP B8I944
C	-16	ASP	-	expression tag	UNP B8I944
C	-15	ASP	-	expression tag	UNP B8I944
C	-14	LYS	-	expression tag	UNP B8I944
C	-13	HIS	-	expression tag	UNP B8I944
C	-12	HIS	-	expression tag	UNP B8I944
C	-11	HIS	-	expression tag	UNP B8I944
C	-10	HIS	-	expression tag	UNP B8I944
C	-9	HIS	-	expression tag	UNP B8I944
C	-8	HIS	-	expression tag	UNP B8I944
C	-7	GLU	-	expression tag	UNP B8I944
C	-6	ASN	-	expression tag	UNP B8I944

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-5	LEU	-	expression tag	UNP B8I944
C	-4	TYR	-	expression tag	UNP B8I944
C	-3	PHE	-	expression tag	UNP B8I944
C	-2	GLN	-	expression tag	UNP B8I944
C	-1	GLY	-	expression tag	UNP B8I944
C	0	GLY	-	expression tag	UNP B8I944
D	-22	MET	-	initiating methionine	UNP B8I944
D	-21	ASP	-	expression tag	UNP B8I944
D	-20	TYR	-	expression tag	UNP B8I944
D	-19	LYS	-	expression tag	UNP B8I944
D	-18	ASP	-	expression tag	UNP B8I944
D	-17	ASP	-	expression tag	UNP B8I944
D	-16	ASP	-	expression tag	UNP B8I944
D	-15	ASP	-	expression tag	UNP B8I944
D	-14	LYS	-	expression tag	UNP B8I944
D	-13	HIS	-	expression tag	UNP B8I944
D	-12	HIS	-	expression tag	UNP B8I944
D	-11	HIS	-	expression tag	UNP B8I944
D	-10	HIS	-	expression tag	UNP B8I944
D	-9	HIS	-	expression tag	UNP B8I944
D	-8	HIS	-	expression tag	UNP B8I944
D	-7	GLU	-	expression tag	UNP B8I944
D	-6	ASN	-	expression tag	UNP B8I944
D	-5	LEU	-	expression tag	UNP B8I944
D	-4	TYR	-	expression tag	UNP B8I944
D	-3	PHE	-	expression tag	UNP B8I944
D	-2	GLN	-	expression tag	UNP B8I944
D	-1	GLY	-	expression tag	UNP B8I944
D	0	GLY	-	expression tag	UNP B8I944

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mn 1 1	0	0
2	B	1	Total Mn 1 1	0	0
2	C	1	Total Mn 1 1	0	0
2	D	1	Total Mn 1 1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	186	Total 186	O 186	0	0
4	B	123	Total 123	O 123	0	0
4	C	149	Total 149	O 149	0	0
4	D	116	Total 116	O 116	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: D-psicose 3-epimerase




- Molecule 1: D-psicose 3-epimerase




- Molecule 1: D-psicose 3-epimerase



- Molecule 1: D-psicose 3-epimerase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	79.13Å 116.08Å 91.82Å 90.00° 105.08° 90.00°	Depositor
Resolution (Å)	48.61 – 2.10 48.56 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.0 (48.61-2.10) 98.0 (48.56-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.94 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.174 , 0.206 0.181 , 0.211	Depositor DCC
R_{free} test set	4559 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	32.8	Xtriage
Anisotropy	0.033	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 37.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9872	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	0/2357	0.71	0/3192
1	B	0.64	0/2365	0.71	0/3203
1	C	0.64	0/2365	0.72	0/3203
1	D	0.63	0/2357	0.71	0/3192
All	All	0.64	0/9444	0.71	0/12790

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2303	0	2235	14	0
1	B	2308	0	2239	11	0
1	C	2308	0	2241	12	0
1	D	2303	0	2235	14	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	18	0	23	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	18	0	23	2	0
3	C	18	0	23	3	0
3	D	18	0	23	0	0
4	A	186	0	0	2	0
4	B	123	0	0	1	0
4	C	149	0	0	2	0
4	D	116	0	0	1	0
All	All	9872	0	9042	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 53 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:25:VAL:HG13	1:B:30:PHE:HB2	1.60	0.82
1:B:186:HIS:HE2	3:B:304:GOL:H31	1.63	0.63
1:A:284:ARG:O	1:A:288:GLU:HB2	2.01	0.61
1:B:181:MET:HE3	1:B:209:HIS:CE1	2.41	0.56
1:B:181:MET:CE	1:B:209:HIS:CE1	2.89	0.56

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	288/316 (91%)	281 (98%)	7 (2%)	0	100	100
1	B	289/316 (92%)	282 (98%)	7 (2%)	0	100	100
1	C	289/316 (92%)	282 (98%)	6 (2%)	1 (0%)	41	41
1	D	288/316 (91%)	278 (96%)	10 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	1154/1264 (91%)	1123 (97%)	30 (3%)	1 (0%)	51 54

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	289	CYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	242/266 (91%)	241 (100%)	1 (0%)	91 94
1	B	243/266 (91%)	239 (98%)	4 (2%)	62 69
1	C	243/266 (91%)	242 (100%)	1 (0%)	91 94
1	D	242/266 (91%)	239 (99%)	3 (1%)	71 77
All	All	970/1064 (91%)	961 (99%)	9 (1%)	81 84

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	73	ASN
1	D	116	TYR
1	B	73	ASN
1	B	116	TYR
1	C	116	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	84	ASN
1	C	238	ASN
1	D	73	ASN
1	B	73	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	277	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 4 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	B	304	2	5,5,5	0.17	0	5,5,5	0.45	0
3	GOL	B	302	-	5,5,5	0.09	0	5,5,5	0.26	0
3	GOL	D	303	-	5,5,5	0.11	0	5,5,5	0.34	0
3	GOL	B	303	-	5,5,5	0.11	0	5,5,5	0.32	0
3	GOL	A	304	2	5,5,5	0.17	0	5,5,5	0.54	0
3	GOL	D	302	-	5,5,5	0.12	0	5,5,5	0.36	0
3	GOL	D	304	2	5,5,5	0.18	0	5,5,5	0.35	0
3	GOL	C	304	2	5,5,5	0.16	0	5,5,5	0.34	0
3	GOL	A	303	-	5,5,5	0.09	0	5,5,5	0.32	0
3	GOL	A	302	-	5,5,5	0.10	0	5,5,5	0.33	0
3	GOL	C	302	-	5,5,5	0.13	0	5,5,5	0.42	0
3	GOL	C	303	-	5,5,5	0.11	0	5,5,5	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	304	2	-	4/4/4/4	-
3	GOL	B	302	-	-	2/4/4/4	-
3	GOL	D	303	-	-	2/4/4/4	-
3	GOL	B	303	-	-	4/4/4/4	-
3	GOL	A	304	2	-	0/4/4/4	-
3	GOL	D	302	-	-	2/4/4/4	-
3	GOL	D	304	2	-	4/4/4/4	-
3	GOL	C	304	2	-	4/4/4/4	-
3	GOL	A	303	-	-	1/4/4/4	-
3	GOL	A	302	-	-	2/4/4/4	-
3	GOL	C	302	-	-	2/4/4/4	-
3	GOL	C	303	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 29 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	302	GOL	C1-C2-C3-O3
3	B	302	GOL	O2-C2-C3-O3
3	B	303	GOL	C1-C2-C3-O3
3	B	304	GOL	O1-C1-C2-C3
3	C	302	GOL	O1-C1-C2-C3

There are no ring outliers.

6 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	304	GOL	1	0
3	B	302	GOL	1	0
3	A	304	GOL	1	0
3	C	304	GOL	2	0
3	A	303	GOL	2	0
3	C	303	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	290/316 (91%)	-0.28	2 (0%) 87 89	23, 30, 44, 79	0
1	B	290/316 (91%)	0.02	4 (1%) 75 78	24, 37, 55, 80	0
1	C	290/316 (91%)	-0.16	5 (1%) 70 74	24, 34, 49, 89	0
1	D	290/316 (91%)	0.19	8 (2%) 53 59	25, 38, 58, 77	0
All	All	1160/1264 (91%)	-0.06	19 (1%) 72 75	23, 34, 55, 89	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	289	CYS	4.5
1	C	238	ASN	4.3
1	A	289	CYS	3.8
1	B	290	HIS	3.6
1	C	289	CYS	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MN	A	301	1/1	0.50	0.19	49,49,49,49	1
3	GOL	B	304	6/6	0.54	0.31	53,61,67,70	0
3	GOL	C	304	6/6	0.63	0.25	51,56,63,65	0
3	GOL	A	304	6/6	0.64	0.25	57,60,65,65	0
2	MN	C	301	1/1	0.75	0.13	48,48,48,48	1
3	GOL	D	304	6/6	0.77	0.28	55,61,64,66	0
3	GOL	B	303	6/6	0.84	0.20	55,59,60,62	0
2	MN	D	301	1/1	0.84	0.10	47,47,47,47	1
2	MN	B	301	1/1	0.90	0.21	49,49,49,49	1
3	GOL	B	302	6/6	0.90	0.14	43,50,52,53	0
3	GOL	C	302	6/6	0.91	0.14	42,47,51,51	0
3	GOL	A	302	6/6	0.92	0.15	31,37,43,45	0
3	GOL	D	302	6/6	0.92	0.16	38,48,51,53	0
3	GOL	C	303	6/6	0.92	0.15	49,50,51,51	0
3	GOL	D	303	6/6	0.93	0.22	50,52,53,54	0
3	GOL	A	303	6/6	0.95	0.10	44,47,48,51	0

6.5 Other polymers [i](#)

There are no such residues in this entry.