



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 6, 2022 – 04:24 PM EDT

PDB ID : 6EA4
Title : ERAP2 bound to Aryl Sulfonamide Uncompetitive Inhibitor
Authors : Maben, Z.; Stern, L.J.
Deposited on : 2018-08-02
Resolution : 2.45 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.28.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.28.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1544 (2.48-2.44)
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)
RSRZ outliers	127900	1523 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	911	<div> <div>2%</div> <div>87%</div> <div>9%</div> <div>.</div> </div>
1	B	911	<div> <div>12%</div> <div>83%</div> <div>12%</div> <div>5%</div> </div>
2	C	5	<div> <div>60%</div> <div>40%</div> </div>
2	D	5	<div> <div>60%</div> <div>40%</div> </div>
2	G	5	<div> <div>60%</div> <div>40%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	2	 100%
3	F	2	 100%
3	I	2	 100%
4	H	3	 67% 33%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	IMD	B	1008	-	-	-	X
2	BMA	D	3	-	-	-	X
2	MAN	D	4	-	-	-	X
2	MAN	G	4	-	-	-	X
3	NAG	F	2	-	-	-	X
3	NAG	I	2	-	-	-	X
4	BMA	H	3	-	-	-	X
7	J2G	A	1003[A]	-	-	-	X
7	J2G	A	1003[B]	-	-	-	X
7	J2G	B	1003[A]	-	-	-	X
7	J2G	B	1003[B]	-	-	-	X
8	LYS	B	1010	-	-	-	X

2 Entry composition [i](#)

There are 13 unique types of molecules in this entry. The entry contains 15097 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Endoplasmic reticulum aminopeptidase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	883	Total	C	N	O	S	0	7	0
			7212	4653	1201	1328	30			
1	B	869	Total	C	N	O	S	0	2	0
			7054	4550	1169	1306	29			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	392	ASN	LYS	variant	UNP Q6P179
A	961	ARG	-	expression tag	UNP Q6P179
A	962	HIS	-	expression tag	UNP Q6P179
A	963	HIS	-	expression tag	UNP Q6P179
B	392	ASN	LYS	variant	UNP Q6P179
B	961	ARG	-	expression tag	UNP Q6P179
B	962	HIS	-	expression tag	UNP Q6P179
B	963	HIS	-	expression tag	UNP Q6P179

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	5	Total	C	N	O	0	0	0
			61	34	2	25			
2	D	5	Total	C	N	O	0	0	0
			61	34	2	25			
2	G	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	F	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	I	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	H	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Zn	0	0
			1	1		
5	B	1	Total	Zn	0	0
			1	1		

- Molecule 6 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total 12	C 6	N 1	O 4	S 1	0	0
6	B	1	Total 12	C 6	N 1	O 4	S 1	0	0

- Molecule 7 is 4-methoxy-3-{[2-(piperidin-1-yl)-4-(trifluoromethyl)phenyl]sulfamoyl}benzoic acid (three-letter code: J2G) (formula: C₂₀H₂₁F₃N₂O₅S) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
7	A	1	Total	C	F	N	O	S	0	1
			62	40	6	4	10	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
7	B	1	Total	C	F	N	O	S	0	1
			62	40	6	4	10	2		

- Molecule 8 is LYSINE (three-letter code: LYS) (formula: $C_6H_{15}N_2O_2$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	N	O	0	0
			10	6	2	2		
8	A	1	Total	C	N	O	0	0
			10	6	2	2		
8	B	1	Total	C	N	O	0	0
			10	6	2	2		
8	B	1	Total	C	N	O	0	0
			10	6	2	2		

- Molecule 9 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	A	1	Total	C	N	O	0	0
			14	8	1	5		
9	A	1	Total	C	N	O	0	0
			14	8	1	5		
9	A	1	Total	C	N	O	0	0
			14	8	1	5		
9	A	1	Total	C	N	O	0	0
			14	8	1	5		
9	B	1	Total	C	N	O	0	0
			14	8	1	5		
9	B	1	Total	C	N	O	0	0
			14	8	1	5		
9	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 10 is IMIDAZOLE (three-letter code: IMD) (formula: C₃H₅N₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	C	N	0	0
			5	3	2		
10	B	1	Total	C	N	0	0
			5	3	2		

- Molecule 11 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	A	1	Total	C	O	0	0
			10	6	4		


- Molecule 12 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
12	A	1	Total	C	O	0	0
			4	2	2		
12	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 13 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	A	175	Total	O	0	0
			175	175		
13	B	34	Total	O	0	0
			34	34		



- Molecule 2: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C: 60% 40%



- Molecule 2: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D: 60% 40%



- Molecule 2: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G: 60% 40%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E: 100%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F: 100%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  100%

MAG1
MAG2

- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  67% 33%

MAG1
MAG2
BNA3

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	75.23Å 135.45Å 127.28Å 90.00° 90.11° 90.00°	Depositor
Resolution (Å)	29.29 – 2.45 29.89 – 2.45	Depositor EDS
% Data completeness (in resolution range)	97.1 (29.29-2.45) 87.2 (29.89-2.45)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.17 (at 2.45Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.208 , 0.236 0.208 , 0.233	Depositor DCC
R_{free} test set	2027 reflections (2.23%)	wwPDB-VP
Wilson B-factor (Å ²)	48.0	Xtriage
Anisotropy	0.386	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 43.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.026 for h,-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	15097	wwPDB-VP
Average B, all atoms (Å ²)	82.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, PGE, ZN, BMA, MES, IMD, NAG, EDO, J2G

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/7408	0.44	0/10041
1	B	0.23	0/7233	0.45	0/9808
All	All	0.23	0/14641	0.45	0/19849

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7212	0	7151	51	0
1	B	7054	0	6957	68	0
2	C	61	0	52	0	0
2	D	61	0	52	0	0
2	G	61	0	52	0	0
3	E	28	0	25	0	0
3	F	28	0	25	0	0
3	I	28	0	25	0	0
4	H	39	0	34	1	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	12	0	12	3	0
6	B	12	0	12	2	0
7	A	62	0	0	1	0
7	B	62	0	0	0	0
8	A	20	0	24	2	0
8	B	20	0	24	2	0
9	A	56	0	52	0	0
9	B	42	0	39	0	0
10	A	5	0	5	0	0
10	B	5	0	5	2	0
11	A	10	0	14	2	0
12	A	4	0	6	0	0
12	B	4	0	6	0	0
13	A	175	0	0	3	0
13	B	34	0	0	0	0
All	All	15097	0	14572	123	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 123 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:777:TRP:HE1	1:A:783:LYS:HD2	1.42	0.83
1:A:383:MET:HE2	1:A:389:ILE:HA	1.66	0.77
1:A:177[B]:GLU:HB3	1:A:203:GLN:HG2	1.67	0.76
1:B:177[B]:GLU:HB3	1:B:203:GLN:HG2	1.72	0.72
1:B:200:GLU:OE1	8:B:1004:LYS:N	2.23	0.71

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	886/911 (97%)	859 (97%)	27 (3%)	0	100	100
1	B	863/911 (95%)	835 (97%)	28 (3%)	0	100	100
All	All	1749/1822 (96%)	1694 (97%)	55 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	792/819 (97%)	786 (99%)	6 (1%)	81	88
1	B	772/819 (94%)	764 (99%)	8 (1%)	76	84
All	All	1564/1638 (96%)	1550 (99%)	14 (1%)	78	86

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	322	LYS
1	B	339	TRP
1	B	779	GLU
1	B	686	TYR
1	B	755	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	869	GLN
1	B	581	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

24 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	2,1	14,14,15	0.28	0	17,19,21	0.37	0
2	NAG	C	2	2	14,14,15	0.26	0	17,19,21	0.44	0
2	BMA	C	3	2	11,11,12	0.54	0	15,15,17	0.83	0
2	MAN	C	4	2	11,11,12	0.76	1 (9%)	15,15,17	1.61	3 (20%)
2	MAN	C	5	2	11,11,12	0.57	0	15,15,17	1.03	2 (13%)
2	NAG	D	1	2,1	14,14,15	0.27	0	17,19,21	0.38	0
2	NAG	D	2	2	14,14,15	0.21	0	17,19,21	0.48	0
2	BMA	D	3	2	11,11,12	0.69	0	15,15,17	0.78	0
2	MAN	D	4	2	11,11,12	0.88	1 (9%)	15,15,17	1.63	3 (20%)
2	MAN	D	5	2	11,11,12	0.66	0	15,15,17	1.03	2 (13%)
3	NAG	E	1	3,1	14,14,15	0.23	0	17,19,21	0.41	0
3	NAG	E	2	3	14,14,15	0.23	0	17,19,21	0.40	0
3	NAG	F	1	3,1	14,14,15	0.20	0	17,19,21	0.38	0
3	NAG	F	2	3	14,14,15	0.23	0	17,19,21	0.41	0
2	NAG	G	1	2,1	14,14,15	0.27	0	17,19,21	0.37	0
2	NAG	G	2	2	14,14,15	0.31	0	17,19,21	0.44	0
2	BMA	G	3	2	11,11,12	0.60	0	15,15,17	0.73	0
2	MAN	G	4	2	11,11,12	0.92	1 (9%)	15,15,17	1.64	3 (20%)
2	MAN	G	5	2	11,11,12	0.67	0	15,15,17	1.04	2 (13%)
4	NAG	H	1	1,4	14,14,15	0.17	0	17,19,21	0.40	0
4	NAG	H	2	4	14,14,15	0.24	0	17,19,21	0.44	0
4	BMA	H	3	4	11,11,12	0.63	0	15,15,17	0.70	0
3	NAG	I	1	3,1	14,14,15	0.25	0	17,19,21	0.44	0
3	NAG	I	2	3	14,14,15	0.24	0	17,19,21	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	MAN	C	4	2	-	2/2/19/22	0/1/1/1
2	MAN	C	5	2	-	0/2/19/22	0/1/1/1
2	NAG	D	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1
2	BMA	D	3	2	-	0/2/19/22	0/1/1/1
2	MAN	D	4	2	-	0/2/19/22	0/1/1/1
2	MAN	D	5	2	-	1/2/19/22	0/1/1/1
3	NAG	E	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	E	2	3	-	2/6/23/26	0/1/1/1
3	NAG	F	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	F	2	3	-	2/6/23/26	0/1/1/1
2	NAG	G	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	G	2	2	-	0/6/23/26	0/1/1/1
2	BMA	G	3	2	-	0/2/19/22	0/1/1/1
2	MAN	G	4	2	-	2/2/19/22	0/1/1/1
2	MAN	G	5	2	-	2/2/19/22	0/1/1/1
4	NAG	H	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	H	2	4	-	0/6/23/26	0/1/1/1
4	BMA	H	3	4	-	0/2/19/22	0/1/1/1
3	NAG	I	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	I	2	3	-	0/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	4	MAN	C1-C2	2.43	1.57	1.52
2	G	4	MAN	C1-C2	2.43	1.57	1.52
2	C	4	MAN	C1-C2	2.09	1.57	1.52

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	4	MAN	C1-O5-C5	5.08	119.07	112.19
2	D	4	MAN	C1-O5-C5	4.93	118.88	112.19
2	C	4	MAN	C1-O5-C5	4.79	118.68	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	5	MAN	C1-O5-C5	2.77	115.95	112.19
2	D	5	MAN	C1-O5-C5	2.68	115.82	112.19

There are no chirality outliers.

5 of 13 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	G	4	MAN	C4-C5-C6-O6
2	G	4	MAN	O5-C5-C6-O6
3	F	2	NAG	O5-C5-C6-O6
2	G	5	MAN	C4-C5-C6-O6
2	C	4	MAN	O5-C5-C6-O6


There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	H	1	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

















5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 2 are monoatomic - leaving 22 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
9	NAG	A	1007	1	14,14,15	0.30	0	17,19,21	0.45	0
7	J2G	A	1003[A]	-	31,33,33	1.28	3 (9%)	45,49,49	1.22	6 (13%)
12	EDO	A	1012	-	3,3,3	0.46	0	2,2,2	0.34	0
10	IMD	A	1010	-	3,5,5	0.41	0	4,5,5	0.58	0
8	LYS	A	1004	5	5,9,9	0.28	0	4,10,10	0.36	0
9	NAG	B	1007	1	14,14,15	0.24	0	17,19,21	0.46	0
11	PGE	A	1011	-	9,9,9	0.30	0	8,8,8	0.29	0
9	NAG	A	1008	1	14,14,15	0.31	0	17,19,21	0.45	0
6	MES	A	1002	-	12,12,12	2.32	1 (8%)	14,16,16	1.83	5 (35%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
9	NAG	B	1005	1	14,14,15	0.21	0	17,19,21	0.40	0
8	LYS	A	1009	-	5,9,9	0.33	0	4,10,10	0.47	0
7	J2G	B	1003[B]	-	31,33,33	1.26	2 (6%)	45,49,49	1.23	4 (8%)
7	J2G	A	1003[B]	-	31,33,33	1.28	2 (6%)	45,49,49	1.22	3 (6%)
8	LYS	B	1010	-	5,9,9	0.32	0	4,10,10	0.39	0
12	EDO	B	1009	-	3,3,3	0.46	0	2,2,2	0.32	0
9	NAG	A	1005	1	14,14,15	0.23	0	17,19,21	0.42	0
8	LYS	B	1004	5	5,9,9	0.28	0	4,10,10	0.43	0
6	MES	B	1002	-	12,12,12	2.26	1 (8%)	14,16,16	1.98	5 (35%)
10	IMD	B	1008	-	3,5,5	0.42	0	4,5,5	0.58	0
9	NAG	B	1006	1	14,14,15	0.23	0	17,19,21	0.42	0
9	NAG	A	1006	1	14,14,15	0.23	0	17,19,21	0.44	0
7	J2G	B	1003[A]	-	31,33,33	1.26	3 (9%)	45,49,49	1.20	6 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	NAG	A	1007	1	-	1/6/23/26	0/1/1/1
7	J2G	A	1003[A]	-	-	14/23/35/35	0/3/3/3
12	EDO	A	1012	-	-	0/1/1/1	-
10	IMD	A	1010	-	-	-	0/1/1/1
8	LYS	A	1004	5	-	0/5/9/9	-
9	NAG	B	1007	1	-	4/6/23/26	0/1/1/1
11	PGE	A	1011	-	-	3/7/7/7	-
9	NAG	A	1008	1	-	0/6/23/26	0/1/1/1
6	MES	A	1002	-	-	3/6/14/14	0/1/1/1
9	NAG	B	1005	1	-	2/6/23/26	0/1/1/1
8	LYS	A	1009	-	-	0/5/9/9	-
7	J2G	B	1003[B]	-	-	9/23/35/35	0/3/3/3
7	J2G	A	1003[B]	-	-	9/23/35/35	0/3/3/3
8	LYS	B	1010	-	-	0/5/9/9	-
12	EDO	B	1009	-	-	0/1/1/1	-
9	NAG	A	1005	1	-	2/6/23/26	0/1/1/1
8	LYS	B	1004	5	-	0/5/9/9	-
6	MES	B	1002	-	-	0/6/14/14	0/1/1/1
10	IMD	B	1008	-	-	-	0/1/1/1
9	NAG	B	1006	1	-	1/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	NAG	A	1006	1	-	2/6/23/26	0/1/1/1
7	J2G	B	1003[A]	-	-	14/23/35/35	0/3/3/3

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	1002	MES	C8-S	-7.75	1.66	1.77
6	B	1002	MES	C8-S	-7.55	1.66	1.77
7	A	1003[B]	J2G	CAN-CAV	5.30	1.52	1.47
7	A	1003[A]	J2G	CAN-CAV	5.25	1.52	1.47
7	B	1003[B]	J2G	CAN-CAV	5.18	1.52	1.47

The worst 5 of 29 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	1002	MES	C5-N4-C3	3.92	117.66	108.83
7	A	1003[B]	J2G	CAF-SAG-NAH	-3.86	102.76	107.27
7	B	1003[B]	J2G	CBC-NAZ-CBA	3.72	119.73	111.52
7	B	1003[B]	J2G	OAC-SAG-CAF	3.60	113.58	107.66
7	A	1003[B]	J2G	OAC-SAG-CAF	3.49	113.40	107.66

There are no chirality outliers.

5 of 64 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	1002	MES	C7-C8-S-O2S
6	A	1002	MES	C7-C8-S-O3S
7	A	1003[A]	J2G	CAE-CAF-SAG-NAH
7	A	1003[A]	J2G	CAE-CAF-SAG-OAC
7	A	1003[A]	J2G	CAE-CAF-SAG-OAP

There are no ring outliers.

8 monomers are involved in 13 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	1004	LYS	1	0
11	A	1011	PGE	2	0
6	A	1002	MES	3	0
8	A	1009	LYS	1	0
7	A	1003[B]	J2G	1	0



Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	B	1004	LYS	2	0
6	B	1002	MES	2	0
10	B	1008	IMD	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







Ligand J2G B 1003 (B)





Ligand J2G A 1003 (B)



Ligand LYS B 1010





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	883/911 (96%)	-0.10	22 (2%) 57 53	38, 59, 95, 142	0
1	B	869/911 (95%)	0.63	107 (12%) 4 2	46, 98, 158, 198	0
All	All	1752/1822 (96%)	0.27	129 (7%) 14 11	38, 73, 144, 198	0

The worst 5 of 129 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	556	GLY	6.1
1	B	570	VAL	5.9
1	B	785	ASN	5.9
1	B	105	THR	5.8
1	B	501	ASN	5.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MAN	D	4	11/12	0.57	0.41	141,143,147,148	0
3	NAG	I	2	14/15	0.66	0.50	135,142,143,143	0
2	MAN	D	5	11/12	0.67	0.37	91,103,112,112	0
3	NAG	F	2	14/15	0.69	0.41	122,136,140,141	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	BMA	H	3	11/12	0.70	0.56	136,139,143,143	0
2	BMA	G	3	11/12	0.71	0.29	104,109,118,123	0
2	BMA	C	3	11/12	0.71	0.36	114,117,123,128	0
2	BMA	D	3	11/12	0.75	0.46	114,120,135,140	0
3	NAG	I	1	14/15	0.76	0.23	112,125,134,139	0
2	MAN	G	4	11/12	0.77	0.47	129,132,135,136	0
2	MAN	G	5	11/12	0.79	0.35	72,96,106,108	0
2	MAN	C	4	11/12	0.81	0.49	124,128,132,134	0
2	MAN	C	5	11/12	0.81	0.41	98,112,119,121	0
3	NAG	E	2	14/15	0.83	0.38	106,117,121,121	0
3	NAG	E	1	14/15	0.84	0.20	71,87,102,111	0
3	NAG	F	1	14/15	0.84	0.29	96,105,123,130	0
4	NAG	H	1	14/15	0.86	0.32	89,100,111,114	0
2	NAG	D	2	14/15	0.86	0.24	65,76,89,108	0
4	NAG	H	2	14/15	0.88	0.39	113,121,126,132	0
2	NAG	C	2	14/15	0.90	0.20	68,86,99,102	0
2	NAG	G	2	14/15	0.92	0.26	63,80,93,100	0
2	NAG	D	1	14/15	0.94	0.13	40,57,62,62	0
2	NAG	G	1	14/15	0.95	0.17	52,63,73,83	0
2	NAG	C	1	14/15	0.95	0.09	53,64,76,78	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain F:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain I:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
10	IMD	B	1008	5/5	0.54	0.59	95,97,103,111	5
9	NAG	A	1008	14/15	0.62	0.35	107,127,134,144	0
8	LYS	B	1010	10/10	0.63	0.41	83,90,93,93	10
7	J2G	B	1003[A]	31/31	0.63	0.50	81,86,89,89	31
7	J2G	B	1003[B]	31/31	0.63	0.50	81,85,88,90	31
7	J2G	A	1003[B]	31/31	0.69	0.43	50,58,62,63	31
7	J2G	A	1003[A]	31/31	0.69	0.43	50,59,63,64	31
9	NAG	B	1005	14/15	0.71	0.35	143,153,157,157	0
9	NAG	A	1007	14/15	0.72	0.34	118,121,127,131	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
9	NAG	B	1007	14/15	0.80	0.52	188,196,199,200	0
9	NAG	B	1006	14/15	0.81	0.30	169,174,181,182	0
11	PGE	A	1011	10/10	0.81	0.26	55,66,72,82	10
9	NAG	A	1005	14/15	0.82	0.31	119,128,137,141	0
10	IMD	A	1010	5/5	0.83	0.33	54,59,62,64	5
9	NAG	A	1006	14/15	0.83	0.34	89,106,117,118	0
8	LYS	A	1009	10/10	0.83	0.30	63,68,71,72	10
12	EDO	B	1009	4/4	0.83	0.23	98,103,108,112	0
6	MES	B	1002	12/12	0.86	0.23	82,87,88,100	12
8	LYS	B	1004	10/10	0.89	0.27	82,85,88,90	0
6	MES	A	1002	12/12	0.92	0.21	51,58,71,81	12
12	EDO	A	1012	4/4	0.94	0.22	64,67,68,77	0
8	LYS	A	1004	10/10	0.96	0.25	49,51,61,65	0
5	ZN	B	1001	1/1	0.98	0.16	73,73,73,73	0
5	ZN	A	1001	1/1	1.00	0.16	41,41,41,41	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around J2G B 1003 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around J2G B 1003 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around J2G A 1003 (B):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around J2G A 1003 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around LYS A 1009:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around LYS B 1004:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)


**Electron density around LYS A 1004:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around ZN B 1001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZN A 1001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.