



Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 05:12 am BST

PDB ID : 6EBP
Title : Crystal Structure of the Class Ie Ribonucleotide Reductase Beta Subunit from *Aerococcus urinae* in Activated Form
Authors : Palowitch, G.M.; Boal, A.K.
Deposited on : 2018-08-06
Resolution : 1.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	355	<div> <div>3%</div> <div>85%</div> <div>15%</div> </div>
1	B	355	<div> <div>4%</div> <div>86%</div> <div>12%</div> </div>
1	C	355	<div> <div>3%</div> <div>85%</div> <div>14%</div> </div>
1	D	355	<div> <div>4%</div> <div>86%</div> <div>12%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 10705 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribonucleoside-diphosphate reductase, beta subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	303	Total	C	N	O	S	0	0	0
			2468	1596	400	466	6			
1	B	311	Total	C	N	O	S	0	0	0
			2538	1637	414	481	6			
1	C	304	Total	C	N	O	S	0	0	0
			2479	1602	404	467	6			
1	D	311	Total	C	N	O	S	0	0	0
			2538	1637	414	481	6			

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-17	MET	-	initiating methionine	UNP F2I8X9
A	-16	GLY	-	expression tag	UNP F2I8X9
A	-15	SER	-	expression tag	UNP F2I8X9
A	-14	SER	-	expression tag	UNP F2I8X9
A	-13	HIS	-	expression tag	UNP F2I8X9
A	-12	HIS	-	expression tag	UNP F2I8X9
A	-11	HIS	-	expression tag	UNP F2I8X9
A	-10	HIS	-	expression tag	UNP F2I8X9
A	-9	HIS	-	expression tag	UNP F2I8X9
A	-8	HIS	-	expression tag	UNP F2I8X9
A	-7	SER	-	expression tag	UNP F2I8X9
A	-6	SER	-	expression tag	UNP F2I8X9
A	-5	GLY	-	expression tag	UNP F2I8X9
A	-4	LEU	-	expression tag	UNP F2I8X9
A	-3	VAL	-	expression tag	UNP F2I8X9
A	-2	PRO	-	expression tag	UNP F2I8X9
A	-1	ARG	-	expression tag	UNP F2I8X9
A	0	GLY	-	expression tag	UNP F2I8X9
A	1	SER	-	expression tag	UNP F2I8X9
A	123	DAH	TYR	see remark 999	UNP F2I8X9
B	-17	MET	-	initiating methionine	UNP F2I8X9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	GLY	-	expression tag	UNP F2I8X9
B	-15	SER	-	expression tag	UNP F2I8X9
B	-14	SER	-	expression tag	UNP F2I8X9
B	-13	HIS	-	expression tag	UNP F2I8X9
B	-12	HIS	-	expression tag	UNP F2I8X9
B	-11	HIS	-	expression tag	UNP F2I8X9
B	-10	HIS	-	expression tag	UNP F2I8X9
B	-9	HIS	-	expression tag	UNP F2I8X9
B	-8	HIS	-	expression tag	UNP F2I8X9
B	-7	SER	-	expression tag	UNP F2I8X9
B	-6	SER	-	expression tag	UNP F2I8X9
B	-5	GLY	-	expression tag	UNP F2I8X9
B	-4	LEU	-	expression tag	UNP F2I8X9
B	-3	VAL	-	expression tag	UNP F2I8X9
B	-2	PRO	-	expression tag	UNP F2I8X9
B	-1	ARG	-	expression tag	UNP F2I8X9
B	0	GLY	-	expression tag	UNP F2I8X9
B	1	SER	-	expression tag	UNP F2I8X9
B	123	DAH	TYR	see remark 999	UNP F2I8X9
C	-17	MET	-	initiating methionine	UNP F2I8X9
C	-16	GLY	-	expression tag	UNP F2I8X9
C	-15	SER	-	expression tag	UNP F2I8X9
C	-14	SER	-	expression tag	UNP F2I8X9
C	-13	HIS	-	expression tag	UNP F2I8X9
C	-12	HIS	-	expression tag	UNP F2I8X9
C	-11	HIS	-	expression tag	UNP F2I8X9
C	-10	HIS	-	expression tag	UNP F2I8X9
C	-9	HIS	-	expression tag	UNP F2I8X9
C	-8	HIS	-	expression tag	UNP F2I8X9
C	-7	SER	-	expression tag	UNP F2I8X9
C	-6	SER	-	expression tag	UNP F2I8X9
C	-5	GLY	-	expression tag	UNP F2I8X9
C	-4	LEU	-	expression tag	UNP F2I8X9
C	-3	VAL	-	expression tag	UNP F2I8X9
C	-2	PRO	-	expression tag	UNP F2I8X9
C	-1	ARG	-	expression tag	UNP F2I8X9
C	0	GLY	-	expression tag	UNP F2I8X9
C	1	SER	-	expression tag	UNP F2I8X9
C	123	DAH	TYR	see remark 999	UNP F2I8X9
D	-17	MET	-	initiating methionine	UNP F2I8X9
D	-16	GLY	-	expression tag	UNP F2I8X9
D	-15	SER	-	expression tag	UNP F2I8X9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-14	SER	-	expression tag	UNP F2I8X9
D	-13	HIS	-	expression tag	UNP F2I8X9
D	-12	HIS	-	expression tag	UNP F2I8X9
D	-11	HIS	-	expression tag	UNP F2I8X9
D	-10	HIS	-	expression tag	UNP F2I8X9
D	-9	HIS	-	expression tag	UNP F2I8X9
D	-8	HIS	-	expression tag	UNP F2I8X9
D	-7	SER	-	expression tag	UNP F2I8X9
D	-6	SER	-	expression tag	UNP F2I8X9
D	-5	GLY	-	expression tag	UNP F2I8X9
D	-4	LEU	-	expression tag	UNP F2I8X9
D	-3	VAL	-	expression tag	UNP F2I8X9
D	-2	PRO	-	expression tag	UNP F2I8X9
D	-1	ARG	-	expression tag	UNP F2I8X9
D	0	GLY	-	expression tag	UNP F2I8X9
D	1	SER	-	expression tag	UNP F2I8X9
D	123	DAH	TYR	see remark 999	UNP F2I8X9

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Ca 1 1	0	0
2	A	3	Total Ca 3 3	0	0
2	D	1	Total Ca 1 1	0	0
2	C	1	Total Ca 1 1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	196	Total	O	0	0
			196	196		
4	B	166	Total	O	0	0
			166	166		
4	C	159	Total	O	0	0
			159	159		
4	D	131	Total	O	0	0
			131	131		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ribonucleoside-diphosphate reductase, beta subunit




- Molecule 1: Ribonucleoside-diphosphate reductase, beta subunit



- Molecule 1: Ribonucleoside-diphosphate reductase, beta subunit



- Molecule 1: Ribonucleoside-diphosphate reductase, beta subunit



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	80.00Å 109.42Å 83.86Å 90.00° 90.06° 90.00°	Depositor
Resolution (Å)	35.18 – 1.59 35.18 – 1.59	Depositor EDS
% Data completeness (in resolution range)	94.9 (35.18-1.59) 95.9 (35.18-1.59)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.56 (at 1.58Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.163 , 0.179 0.180 , 0.198	Depositor DCC
R_{free} test set	9317 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	13.2	Xtriage
Anisotropy	0.055	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 42.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.000 for l,k,-h 0.146 for h,-k,-l 0.006 for l,-k,h	Xtriage
Reported twinning fraction	0.544 for H, K, L 0.456 for -h,-k,l	Depositor
Outliers	2 of 186690 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	10705	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA, DAH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/2516	0.55	0/3416
1	B	0.37	0/2588	0.56	0/3513
1	C	0.38	0/2527	0.54	0/3430
1	D	0.38	0/2588	0.53	0/3513
All	All	0.38	0/10219	0.55	0/13872

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2468	0	2404	0	0
1	B	2538	0	2457	2	0
1	C	2479	0	2416	1	0
1	D	2538	0	2457	2	0
2	A	3	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	12	0	14	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	12	0	12	0	0
4	A	196	0	0	0	0
4	B	166	0	0	0	0
4	C	159	0	0	0	0
4	D	131	0	0	0	0
All	All	10705	0	9760	5	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (5) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:85:ASP:OD2	1:C:123:DAH:OE2	2.27	0.52
1:B:85:ASP:OD2	1:B:123:DAH:OE2	2.33	0.46
1:D:85:ASP:OD2	1:D:123:DAH:OE2	2.35	0.45
1:D:221:TYR:CZ	1:D:225:VAL:HG11	2.52	0.44
1:B:221:TYR:CZ	1:B:225:VAL:HG11	2.55	0.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	300/355 (84%)	298 (99%)	2 (1%)	0	100	100
1	B	308/355 (87%)	305 (99%)	3 (1%)	0	100	100
1	C	301/355 (85%)	299 (99%)	2 (1%)	0	100	100
1	D	308/355 (87%)	306 (99%)	2 (1%)	0	100	100
All	All	1217/1420 (86%)	1208 (99%)	9 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	265/310 (86%)	263 (99%)	2 (1%)	81	70
1	B	272/310 (88%)	272 (100%)	0	100	100
1	C	266/310 (86%)	264 (99%)	2 (1%)	81	70
1	D	272/310 (88%)	271 (100%)	1 (0%)	91	84
All	All	1075/1240 (87%)	1070 (100%)	5 (0%)	88	80

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	53	ASN
1	A	121	ARG
1	C	4	ASN
1	C	121	ARG
1	D	121	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	20	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	DAH	C	123	1	12,13,14	1.77	1 (8%)	14,17,19	1.29	2 (14%)
1	DAH	A	123	1	12,13,14	1.62	1 (8%)	14,17,19	0.92	1 (7%)
1	DAH	D	123	1	12,13,14	1.59	1 (8%)	14,17,19	1.04	1 (7%)
1	DAH	B	123	1	12,13,14	1.71	1 (8%)	14,17,19	1.14	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	DAH	C	123	1	-	0/5/6/8	0/1/1/1
1	DAH	A	123	1	-	0/5/6/8	0/1/1/1
1	DAH	D	123	1	-	0/5/6/8	0/1/1/1
1	DAH	B	123	1	-	0/5/6/8	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	123	DAH	CZ-CE2	5.91	1.49	1.40
1	B	123	DAH	CZ-CE2	5.70	1.49	1.40
1	D	123	DAH	CZ-CE2	5.28	1.48	1.40
1	A	123	DAH	CZ-CE2	5.17	1.48	1.40

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	123	DAH	CB-CA-C	-2.62	106.55	111.47
1	D	123	DAH	CG-CB-CA	-2.47	109.10	114.10
1	C	123	DAH	CD1-CG-CD2	2.17	121.58	118.54
1	B	123	DAH	CD1-CG-CD2	2.03	121.39	118.54
1	A	123	DAH	CB-CG-CD2	-2.03	116.96	120.44
1	B	123	DAH	CB-CA-C	-2.02	107.68	111.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	C	123	DAH	1	0
1	D	123	DAH	1	0
1	B	123	DAH	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 6 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	C	403	2	5,5,5	0.27	0	5,5,5	0.23	0
3	GOL	C	402	2	5,5,5	0.25	0	5,5,5	0.20	0
3	GOL	A	405	2	5,5,5	0.29	0	5,5,5	0.21	0
3	GOL	A	404	2	5,5,5	0.29	0	5,5,5	0.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	403	2	-	0/4/4/4	-
3	GOL	C	402	2	-	0/4/4/4	-
3	GOL	A	405	2	-	0/4/4/4	-
3	GOL	A	404	2	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	404	GOL	O2-C2-C3-O3
3	A	404	GOL	C1-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	302/355 (85%)	0.31	11 (3%) 42 40	6, 11, 21, 26	0
1	B	310/355 (87%)	0.38	13 (4%) 36 33	6, 13, 23, 31	0
1	C	303/355 (85%)	0.31	11 (3%) 42 40	8, 12, 21, 34	0
1	D	310/355 (87%)	0.45	13 (4%) 36 33	8, 15, 26, 35	0
All	All	1225/1420 (86%)	0.36	48 (3%) 39 36	6, 13, 23, 35	0

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	307	ARG	4.9
1	B	310	GLU	4.7
1	A	53	ASN	4.6
1	A	305	SER	4.6
1	D	311	ASN	3.8
1	B	307	ARG	3.3
1	D	231	GLU	3.3
1	B	311	ASN	3.2
1	A	232	LYS	3.2
1	B	4	ASN	3.2
1	B	53	ASN	3.1
1	D	53	ASN	3.0
1	A	231	GLU	3.0
1	D	227	LYS	3.0
1	D	291	LYS	2.9
1	B	292	GLU	2.9
1	D	314	PHE	2.8
1	D	292	GLU	2.8
1	A	163	ASP	2.8
1	C	231	GLU	2.7
1	A	306	ALA	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	291	LYS	2.6
1	D	228	LEU	2.6
1	C	4	ASN	2.6
1	C	7	ASP	2.5
1	B	314	PHE	2.5
1	D	87	VAL	2.4
1	C	292	GLU	2.4
1	D	310	GLU	2.4
1	C	305	SER	2.3
1	A	304	LEU	2.3
1	B	43	ARG	2.3
1	C	301	PHE	2.3
1	D	232	LYS	2.3
1	B	55	PRO	2.3
1	B	65	GLU	2.3
1	C	53	ASN	2.2
1	A	301	PHE	2.2
1	B	81	LEU	2.1
1	C	288	PRO	2.1
1	C	163	ASP	2.1
1	C	291	LYS	2.1
1	A	84	LEU	2.1
1	A	291	LYS	2.1
1	B	21	SER	2.0
1	D	4	ASN	2.0
1	D	56	VAL	2.0
1	A	116	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	DAH	B	123	13/14	0.89	0.14	12,15,18,18	0
1	DAH	A	123	13/14	0.90	0.13	11,13,15,15	0
1	DAH	C	123	13/14	0.90	0.13	11,13,16,16	0
1	DAH	D	123	13/14	0.91	0.14	14,17,19,20	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	C	403	6/6	0.77	0.19	21,21,21,21	0
3	GOL	A	405	6/6	0.84	0.12	24,24,24,24	0
3	GOL	C	402	6/6	0.90	0.14	18,19,19,19	0
3	GOL	A	404	6/6	0.91	0.12	19,20,20,20	0
2	CA	A	403	1/1	0.99	0.06	20,20,20,20	0
2	CA	D	401	1/1	0.99	0.04	13,13,13,13	0
2	CA	A	402	1/1	0.99	0.04	16,16,16,16	0
2	CA	B	401	1/1	0.99	0.06	11,11,11,11	0
2	CA	C	401	1/1	1.00	0.04	11,11,11,11	0
2	CA	A	401	1/1	1.00	0.04	12,12,12,12	0

6.5 Other polymers [i](#)

There are no such residues in this entry.