



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 08:35 pm BST

PDB ID : 4ECI
Title : Crystal structure of glutathione s-transferase prk13972 (target efi-501853) from pseudomonas aeruginosa pacs2 complexed with acetate
Authors : Patskovsky, Y.; Toro, R.; Bhosle, R.; Zencheck, W.D.; Hillerich, B.; Seidel, R.D.; Washington, E.; Scott Glenn, A.; Chowdhury, S.; Evans, B.; Hammonds, J.; Imker, H.J.; Armstrong, R.N.; Gerlt, J.A.; Almo, S.C.; Enzyme Function Initiative (EFI)
Deposited on : 2012-03-26
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	244	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%;"></div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px;"> <div style="width: 1%; height: 10px; background-color: red;"></div> <div style="width: 77%, height: 10px; background-color: green;"></div> <div style="width: 6%, height: 10px; background-color: yellow;"></div> <div style="width: 16%, height: 10px; background-color: grey;"></div> </div> </div> <div> 77% 6% 16% </div> </div>
1	B	244	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%;"></div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px;"> <div style="width: 4%, height: 10px; background-color: red;"></div> <div style="width: 76%, height: 10px; background-color: green;"></div> <div style="width: 7%, height: 10px; background-color: yellow;"></div> <div style="width: 16%, height: 10px; background-color: grey;"></div> </div> </div> <div> 76% 7% 16% </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3516 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called glutathione S-transferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	204	Total	C	N	O	S	0	2	0
			1646	1052	296	292	6			
1	B	204	Total	C	N	O	S	0	4	0
			1651	1057	294	293	7			

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	EXPRESSION TAG	UNP Q02KA8
A	0	VAL	-	EXPRESSION TAG	UNP Q02KA8
A	221	ALA	-	EXPRESSION TAG	UNP Q02KA8
A	222	GLU	-	EXPRESSION TAG	UNP Q02KA8
A	223	ASN	-	EXPRESSION TAG	UNP Q02KA8
A	224	LEU	-	EXPRESSION TAG	UNP Q02KA8
A	225	TYR	-	EXPRESSION TAG	UNP Q02KA8
A	226	PHE	-	EXPRESSION TAG	UNP Q02KA8
A	227	GLN	-	EXPRESSION TAG	UNP Q02KA8
A	228	SER	-	EXPRESSION TAG	UNP Q02KA8
A	229	HIS	-	EXPRESSION TAG	UNP Q02KA8
A	230	HIS	-	EXPRESSION TAG	UNP Q02KA8
A	231	HIS	-	EXPRESSION TAG	UNP Q02KA8
A	232	HIS	-	EXPRESSION TAG	UNP Q02KA8
A	233	HIS	-	EXPRESSION TAG	UNP Q02KA8
A	234	HIS	-	EXPRESSION TAG	UNP Q02KA8
A	235	TRP	-	EXPRESSION TAG	UNP Q02KA8
A	236	SER	-	EXPRESSION TAG	UNP Q02KA8
A	237	HIS	-	EXPRESSION TAG	UNP Q02KA8
A	238	PRO	-	EXPRESSION TAG	UNP Q02KA8
A	239	GLN	-	EXPRESSION TAG	UNP Q02KA8
A	240	PHE	-	EXPRESSION TAG	UNP Q02KA8
A	241	GLU	-	EXPRESSION TAG	UNP Q02KA8
A	242	LYS	-	EXPRESSION TAG	UNP Q02KA8
B	-1	MET	-	EXPRESSION TAG	UNP Q02KA8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	VAL	-	EXPRESSION TAG	UNP Q02KA8
B	221	ALA	-	EXPRESSION TAG	UNP Q02KA8
B	222	GLU	-	EXPRESSION TAG	UNP Q02KA8
B	223	ASN	-	EXPRESSION TAG	UNP Q02KA8
B	224	LEU	-	EXPRESSION TAG	UNP Q02KA8
B	225	TYR	-	EXPRESSION TAG	UNP Q02KA8
B	226	PHE	-	EXPRESSION TAG	UNP Q02KA8
B	227	GLN	-	EXPRESSION TAG	UNP Q02KA8
B	228	SER	-	EXPRESSION TAG	UNP Q02KA8
B	229	HIS	-	EXPRESSION TAG	UNP Q02KA8
B	230	HIS	-	EXPRESSION TAG	UNP Q02KA8
B	231	HIS	-	EXPRESSION TAG	UNP Q02KA8
B	232	HIS	-	EXPRESSION TAG	UNP Q02KA8
B	233	HIS	-	EXPRESSION TAG	UNP Q02KA8
B	234	HIS	-	EXPRESSION TAG	UNP Q02KA8
B	235	TRP	-	EXPRESSION TAG	UNP Q02KA8
B	236	SER	-	EXPRESSION TAG	UNP Q02KA8
B	237	HIS	-	EXPRESSION TAG	UNP Q02KA8
B	238	PRO	-	EXPRESSION TAG	UNP Q02KA8
B	239	GLN	-	EXPRESSION TAG	UNP Q02KA8
B	240	PHE	-	EXPRESSION TAG	UNP Q02KA8
B	241	GLU	-	EXPRESSION TAG	UNP Q02KA8
B	242	LYS	-	EXPRESSION TAG	UNP Q02KA8

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total 4	C 2	O 2	0	0
2	B	1	Total 4	C 2	O 2	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	116	Total 116	O 116	0	0
3	B	95	Total 95	O 95	0	0

- Molecule 1: glutathione S-transferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	84.90Å 99.95Å 51.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.80 45.78 – 1.76	Depositor EDS
% Data completeness (in resolution range)	99.1 (50.00-1.80) 98.1 (45.78-1.76)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.88 (at 1.76Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.175 , 0.209 0.175 , 0.210	Depositor DCC
R_{free} test set	1354 reflections (3.10%)	wwPDB-VP
Wilson B-factor (Å ²)	29.9	Xtriage
Anisotropy	0.592	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 42.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3516	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	2/1691 (0.1%)	0.67	0/2290
1	B	0.60	2/1702 (0.1%)	0.66	0/2305
All	All	0.59	4/3393 (0.1%)	0.66	0/4595

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	164	TRP	CD2-CE2	5.47	1.48	1.41
1	B	170	TRP	CD2-CE2	5.29	1.47	1.41
1	A	170	TRP	CD2-CE2	5.14	1.47	1.41
1	B	184	TRP	CD2-CE2	5.13	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1646	0	1642	10	0
1	B	1651	0	1648	12	0
2	A	4	0	3	0	0
2	B	4	0	3	0	0
3	A	116	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	95	0	0	0	0
All	All	3516	0	3296	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:35:LYS:C	1:B:36:LYS:HG2	1.63	1.13
1:B:35:LYS:O	1:B:36:LYS:HG2	1.65	0.96
1:B:41:PRO:O	1:B:45:ARG:HG3	1.84	0.78
1:B:35:LYS:O	1:B:36:LYS:CG	2.37	0.73
1:B:52:ILE:C	1:B:52:ILE:HD12	2.15	0.67
1:B:35:LYS:C	1:B:36:LYS:CG	2.49	0.66
1:A:1:MET:HB3	1:A:59:ASP:HB2	1.78	0.65
1:A:42:GLU:CD	1:A:42:GLU:H	2.05	0.59
1:A:174:ALA:HB1	1:A:176:ASP:OD1	2.03	0.59
1:A:52:ILE:HB	1:A:53:PRO:HA	1.90	0.53
1:B:32:SER:OG	1:B:35:LYS:HG3	2.09	0.52
1:A:0:VAL:HA	1:A:25:PRO:HB2	1.91	0.52
1:B:18:ALA:O	1:B:22:MET:HG3	2.13	0.49
1:A:128:ARG:NH1	1:B:113:VAL:HG21	2.28	0.48
1:A:36:LYS:HE2	3:A:490:HOH:O	2.17	0.45
1:A:202:ARG:HD3	1:A:202:ARG:HA	1.81	0.43
1:A:52:ILE:HD12	1:A:52:ILE:C	2.39	0.42
1:B:118:PHE:CG	1:B:119:PRO:HD2	2.55	0.42
1:B:39:LYS:NZ	1:B:51[B]:ARG:NH2	2.68	0.42
1:A:54:ALA:HA	1:A:71:ILE:HD11	2.02	0.41
1:B:88:LYS:O	1:B:92[A]:ARG:HG3	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	204/244 (84%)	197 (97%)	7 (3%)	0	100	100
1	B	206/244 (84%)	201 (98%)	5 (2%)	0	100	100
All	All	410/488 (84%)	398 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	168/202 (83%)	165 (98%)	3 (2%)	59	48
1	B	169/202 (84%)	166 (98%)	3 (2%)	59	48
All	All	337/404 (83%)	331 (98%)	6 (2%)	59	48

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	0	VAL
1	A	1	MET
1	A	120	GLU
1	B	36	LYS
1	B	42	GLU
1	B	176	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	60	ASN
1	B	29	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACT	B	301	-	1,3,3	2.08	1 (100%)	0,3,3	0.00	-
2	ACT	A	301	-	1,3,3	2.17	1 (100%)	0,3,3	0.00	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	ACT	CH3-C	2.17	1.51	1.48
2	B	301	ACT	CH3-C	2.08	1.51	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	204/244 (83%)	-0.48	2 (0%) 82 80	25, 34, 58, 101	0
1	B	204/244 (83%)	-0.30	10 (4%) 29 24	26, 36, 71, 125	0
All	All	408/488 (83%)	-0.39	12 (2%) 51 46	25, 35, 66, 125	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	33	PHE	5.7
1	B	32	SER	3.5
1	A	1	MET	3.5
1	B	36	LYS	3.2
1	B	34	ASP	3.1
1	B	35	LYS	3.1
1	B	-1	MET	2.8
1	B	30	ALA	2.3
1	B	117	TYR	2.3
1	B	120	GLU	2.3
1	A	202	ARG	2.2
1	B	31	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ACT	A	301	4/4	0.98	0.08	31,35,35,38	0
2	ACT	B	301	4/4	0.99	0.06	25,30,31,44	0

6.5 Other polymers [i](#)

There are no such residues in this entry.