



# Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 04:23 pm BST

PDB ID : 6EC2  
Title : Structure of HIV-1 CA 1/3-hexamer  
Authors : Summers, B.J.; Xiong, Y.  
Deposited on : 2018-08-07  
Resolution : 3.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1026 (3.48-3.32)
Clashscore	141614	1055 (3.48-3.32)
Ramachandran outliers	138981	1038 (3.48-3.32)
Sidechain outliers	138945	1038 (3.48-3.32)
RSRZ outliers	127900	2173 (3.50-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	231	
1	F	231	
2	C	231	
2	G	231	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6405 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Capsid protein p24.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	212	Total	C	N	O	S	0	9	0
			1633	1029	280	311	13			
1	F	212	Total	C	N	O	S	0	10	0
			1639	1031	286	309	13			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	6	LEU	ILE	conflict	UNP P04591
A	45	CYS	GLU	engineered mutation	UNP P04591
A	54	GLU	THR	engineered mutation	UNP P04591
A	83	LEU	VAL	conflict	UNP P04591
A	120	HIS	ASN	conflict	UNP P04591
A	184	ALA	TRP	engineered mutation	UNP P04591
A	185	ALA	MET	engineered mutation	UNP P04591
A	208	GLY	ALA	conflict	UNP P04591
F	6	LEU	ILE	conflict	UNP P04591
F	45	CYS	GLU	engineered mutation	UNP P04591
F	54	GLU	THR	engineered mutation	UNP P04591
F	83	LEU	VAL	conflict	UNP P04591
F	120	HIS	ASN	conflict	UNP P04591
F	184	ALA	TRP	engineered mutation	UNP P04591
F	185	ALA	MET	engineered mutation	UNP P04591
F	208	GLY	ALA	conflict	UNP P04591


- Molecule 2 is a protein called Capsid protein p24.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	201	Total	C	N	O	S	0	9	0
			1568	993	267	295	13			
2	G	201	Total	C	N	O	S	0	9	0
			1539	977	262	288	12			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	6	LEU	ILE	conflict	UNP P04591
C	14	CYS	ALA	engineered mutation	UNP P04591
C	42	GLU	ALA	engineered mutation	UNP P04591
C	83	LEU	VAL	conflict	UNP P04591
C	120	HIS	ASN	conflict	UNP P04591
C	184	ALA	TRP	engineered mutation	UNP P04591
C	185	ALA	MET	engineered mutation	UNP P04591
C	208	GLY	ALA	conflict	UNP P04591
G	6	LEU	ILE	conflict	UNP P04591
G	14	CYS	ALA	engineered mutation	UNP P04591
G	42	GLU	ALA	engineered mutation	UNP P04591
G	83	LEU	VAL	conflict	UNP P04591
G	120	HIS	ASN	conflict	UNP P04591
G	184	ALA	TRP	engineered mutation	UNP P04591
G	185	ALA	MET	engineered mutation	UNP P04591
G	208	GLY	ALA	conflict	UNP P04591

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	F	1	Total C O 4 2 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	4	Total 4	O 4	0	0
4	C	4	Total 4	O 4	0	0
4	F	6	Total 6	O 6	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Capsid protein p24




#### • Molecule 1: Capsid protein p24




#### • Molecule 2: Capsid protein p24



#### • Molecule 2: Capsid protein p24





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.80 Å 97.47 Å 150.89 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.01 – 3.40 75.45 – 3.40	Depositor EDS
% Data completeness (in resolution range)	99.3 (50.01-3.40) 99.4 (75.45-3.40)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.58 (at 3.41 Å)	Xtriage
Refinement program	REFMAC 5.8.0103, PHENIX 1.12_2829	Depositor
R, $R_{free}$	0.211 , 0.275 0.213 , 0.278	Depositor DCC
$R_{free}$ test set	806 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	66.0	Xtriage
Anisotropy	0.810	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 59.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	6405	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.84% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.81	1/1669 (0.1%)	1.04	7/2271 (0.3%)
1	F	0.76	0/1675	0.90	3/2277 (0.1%)
2	C	0.73	0/1603	0.93	3/2177 (0.1%)
2	G	0.64	0/1573	0.83	0/2139
All	All	0.74	1/6520 (0.0%)	0.93	13/8864 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	150[A]	ILE	C-O	5.49	1.33	1.23

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	39	MET	CG-SD-CE	9.12	114.78	100.20
1	F	143	ARG	NE-CZ-NH1	-7.00	116.80	120.30
2	C	150[A]	ILE	CA-C-N	-6.62	102.64	117.20
2	C	96	MET	CG-SD-CE	6.39	110.42	100.20
1	A	150[A]	ILE	CA-C-O	6.31	133.36	120.10
1	A	136	LEU	CB-CG-CD1	-6.22	100.43	111.00
1	A	150[A]	ILE	CA-C-N	-5.90	104.23	117.20
1	A	150[A]	ILE	N-CA-CB	-5.44	98.30	110.80

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	63	GLN	CB-CA-C	-5.38	99.64	110.40
1	F	198	CYS	CA-CB-SG	-5.33	104.41	114.00
1	F	20	LEU	CB-CG-CD1	-5.28	102.03	111.00
1	A	18	ARG	CB-CA-C	-5.11	100.18	110.40
1	A	198	CYS	CA-CB-SG	-5.11	104.81	114.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	156	GLY	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1633	0	1592	24	0
1	F	1639	0	1602	25	0
2	C	1568	0	1532	15	0
2	G	1539	0	1498	14	0
3	A	4	0	3	0	0
3	C	4	0	3	0	0
3	F	4	0	3	0	0
4	A	4	0	0	0	0
4	C	4	0	0	1	0
4	F	6	0	0	1	0
All	All	6405	0	6233	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:15:ILE:CG2	1:F:54:GLU:OE1	1.87	1.22

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:146:SER:OG	1:A:175:GLU:HG2	1.37	1.19
1:A:148:THR:O	1:A:175:GLU:HG3	1.57	1.04
1:F:15:ILE:HG22	1:F:54:GLU:OE1	1.55	1.04
1:F:15:ILE:HG21	1:F:54:GLU:OE1	1.57	1.03
2:C:150[A]:ILE:HG23	2:C:151:LEU:N	1.78	0.97
2:C:147:PRO:HB2	2:C:175:GLU:OE2	1.64	0.95
2:C:150[A]:ILE:CG2	2:C:151:LEU:H	1.79	0.95
1:A:148:THR:O	1:A:175:GLU:CG	2.21	0.88
1:F:84:HIS:O	1:F:100:ARG:NH1	2.09	0.85
2:C:150[A]:ILE:HG23	2:C:151:LEU:H	1.37	0.81
1:F:157[A]:PRO:HD2	4:F:402:HOH:O	1.83	0.78
1:A:146:SER:HG	1:A:175:GLU:HG2	1.54	0.72
1:A:146:SER:OG	1:A:175:GLU:CG	2.29	0.72
1:A:172:LEU:O	1:A:176:GLN:HB3	1.91	0.70
2:C:144:MET:HE1	1:F:212:GLU:HG3	1.77	0.66
1:A:37:ILE:HD12	1:A:138:LEU:HB3	1.79	0.65
1:A:180:GLU:HG2	2:G:70:LYS:HE2	1.78	0.64
1:A:212:GLU:HG3	2:G:144:MET:HE1	1.81	0.63
2:G:30:LYS:HE2	2:G:163:ASP:OD2	2.02	0.59
2:G:37:ILE:HD12	2:G:138:LEU:HB3	1.84	0.59
2:C:144:MET:CE	1:F:212:GLU:HG3	2.33	0.58
1:A:144:MET:O	1:F:143:ARG:NH1	2.37	0.57
2:G:147:PRO:HG2	2:G:153:ILE:CD1	2.34	0.57
1:A:2:ILE:HD12	1:A:115:ILE:HG23	1.88	0.55
1:A:148:THR:O	1:A:175:GLU:HG2	2.07	0.54
1:A:68:MET:SD	1:A:144:MET:HE1	2.46	0.54
1:A:149:SER:O	1:A:151:LEU:N	2.40	0.54
2:C:147:PRO:CB	2:C:175:GLU:OE2	2.48	0.53
1:A:18:ARG:CZ	2:G:18:ARG:HD2	2.40	0.52
2:G:29:GLU:HB3	2:G:160:PRO:HD3	1.92	0.52
1:F:155:GLN:HE22	1:F:161:PHE:HD1	1.58	0.52
1:F:37:ILE:HD12	1:F:138:LEU:HB3	1.92	0.51
1:F:26[A]:VAL:HG21	1:F:39:MET:HG2	1.93	0.51
1:F:175:GLU:O	1:F:177:ALA:N	2.43	0.50
2:G:37:ILE:HB	2:G:38[A]:PRO:HD3	1.92	0.50
1:A:150[A]:ILE:HG23	1:A:151:LEU:HD23	1.91	0.50
2:G:147:PRO:HG2	2:G:153:ILE:HD11	1.94	0.49
2:C:150[A]:ILE:HG22	2:C:151:LEU:H	1.75	0.48
2:G:30:LYS:HE2	2:G:163:ASP:CG	2.35	0.47
1:A:20:LEU:HD22	1:A:58:THR:HG21	1.97	0.47
1:A:172:LEU:O	1:A:176:GLN:CB	2.61	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:144:MET:HE2	1:A:144:MET:HB2	1.63	0.46
4:C:401:HOH:O	1:F:173:ARG:HD3	2.16	0.45
2:G:170:LYS:HA	2:G:170:LYS:HD2	1.73	0.45
1:F:191:VAL:HG22	1:F:214[A]:MET:HE1	1.98	0.45
1:F:37:ILE:HB	1:F:38[A]:PRO:HD3	1.97	0.45
2:G:147:PRO:HG2	2:G:153:ILE:HD13	1.98	0.44
2:C:140:LYS:HE3	1:F:212:GLU:OE1	2.17	0.44
2:G:37:ILE:CD1	2:G:138:LEU:HB3	2.47	0.44
1:A:172:LEU:HG	1:A:181:VAL:HG11	1.99	0.44
1:A:68:MET:SD	1:A:144:MET:CE	3.06	0.43
1:F:110:THR:OG1	1:F:113:GLU:HG3	2.18	0.43
1:F:191:VAL:CG2	1:F:214[A]:MET:HE1	2.48	0.43
1:F:153:ILE:HG21	1:F:168:PHE:HA	2.02	0.42
2:C:37:ILE:HB	2:C:38:PRO:HD3	2.01	0.42
1:F:69:LEU:O	1:F:73:ILE:HG13	2.19	0.42
1:A:202:LEU:O	1:A:205:LEU:HB2	2.20	0.42
1:F:21:ASN:O	1:F:25:LYS:HG2	2.20	0.42
1:A:161:PHE:O	1:A:165:VAL:HG23	2.20	0.41
2:C:37:ILE:HD11	2:C:142:VAL:HG21	2.02	0.41
2:C:100:ARG:HG2	2:C:103:ASP:OD2	2.21	0.41
1:F:1[A]:PRO:HD3	1:F:15:ILE:HA	2.02	0.41
1:F:7:GLN:OE1	1:F:7:GLN:N	2.53	0.41
1:A:212:GLU:OE2	2:G:140:LYS:HE3	2.21	0.41
1:F:48:THR:HG22	1:F:118:MET:SD	2.61	0.41
1:F:202:LEU:HD22	1:F:214[A]:MET:HB3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	209/231 (90%)	203 (97%)	3 (1%)	3 (1%)	11 37

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	F	209/231 (90%)	199 (95%)	10 (5%)	0	100	100
2	C	195/231 (84%)	193 (99%)	1 (0%)	1 (0%)	29	61
2	G	194/231 (84%)	192 (99%)	2 (1%)	0	100	100
All	All	807/924 (87%)	787 (98%)	16 (2%)	4 (0%)	29	61

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	181	VAL
1	A	157[A]	PRO
2	C	151	LEU
1	A	150[A]	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	174/192 (91%)	168 (97%)	6 (3%)	37	65
1	F	175/192 (91%)	172 (98%)	3 (2%)	60	80
2	C	169/194 (87%)	167 (99%)	2 (1%)	71	85
2	G	162/194 (84%)	159 (98%)	3 (2%)	57	78
All	All	680/772 (88%)	666 (98%)	14 (2%)	53	76

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	PRO
1	A	148	THR
1	A	170	LYS
1	A	183	ASN
1	A	198	CYS
1	A	219	GLN
2	C	98	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	C	150[A]	ILE
1	F	70	LYS
1	F	198	CYS
1	F	219	GLN
2	G	6	LEU
2	G	144	MET
2	G	170	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	183	ASN
2	C	7	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	ACT	A	301	-	1,3,3	2.74	1 (100%)	0,3,3	0.00	-
3	ACT	F	301	-	1,3,3	3.71	1 (100%)	0,3,3	0.00	-
3	ACT	C	301	-	1,3,3	2.66	1 (100%)	0,3,3	0.00	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	301	ACT	CH3-C	3.71	1.53	1.48
3	A	301	ACT	CH3-C	2.74	1.52	1.48
3	C	301	ACT	CH3-C	2.66	1.52	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	212/231 (91%)	-0.10	1 (0%) 91 90	38, 58, 91, 112	1 (0%)
1	F	212/231 (91%)	-0.03	2 (0%) 84 83	38, 59, 100, 133	1 (0%)
2	C	201/231 (87%)	0.76	36 (17%) 1 1	36, 57, 178, 215	2 (0%)
2	G	201/231 (87%)	0.87	35 (17%) 1 1	46, 84, 166, 213	2 (0%)
All	All	826/924 (89%)	0.36	74 (8%) 9 11	36, 62, 160, 215	6 (0%)

All (74) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	197	ASP	8.9
2	C	209	ALA	8.8
2	C	218	CYS	7.6
2	C	210	THR	7.5
2	C	198	CYS	7.4
2	C	212	GLU	7.1
2	C	196[A]	PRO	6.5
2	G	184	ALA	6.1
2	C	168	PHE	5.0
2	G	205	LEU	4.9
2	C	211	LEU	4.7
2	G	153	ILE	4.7
2	C	202	LEU	4.6
2	C	188	THR	4.5
2	C	217	ALA	4.5
2	C	199	LYS	4.2
2	C	208	GLY	4.2
2	G	160	PRO	4.2
2	G	206	GLY	4.2
2	G	218	CYS	4.1
2	G	219	GLN	4.1

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	C	214[A]	MET	4.1
2	C	207	PRO	4.0
2	C	148	THR	3.9
2	G	194	ALA	3.9
2	G	217	ALA	3.8
2	G	207	PRO	3.8
2	C	195	ASN	3.5
2	C	189	LEU	3.3
2	C	173	ARG	3.1
2	C	192	GLN	3.0
2	G	152	ASP	3.0
2	G	189	LEU	3.0
2	G	191	VAL	2.9
2	G	193	ASN	2.9
2	C	169	TYR	2.9
2	G	216	THR	2.9
2	C	215	MET	2.8
2	G	190	LEU	2.8
2	G	169	TYR	2.8
2	G	200	THR	2.8
2	G	161	PHE	2.8
2	G	199	LYS	2.8
2	G	159	GLU	2.7
2	C	201	ILE	2.7
2	G	172	LEU	2.7
2	G	165	VAL	2.7
2	G	198	CYS	2.7
2	C	171	THR	2.6
2	C	213	GLU	2.6
2	C	153	ILE	2.5
2	C	193	ASN	2.5
2	C	164	TYR	2.5
2	G	123	PRO	2.4
2	G	99	PRO	2.4
2	C	172	LEU	2.4
2	G	212	GLU	2.4
2	G	192	GLN	2.3
2	C	155	GLN	2.3
2	G	202	LEU	2.3
2	C	200	THR	2.3
2	G	215	MET	2.3
2	G	173	ARG	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	C	152	ASP	2.3
2	G	164	TYR	2.3
2	C	203	LYS	2.2
1	A	60	GLY	2.2
2	G	197	ASP	2.2
2	G	162	ARG	2.2
2	C	161	PHE	2.1
1	F	87	HIS	2.1
2	C	191	VAL	2.1
2	G	201	ILE	2.0
1	F	80	TRP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	ACT	C	301	4/4	0.68	0.37	89,95,98,100	0
3	ACT	F	301	4/4	0.88	0.20	59,63,63,64	0
3	ACT	A	301	4/4	0.92	0.27	69,76,76,76	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.