



Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 03:26 am BST

PDB ID : 3EEV
Title : Crystal Structure of Chloramphenicol Acetyltransferase VCA0300 from *Vibrio cholerae* O1 biovar eltor
Authors : Kim, Y.; Maltseva, N.; Kwon, K.; Anderson, W.F.; Joachimiak, A.; Center for Structural Genomics of Infectious Diseases (CSGID)
Deposited on : 2008-09-05
Resolution : 2.61 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 2.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3797 (2.64-2.60)
Clashscore	141614	4168 (2.64-2.60)
Ramachandran outliers	138981	4093 (2.64-2.60)
Sidechain outliers	138945	4093 (2.64-2.60)
RSRZ outliers	127900	3731 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	212	
1	B	212	
1	C	212	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MPD	C	210	X	-	-	-
2	MPD	C	211	X	-	-	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5176 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Chloramphenicol acetyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	207	Total	C	N	O	S	0	1	0
			1653	1056	278	309	10			
1	B	205	Total	C	N	O	S	0	0	0
			1628	1042	271	304	11			
1	C	208	Total	C	N	O	S	0	0	0
			1653	1057	278	307	11			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	EXPRESSION TAG	UNP Q9KMN1
A	-1	ASN	-	EXPRESSION TAG	UNP Q9KMN1
A	0	ALA	-	EXPRESSION TAG	UNP Q9KMN1
B	-2	SER	-	EXPRESSION TAG	UNP Q9KMN1
B	-1	ASN	-	EXPRESSION TAG	UNP Q9KMN1
B	0	ALA	-	EXPRESSION TAG	UNP Q9KMN1
C	-2	SER	-	EXPRESSION TAG	UNP Q9KMN1
C	-1	ASN	-	EXPRESSION TAG	UNP Q9KMN1
C	0	ALA	-	EXPRESSION TAG	UNP Q9KMN1

- Molecule 2 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			8	6	2		
2	C	1	Total	C	O	0	0
			8	6	2		
2	C	1	Total	C	O	0	0
			8	6	2		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	66	Total	O	0	0
			66	66		
3	B	65	Total	O	0	0
			65	65		
3	C	87	Total	O	0	0
			87	87		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Chloramphenicol acetyltransferase

Chain A: 





• Molecule 1: Chloramphenicol acetyltransferase

Chain B: 



• Molecule 1: Chloramphenicol acetyltransferase

Chain C: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	99.97Å 99.97Å 127.37Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	32.70 – 2.61 32.72 – 2.61	Depositor EDS
% Data completeness (in resolution range)	99.7 (32.70-2.61) 99.7 (32.72-2.61)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.48 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.5.0053	Depositor
R, R_{free}	0.175 , 0.242 0.171 , 0.238	Depositor DCC
R_{free} test set	1177 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	48.6	Xtriage
Anisotropy	0.421	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 50.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.019 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5176	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.81	0/1704	0.82	2/2313 (0.1%)
1	B	0.81	0/1679	0.77	0/2280
1	C	0.86	0/1704	0.78	2/2312 (0.1%)
All	All	0.83	0/5087	0.79	4/6905 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	108	ASP	CB-CG-OD1	5.65	123.38	118.30
1	A	208	ARG	NE-CZ-NH2	5.64	123.12	120.30
1	C	124	MET	CG-SD-CE	5.12	108.39	100.20
1	C	184	LEU	CA-CB-CG	-5.02	103.76	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1653	0	1549	52	0
1	B	1628	0	1527	39	0
1	C	1653	0	1558	38	0
2	A	8	0	14	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	16	0	28	6	0
3	A	66	0	0	4	0
3	B	65	0	0	3	0
3	C	87	0	0	4	0
All	All	5176	0	4676	116	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (116) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:149:GLU:OE2	1:A:158:HIS:HD2	1.43	1.00
1:A:5:THR:HG22	3:A:215:HOH:O	1.71	0.91
1:B:196:ILE:HD13	1:C:83:ILE:HG12	1.59	0.83
1:A:113:HIS:H	1:A:131:HIS:HD2	1.26	0.81
1:B:122:MET:HG2	1:B:124:MET:CE	2.12	0.79
1:A:194:SER:HB3	3:A:275:HOH:O	1.83	0.79
1:C:171:LEU:HD11	1:C:196:ILE:HD12	1.64	0.77
1:B:37:HIS:HE1	3:B:321:HOH:O	1.65	0.77
1:B:149:GLU:OE2	1:B:158:HIS:HD2	1.68	0.76
1:A:149:GLU:OE2	1:A:158:HIS:CD2	2.35	0.74
1:C:116:TRP:CZ2	2:C:210:MPD:HM1	2.24	0.72
1:C:71:MET:HG3	1:C:124:MET:CE	2.20	0.72
1:A:166:VAL:HG12	1:A:170:MET:CE	2.21	0.71
1:C:1:MET:HB2	3:C:399:HOH:O	1.91	0.71
1:C:116:TRP:HZ2	2:C:210:MPD:HM1	1.58	0.69
1:A:124:MET:HE2	1:A:124:MET:HA	1.75	0.69
1:B:60:SER:H	1:B:113:HIS:HD2	1.38	0.68
1:C:206:LYS:HG3	2:C:211:MPD:HM2	1.75	0.68
1:A:166:VAL:HG12	1:A:170:MET:HE3	1.74	0.66
1:C:195:ASP:HB3	3:C:416:HOH:O	1.95	0.66
1:B:171:LEU:HD11	1:B:196:ILE:HD12	1.76	0.66
1:A:188:MET:HE3	1:B:96:PHE:CD1	2.31	0.66
1:A:113:HIS:H	1:A:131:HIS:CD2	2.13	0.66
1:C:49:GLU:CG	3:C:236:HOH:O	2.43	0.65
1:A:99:ALA:HB2	1:C:188:MET:HE1	1.77	0.65
1:A:94:ASP:OD1	1:A:94:ASP:N	2.30	0.64
1:A:122:MET:HG2	1:A:124:MET:HE3	1.79	0.64
1:C:174:MET:HE2	1:C:200:TYR:HD1	1.63	0.63
1:B:113:HIS:H	1:B:131:HIS:HD2	1.48	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:196:ILE:HD13	1:B:83:ILE:HG12	1.82	0.62
1:C:71:MET:HG3	1:C:124:MET:HE2	1.83	0.61
2:A:210:MPD:H13	2:A:210:MPD:O4	2.03	0.59
1:C:206:LYS:HG3	2:C:211:MPD:CM	2.35	0.56
1:C:60:SER:H	1:C:113:HIS:HD2	1.53	0.56
1:B:37:HIS:HD2	1:B:38:SER:O	1.89	0.55
1:B:122:MET:HG2	1:B:124:MET:HE1	1.85	0.55
1:A:39:PHE:O	1:A:42:CYS:HB2	2.06	0.55
1:C:152:GLY:O	1:C:153:SER:HB3	2.07	0.54
1:A:152:GLY:O	1:A:153:SER:CB	2.56	0.54
1:A:83:ILE:HG12	1:C:196:ILE:HD13	1.89	0.53
1:B:149:GLU:OE2	1:B:158:HIS:CD2	2.55	0.53
1:C:183:TRP:O	1:C:187:SER:OG	2.26	0.53
1:B:60:SER:H	1:B:113:HIS:CD2	2.23	0.53
1:A:166:VAL:O	1:A:170:MET:HG3	2.09	0.53
1:A:60:SER:HB2	1:A:113:HIS:ND1	2.24	0.52
1:A:60:SER:HB3	1:A:61:PHE:CD2	2.45	0.52
1:A:204:GLN:O	1:A:208:ARG:HG2	2.10	0.51
1:B:113:HIS:H	1:B:131:HIS:CD2	2.28	0.51
1:A:174:MET:HE1	1:A:199:LEU:HD23	1.92	0.51
1:C:174:MET:HE2	1:C:200:TYR:CD1	2.46	0.51
1:A:174:MET:CE	1:A:199:LEU:HD23	2.41	0.51
1:A:171:LEU:HD11	1:A:196:ILE:HD12	1.93	0.50
1:B:113:HIS:N	1:B:131:HIS:HD2	2.10	0.50
1:C:152:GLY:O	1:C:153:SER:CB	2.59	0.50
1:A:26:LYS:HD2	1:A:27:HIS:CE1	2.46	0.50
1:A:166:VAL:HG12	1:A:170:MET:HE2	1.92	0.49
1:A:122:MET:SD	1:A:124:MET:HE1	2.53	0.49
1:C:124:MET:CE	1:C:124:MET:HA	2.43	0.49
1:A:188:MET:HE3	1:B:96:PHE:HD1	1.77	0.48
1:C:49:GLU:HG3	3:C:236:HOH:O	2.07	0.48
1:A:113:HIS:N	1:A:131:HIS:HD2	2.04	0.48
1:C:174:MET:CE	1:C:200:TYR:HD1	2.26	0.48
1:B:22:ILE:HG22	1:B:24:VAL:HG13	1.96	0.48
1:A:202:ASN:OD1	1:A:206:LYS:HE3	2.14	0.47
1:C:37:HIS:HD2	1:C:38:SER:O	1.96	0.47
1:B:174:MET:HE2	1:B:179:TRP:CH2	2.49	0.47
1:A:152:GLY:O	1:A:156:ALA:HA	2.14	0.47
1:C:171:LEU:HD11	1:C:196:ILE:CD1	2.39	0.47
1:A:88:PHE:CB	1:C:188:MET:HE1	2.45	0.47
1:A:152:GLY:O	1:A:153:SER:HB3	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:139:SER:OG	1:C:152:GLY:HA2	2.14	0.46
1:A:188:MET:CE	1:B:96:PHE:HA	2.45	0.46
1:B:174:MET:HE2	1:B:179:TRP:HH2	1.80	0.46
2:C:210:MPD:H52	2:C:210:MPD:O2	2.16	0.45
1:B:149:GLU:HB2	1:B:161:PHE:CE1	2.52	0.45
1:B:26:LYS:HD2	1:B:27:HIS:CD2	2.52	0.45
1:A:158:HIS:HE1	3:A:264:HOH:O	2.01	0.44
1:B:180:PRO:HD2	1:B:183:TRP:CD2	2.51	0.44
1:C:184:LEU:HD12	1:C:184:LEU:HA	1.77	0.44
1:A:122:MET:HG2	1:A:124:MET:CE	2.47	0.44
1:C:186:GLU:OE1	2:C:211:MPD:H31	2.17	0.44
1:C:174:MET:HE2	1:C:200:TYR:HA	1.99	0.44
1:A:22:ILE:HD13	1:A:43:VAL:HG21	1.99	0.44
1:B:174:MET:HB2	1:B:203:TRP:CD2	2.53	0.43
1:A:125:PRO:HB3	3:A:227:HOH:O	2.18	0.43
1:B:122:MET:CG	1:B:124:MET:CE	2.90	0.43
1:B:37:HIS:CE1	3:B:321:HOH:O	2.53	0.43
1:A:30:TYR:CE2	1:A:32:GLY:HA3	2.54	0.43
1:B:60:SER:N	1:B:113:HIS:HD2	2.12	0.43
1:C:140:VAL:HG23	1:C:154:ASN:HB2	2.01	0.43
1:A:207:ALA:C	1:A:208:ARG:HD2	2.39	0.43
1:B:164:SER:HA	3:B:368:HOH:O	2.17	0.43
1:B:122:MET:SD	1:B:124:MET:HE3	2.59	0.43
1:C:60:SER:H	1:C:113:HIS:CD2	2.33	0.43
1:A:139:SER:OG	1:A:152:GLY:HA2	2.18	0.42
1:C:174:MET:O	1:C:175:ALA:C	2.58	0.42
1:A:93:ASN:HB3	1:A:96:PHE:CD2	2.53	0.42
1:A:181:GLU:OE2	1:B:95:ASN:ND2	2.52	0.42
1:A:19:ASN:HA	1:A:20:PRO:HD3	1.85	0.42
1:B:174:MET:HA	1:B:203:TRP:CZ2	2.54	0.42
1:C:174:MET:HE1	1:C:200:TYR:HB2	2.02	0.42
1:B:139:SER:OG	1:B:152:GLY:HA2	2.20	0.41
1:A:99:ALA:HB2	1:C:188:MET:CE	2.48	0.41
1:A:82:TRP:CZ3	1:C:192:CYS:HB3	2.55	0.41
1:B:72:MET:O	1:B:73:ALA:HB3	2.20	0.41
1:B:86:PHE:HA	1:B:87:PRO:HD3	1.96	0.41
1:A:134:ILE:HB	1:A:150:VAL:HG22	2.03	0.41
1:B:152:GLY:O	1:B:153:SER:CB	2.69	0.41
1:A:116:TRP:CZ2	2:A:210:MPD:H11	2.55	0.41
1:B:122:MET:SD	1:B:124:MET:CE	3.09	0.41
1:A:176:TRP:HA	1:A:179:TRP:CE2	2.57	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:116:TRP:CH2	1:C:78:HIS:HB2	2.55	0.40
1:A:208:ARG:HD2	1:A:208:ARG:N	2.36	0.40
1:A:88:PHE:HB3	1:C:188:MET:HE1	2.03	0.40
1:A:188:MET:HE3	1:B:96:PHE:HA	2.03	0.40
1:B:181:GLU:HG2	1:C:95:ASN:HD22	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	206/212 (97%)	196 (95%)	9 (4%)	1 (0%)	29	50
1	B	203/212 (96%)	196 (97%)	5 (2%)	2 (1%)	15	30
1	C	206/212 (97%)	194 (94%)	11 (5%)	1 (0%)	29	50
All	All	615/636 (97%)	586 (95%)	25 (4%)	4 (1%)	22	41

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	153	SER
1	C	153	SER
1	B	89	PHE
1	A	153	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	178/181 (98%)	168 (94%)	10 (6%)	21	40
1	B	176/181 (97%)	165 (94%)	11 (6%)	18	35
1	C	178/181 (98%)	164 (92%)	14 (8%)	12	23
All	All	532/543 (98%)	497 (93%)	35 (7%)	16	32

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	THR
1	A	6	SER
1	A	18	THR
1	A	91	GLN
1	A	94	ASP
1	A	106	SER
1	A	127	VAL
1	A	186	GLU
1	A	188	MET
1	A	190	SER
1	B	1	MET
1	B	6	SER
1	B	50	ARG
1	B	91	GLN
1	B	92	ASP
1	B	106	SER
1	B	127	VAL
1	B	128	LYS
1	B	181	GLU
1	B	189	GLN
1	B	190	SER
1	C	1	MET
1	C	18	THR
1	C	49	GLU
1	C	50	ARG
1	C	92	ASP
1	C	124	MET
1	C	127	VAL
1	C	157	LYS
1	C	160	LYS
1	C	186	GLU
1	C	187	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	196	ILE
1	C	202	ASN
1	C	208	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	GLN
1	A	27	HIS
1	A	78	HIS
1	A	91	GLN
1	A	131	HIS
1	A	158	HIS
1	B	15	GLN
1	B	27	HIS
1	B	37	HIS
1	B	113	HIS
1	B	131	HIS
1	B	158	HIS
1	B	189	GLN
1	C	113	HIS
1	C	178	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MPD	C	210	-	7,7,7	0.67	0	9,10,10	1.38	1 (11%)
2	MPD	C	211	-	7,7,7	0.28	0	9,10,10	0.80	0
2	MPD	A	210	-	7,7,7	0.84	0	9,10,10	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MPD	C	210	-	1/1/2/2	4/5/5/5	-
2	MPD	C	211	-	1/1/2/2	1/5/5/5	-
2	MPD	A	210	-	-	2/5/5/5	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	210	MPD	CM-C2-C1	-3.21	103.88	110.57

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	C	210	MPD	C4
2	C	211	MPD	C4

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	210	MPD	C2-C3-C4-O4
2	C	210	MPD	O2-C2-C3-C4
2	C	210	MPD	C2-C3-C4-C5
2	C	210	MPD	C1-C2-C3-C4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	C	210	MPD	CM-C2-C3-C4
2	C	211	MPD	C2-C3-C4-C5
2	A	210	MPD	C2-C3-C4-C5

There are no ring outliers.

3 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	210	MPD	3	0
2	C	211	MPD	3	0
2	A	210	MPD	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	207/212 (97%)	-0.29	1 (0%) 91 89	35, 43, 57, 69	0
1	B	205/212 (96%)	-0.29	1 (0%) 91 89	36, 45, 57, 66	0
1	C	208/212 (98%)	-0.35	0 100 100	33, 40, 55, 67	0
All	All	620/636 (97%)	-0.31	2 (0%) 94 93	33, 43, 57, 69	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	49	GLU	3.1
1	B	97	ALA	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MPD	C	210	8/8	0.94	0.20	44,47,48,48	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MPD	A	210	8/8	0.95	0.12	38,43,44,46	0
2	MPD	C	211	8/8	0.96	0.32	56,59,61,63	0

6.5 Other polymers [i](#)

There are no such residues in this entry.