



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 09:11 pm BST

PDB ID : 4EGJ
Title : Crystal structure of D-alanine-D-alanine ligase from Burkholderia xenovorans
Authors : SSGCID; Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2012-03-31
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	334	<div> <div>3%</div> <div> <div></div> <div>81%</div> <div>•</div> <div>15%</div> </div> </div>
1	B	334	<div> <div>2%</div> <div> <div></div> <div>78%</div> <div>7%</div> <div>•</div> <div>15%</div> </div> </div>
1	C	334	<div> <div>12%</div> <div> <div></div> <div>81%</div> <div>•</div> <div>15%</div> </div> </div>
1	D	334	<div> <div>3%</div> <div> <div></div> <div>70%</div> <div>•</div> <div>26%</div> </div> </div>

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 8430 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-alanine–D-alanine ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	285	Total	C	N	O	S	0	3	0
			2118	1357	365	390	6			
1	B	285	Total	C	N	O	S	0	2	0
			2099	1344	362	387	6			
1	C	283	Total	C	N	O	S	0	0	0
			2033	1291	352	385	5			
1	D	246	Total	C	N	O	S	0	0	0
			1822	1162	314	341	5			

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	INITIATING METHIONINE	UNP Q13TZ4
A	-19	ALA	-	EXPRESSION TAG	UNP Q13TZ4
A	-18	HIS	-	EXPRESSION TAG	UNP Q13TZ4
A	-17	HIS	-	EXPRESSION TAG	UNP Q13TZ4
A	-16	HIS	-	EXPRESSION TAG	UNP Q13TZ4
A	-15	HIS	-	EXPRESSION TAG	UNP Q13TZ4
A	-14	HIS	-	EXPRESSION TAG	UNP Q13TZ4
A	-13	HIS	-	EXPRESSION TAG	UNP Q13TZ4
A	-12	MET	-	EXPRESSION TAG	UNP Q13TZ4
A	-11	GLY	-	EXPRESSION TAG	UNP Q13TZ4
A	-10	THR	-	EXPRESSION TAG	UNP Q13TZ4
A	-9	LEU	-	EXPRESSION TAG	UNP Q13TZ4
A	-8	GLU	-	EXPRESSION TAG	UNP Q13TZ4
A	-7	ALA	-	EXPRESSION TAG	UNP Q13TZ4
A	-6	GLN	-	EXPRESSION TAG	UNP Q13TZ4
A	-5	THR	-	EXPRESSION TAG	UNP Q13TZ4
A	-4	GLN	-	EXPRESSION TAG	UNP Q13TZ4
A	-3	GLY	-	EXPRESSION TAG	UNP Q13TZ4
A	-2	PRO	-	EXPRESSION TAG	UNP Q13TZ4
A	-1	GLY	-	EXPRESSION TAG	UNP Q13TZ4
A	0	SER	-	EXPRESSION TAG	UNP Q13TZ4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-20	MET	-	INITIATING METHIONINE	UNP Q13TZ4
B	-19	ALA	-	EXPRESSION TAG	UNP Q13TZ4
B	-18	HIS	-	EXPRESSION TAG	UNP Q13TZ4
B	-17	HIS	-	EXPRESSION TAG	UNP Q13TZ4
B	-16	HIS	-	EXPRESSION TAG	UNP Q13TZ4
B	-15	HIS	-	EXPRESSION TAG	UNP Q13TZ4
B	-14	HIS	-	EXPRESSION TAG	UNP Q13TZ4
B	-13	HIS	-	EXPRESSION TAG	UNP Q13TZ4
B	-12	MET	-	EXPRESSION TAG	UNP Q13TZ4
B	-11	GLY	-	EXPRESSION TAG	UNP Q13TZ4
B	-10	THR	-	EXPRESSION TAG	UNP Q13TZ4
B	-9	LEU	-	EXPRESSION TAG	UNP Q13TZ4
B	-8	GLU	-	EXPRESSION TAG	UNP Q13TZ4
B	-7	ALA	-	EXPRESSION TAG	UNP Q13TZ4
B	-6	GLN	-	EXPRESSION TAG	UNP Q13TZ4
B	-5	THR	-	EXPRESSION TAG	UNP Q13TZ4
B	-4	GLN	-	EXPRESSION TAG	UNP Q13TZ4
B	-3	GLY	-	EXPRESSION TAG	UNP Q13TZ4
B	-2	PRO	-	EXPRESSION TAG	UNP Q13TZ4
B	-1	GLY	-	EXPRESSION TAG	UNP Q13TZ4
B	0	SER	-	EXPRESSION TAG	UNP Q13TZ4
C	-20	MET	-	INITIATING METHIONINE	UNP Q13TZ4
C	-19	ALA	-	EXPRESSION TAG	UNP Q13TZ4
C	-18	HIS	-	EXPRESSION TAG	UNP Q13TZ4
C	-17	HIS	-	EXPRESSION TAG	UNP Q13TZ4
C	-16	HIS	-	EXPRESSION TAG	UNP Q13TZ4
C	-15	HIS	-	EXPRESSION TAG	UNP Q13TZ4
C	-14	HIS	-	EXPRESSION TAG	UNP Q13TZ4
C	-13	HIS	-	EXPRESSION TAG	UNP Q13TZ4
C	-12	MET	-	EXPRESSION TAG	UNP Q13TZ4
C	-11	GLY	-	EXPRESSION TAG	UNP Q13TZ4
C	-10	THR	-	EXPRESSION TAG	UNP Q13TZ4
C	-9	LEU	-	EXPRESSION TAG	UNP Q13TZ4
C	-8	GLU	-	EXPRESSION TAG	UNP Q13TZ4
C	-7	ALA	-	EXPRESSION TAG	UNP Q13TZ4
C	-6	GLN	-	EXPRESSION TAG	UNP Q13TZ4
C	-5	THR	-	EXPRESSION TAG	UNP Q13TZ4
C	-4	GLN	-	EXPRESSION TAG	UNP Q13TZ4
C	-3	GLY	-	EXPRESSION TAG	UNP Q13TZ4
C	-2	PRO	-	EXPRESSION TAG	UNP Q13TZ4
C	-1	GLY	-	EXPRESSION TAG	UNP Q13TZ4
C	0	SER	-	EXPRESSION TAG	UNP Q13TZ4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-20	MET	-	INITIATING METHIONINE	UNP Q13TZ4
D	-19	ALA	-	EXPRESSION TAG	UNP Q13TZ4
D	-18	HIS	-	EXPRESSION TAG	UNP Q13TZ4
D	-17	HIS	-	EXPRESSION TAG	UNP Q13TZ4
D	-16	HIS	-	EXPRESSION TAG	UNP Q13TZ4
D	-15	HIS	-	EXPRESSION TAG	UNP Q13TZ4
D	-14	HIS	-	EXPRESSION TAG	UNP Q13TZ4
D	-13	HIS	-	EXPRESSION TAG	UNP Q13TZ4
D	-12	MET	-	EXPRESSION TAG	UNP Q13TZ4
D	-11	GLY	-	EXPRESSION TAG	UNP Q13TZ4
D	-10	THR	-	EXPRESSION TAG	UNP Q13TZ4
D	-9	LEU	-	EXPRESSION TAG	UNP Q13TZ4
D	-8	GLU	-	EXPRESSION TAG	UNP Q13TZ4
D	-7	ALA	-	EXPRESSION TAG	UNP Q13TZ4
D	-6	GLN	-	EXPRESSION TAG	UNP Q13TZ4
D	-5	THR	-	EXPRESSION TAG	UNP Q13TZ4
D	-4	GLN	-	EXPRESSION TAG	UNP Q13TZ4
D	-3	GLY	-	EXPRESSION TAG	UNP Q13TZ4
D	-2	PRO	-	EXPRESSION TAG	UNP Q13TZ4
D	-1	GLY	-	EXPRESSION TAG	UNP Q13TZ4
D	0	SER	-	EXPRESSION TAG	UNP Q13TZ4


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	113	Total	O	0	0
			113	113		
2	B	98	Total	O	0	0
			98	98		
2	C	69	Total	O	0	0
			69	69		
2	D	78	Total	O	0	0
			78	78		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: D-alanine–D-alanine ligase




- Molecule 1: D-alanine–D-alanine ligase



- Molecule 1: D-alanine–D-alanine ligase



- Molecule 1: D-alanine–D-alanine ligase



4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	78.93Å 78.93Å 224.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.99 – 2.30 44.76 – 2.30	Depositor EDS
% Data completeness (in resolution range)	97.8 (49.99-2.30) 97.8 (44.76-2.30)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.91 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.224 , 0.260 0.222 , 0.259	Depositor DCC
R_{free} test set	3006 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	29.6	Xtriage
Anisotropy	0.429	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 40.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.069 for h,-k,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8430	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 61.77 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.2460e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	1/2167 (0.0%)	0.86	1/2948 (0.0%)
1	B	0.78	0/2144	0.87	3/2917 (0.1%)
1	C	0.78	0/2071	0.90	5/2820 (0.2%)
1	D	0.84	1/1858 (0.1%)	0.90	2/2525 (0.1%)
All	All	0.79	2/8240 (0.0%)	0.88	11/11210 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	258	TRP	CD2-CE2	5.17	1.47	1.41
1	D	111	TRP	CD2-CE2	5.14	1.47	1.41

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	31	ARG	NE-CZ-NH1	-7.43	116.58	120.30
1	C	31	ARG	NE-CZ-NH2	7.04	123.82	120.30
1	D	106	ARG	NE-CZ-NH2	-6.33	117.14	120.30
1	B	301	GLU	OE1-CD-OE2	-6.22	115.84	123.30
1	C	229	LEU	CB-CG-CD2	-5.90	100.97	111.00
1	A	106	ARG	NE-CZ-NH2	-5.42	117.59	120.30
1	C	241	ARG	CB-CA-C	-5.36	99.69	110.40
1	D	31	ARG	NE-CZ-NH1	5.34	122.97	120.30
1	B	89	ARG	NE-CZ-NH2	-5.18	117.71	120.30
1	C	106	ARG	NE-CZ-NH2	-5.16	117.72	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	247	ARG	NE-CZ-NH1	-5.13	117.73	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	256	THR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2118	0	2109	8	0
1	B	2099	0	2079	13	0
1	C	2033	0	1925	3	0
1	D	1822	0	1781	4	0
2	A	113	0	0	2	0
2	B	98	0	0	1	0
2	C	69	0	0	0	0
2	D	78	0	0	1	0
All	All	8430	0	7894	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:175:VAL:O	1:A:175:VAL:HG12	1.93	0.69
1:D:145:LEU:HD13	1:D:183:VAL:CG1	2.23	0.68
1:A:175:VAL:O	1:A:175:VAL:CG1	2.47	0.63
1:B:143:LEU:HD13	1:B:162:LYS:O	1.99	0.62
1:B:228:TYR:O	1:B:290:LYS:NZ	2.37	0.58
1:B:45:HIS:ND1	2:B:417:HOH:O	2.34	0.54
1:C:145:LEU:HD13	1:C:183:VAL:CG1	2.38	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:264:MET:HG2	1:C:274:LEU:HD11	1.90	0.52
1:B:175:VAL:HG12	1:B:175:VAL:O	2.10	0.52
1:B:175:VAL:CG1	1:B:175:VAL:O	2.59	0.51
1:B:169:ALA:O	1:B:172:ILE:HG22	2.14	0.48
1:B:126:ARG:HB2	1:B:175:VAL:HG13	1.96	0.47
1:A:126:ARG:HB2	1:A:175:VAL:HG13	1.97	0.47
1:D:145:LEU:HD13	1:D:183:VAL:HG12	1.97	0.47
1:D:199:GLY:HA2	2:D:407:HOH:O	2.16	0.46
1:A:169:ALA:HB3	2:A:489:HOH:O	2.16	0.45
1:B:134:ALA:O	1:B:138:VAL:HG23	2.17	0.45
1:D:205:VAL:CG2	1:D:239:GLU:HG3	2.47	0.44
1:A:22:GLU:HA	2:A:475:HOH:O	2.17	0.44
1:B:5:ASP:HA	1:B:6:PRO:HD2	1.89	0.44
1:B:102:LEU:HA	1:B:102:LEU:HD12	1.87	0.43
1:A:194:THR:HG21	1:A:287:LEU:HD13	2.02	0.42
1:A:126:ARG:HA	1:A:175:VAL:HG11	2.02	0.42
1:B:228:TYR:CD2	1:B:287:LEU:HD23	2.56	0.41
1:B:54:LEU:HD23	1:B:82:ALA:HB1	2.02	0.41
1:A:300:GLN:O	1:A:303:VAL:HG22	2.21	0.41
1:B:96:LEU:HD11	1:B:100:LEU:HD22	2.02	0.41
1:C:102:LEU:HA	1:C:102:LEU:HD12	1.97	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	282/334 (84%)	277 (98%)	5 (2%)	0	100	100
1	B	281/334 (84%)	276 (98%)	5 (2%)	0	100	100
1	C	275/334 (82%)	268 (98%)	7 (2%)	0	100	100
1	D	238/334 (71%)	234 (98%)	4 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	1076/1336 (80%)	1055 (98%)	21 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	210/258 (81%)	205 (98%)	5 (2%)	49	66
1	B	206/258 (80%)	201 (98%)	5 (2%)	49	66
1	C	189/258 (73%)	185 (98%)	4 (2%)	53	70
1	D	181/258 (70%)	178 (98%)	3 (2%)	60	76
All	All	786/1032 (76%)	769 (98%)	17 (2%)	52	69

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	GLN
1	A	16	LEU
1	A	73	TYR
1	A	125	LEU
1	A	241	ARG
1	B	8	GLN
1	B	16	LEU
1	B	54	LEU
1	B	172	ILE
1	B	241	ARG
1	C	16	LEU
1	C	24	GLU
1	C	73	TYR
1	C	284	ASP
1	D	16	LEU
1	D	59	GLU
1	D	284	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	285/334 (85%)	0.15	9 (3%) 47 54	19, 27, 55, 72	0
1	B	285/334 (85%)	0.13	8 (2%) 53 60	18, 26, 63, 82	0
1	C	283/334 (84%)	0.84	41 (14%) 2 3	21, 31, 91, 112	0
1	D	246/334 (73%)	0.07	10 (4%) 37 44	18, 25, 61, 82	0
All	All	1099/1336 (82%)	0.30	68 (6%) 20 26	18, 27, 69, 112	0

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	170	ALA	13.4
1	C	166	ALA	10.0
1	C	177	PHE	9.8
1	C	172	ILE	9.7
1	C	174	ALA	8.7
1	C	175	VAL	7.8
1	C	169	ALA	7.4
1	C	173	GLU	6.8
1	C	165	ASP	6.7
1	C	129	ASP	6.6
1	C	127	GLY	6.5
1	C	164	ALA	6.4
1	C	124	VAL	6.3
1	C	132	ALA	6.0
1	C	171	LEU	6.0
1	C	131	GLU	5.7
1	D	147	VAL	5.4
1	C	163	SER	5.4
1	C	167	LEU	5.4
1	D	182	VAL	5.1
1	C	125	LEU	5.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	183	VAL	4.9
1	C	176	LYS	4.8
1	C	128	ASP	4.7
1	C	183	VAL	4.6
1	C	179	ARG	4.6
1	C	133	ARG	4.6
1	C	178	ASP	4.6
1	C	123	ALA	4.5
1	C	168	PRO	4.4
1	A	137	ILE	4.3
1	C	130	TYR	4.1
1	D	146	PHE	3.9
1	C	18	GLY	3.9
1	A	174	ALA	3.8
1	C	21	ALA	3.7
1	C	182	VAL	3.7
1	D	143	LEU	3.6
1	C	19	ASN	3.6
1	C	159	ILE	3.6
1	A	177	PHE	3.4
1	B	130	TYR	3.4
1	B	4	ILE	3.4
1	B	132	ALA	3.3
1	C	134	ALA	3.2
1	B	175	VAL	3.1
1	A	171	LEU	3.1
1	C	180	ILE	3.0
1	A	170	ALA	3.0
1	C	146	PHE	3.0
1	C	55	ALA	2.9
1	A	130	TYR	2.9
1	D	121	PHE	2.9
1	C	86	TYR	2.9
1	C	59	GLU	2.8
1	D	145	LEU	2.8
1	C	121	PHE	2.7
1	B	138	VAL	2.6
1	D	181	VAL	2.5
1	A	133	ARG	2.5
1	B	166	ALA	2.5
1	A	128	ASP	2.3
1	B	172	ILE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	127	GLY	2.2
1	A	5	ASP	2.2
1	C	147	VAL	2.1
1	D	3	SER	2.1
1	D	120	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.