



Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 05:18 pm BST

PDB ID : 5EG9
Title : The cap binding site of influenza virus protein PB2 as a drug target
Authors : Severin, C.; Rocha de Moura, T.; Liu, Y.; Li, K.; Zheng, X.; Luo, M.
Deposited on : 2015-10-26
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	183	<div> <div>3%</div> <div> <div></div> <div>71%</div> <div>11%</div> <div>•</div> <div>14%</div> </div> </div>
1	B	183	<div> <div>2%</div> <div> <div></div> <div>77%</div> <div>6%</div> <div>•</div> <div>15%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2522 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polymerase basic protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	157	Total	C	N	O	S	0	1	0
			1237	778	222	228	9			
1	B	156	Total	C	N	O	S	0	0	0
			1222	769	219	225	9			

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	297	MET	-	initiating methionine	UNP Q6DNL8
A	298	GLY	-	expression tag	UNP Q6DNL8
A	299	SER	-	expression tag	UNP Q6DNL8
A	300	SER	-	expression tag	UNP Q6DNL8
A	301	HIS	-	expression tag	UNP Q6DNL8
A	302	HIS	-	expression tag	UNP Q6DNL8
A	303	HIS	-	expression tag	UNP Q6DNL8
A	304	HIS	-	expression tag	UNP Q6DNL8
A	305	HIS	-	expression tag	UNP Q6DNL8
A	306	HIS	-	expression tag	UNP Q6DNL8
A	307	SER	-	expression tag	UNP Q6DNL8
A	308	SER	-	expression tag	UNP Q6DNL8
A	309	GLY	-	expression tag	UNP Q6DNL8
A	310	LEU	-	expression tag	UNP Q6DNL8
A	311	VAL	-	expression tag	UNP Q6DNL8
A	312	PRO	-	expression tag	UNP Q6DNL8
A	313	ARG	-	expression tag	UNP Q6DNL8
A	314	GLY	-	expression tag	UNP Q6DNL8
A	315	SER	-	expression tag	UNP Q6DNL8
A	316	HIS	-	expression tag	UNP Q6DNL8
A	317	MET	-	expression tag	UNP Q6DNL8
A	421	GLY	-	linker	UNP Q6DNL8
A	422	SER	-	linker	UNP Q6DNL8
A	423	GLY	-	linker	UNP Q6DNL8
A	449	SER	PRO	conflict	UNP Q6DNL8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	474	ILE	VAL	conflict	UNP Q6DNL8
B	297	MET	-	initiating methionine	UNP Q6DNL8
B	298	GLY	-	expression tag	UNP Q6DNL8
B	299	SER	-	expression tag	UNP Q6DNL8
B	300	SER	-	expression tag	UNP Q6DNL8
B	301	HIS	-	expression tag	UNP Q6DNL8
B	302	HIS	-	expression tag	UNP Q6DNL8
B	303	HIS	-	expression tag	UNP Q6DNL8
B	304	HIS	-	expression tag	UNP Q6DNL8
B	305	HIS	-	expression tag	UNP Q6DNL8
B	306	HIS	-	expression tag	UNP Q6DNL8
B	307	SER	-	expression tag	UNP Q6DNL8
B	308	SER	-	expression tag	UNP Q6DNL8
B	309	GLY	-	expression tag	UNP Q6DNL8
B	310	LEU	-	expression tag	UNP Q6DNL8
B	311	VAL	-	expression tag	UNP Q6DNL8
B	312	PRO	-	expression tag	UNP Q6DNL8
B	313	ARG	-	expression tag	UNP Q6DNL8
B	314	GLY	-	expression tag	UNP Q6DNL8
B	315	SER	-	expression tag	UNP Q6DNL8
B	316	HIS	-	expression tag	UNP Q6DNL8
B	317	MET	-	expression tag	UNP Q6DNL8
B	421	GLY	-	linker	UNP Q6DNL8
B	422	SER	-	linker	UNP Q6DNL8
B	423	GLY	-	linker	UNP Q6DNL8
B	449	SER	PRO	conflict	UNP Q6DNL8
B	474	ILE	VAL	conflict	UNP Q6DNL8


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	35	Total O 35 35	0	0
2	B	28	Total O 28 28	0	0



3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Polymerase basic protein 2



- Molecule 1: Polymerase basic protein 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	37.13Å 109.01Å 39.66Å 90.00° 90.54° 90.00°	Depositor
Resolution (Å)	20.00 – 2.30 27.23 – 2.30	Depositor EDS
% Data completeness (in resolution range)	87.2 (20.00-2.30) 87.3 (27.23-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.24 (at 2.31Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.191 , 0.258 0.199 , 0.260	Depositor DCC
R_{free} test set	602 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	29.4	Xtriage
Anisotropy	0.098	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 36.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.166 for h,-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2522	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	0/1254	0.91	2/1679 (0.1%)
1	B	0.67	0/1239	0.82	0/1660
All	All	0.69	0/2493	0.87	2/3339 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	375	ARG	NE-CZ-NH2	-11.33	114.64	120.30
1	A	375	ARG	NE-CZ-NH1	10.58	125.59	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1237	0	1266	13	0
1	B	1222	0	1249	8	0
2	A	35	0	0	0	0
2	B	28	0	0	3	0
All	All	2522	0	2515	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:371:THR:HG23	2:B:526:HOH:O	1.79	0.83
1:A:437:ASP:OD1	1:A:439:LYS:HD2	1.81	0.78
1:B:422:SER:CB	1:B:423:GLY:HA3	2.22	0.70
1:A:422:SER:CB	1:A:423:GLY:HA3	2.31	0.61
1:B:422:SER:HB2	1:B:423:GLY:HA3	1.84	0.60
1:A:454:MET:SD	1:A:454:MET:N	2.74	0.59
1:B:371:THR:CG2	2:B:526:HOH:O	2.46	0.57
1:A:449:SER:HA	1:A:469:MET:HG2	1.90	0.53
1:A:423:GLY:O	1:A:424:LEU:HD23	2.10	0.52
1:B:366:VAL:HG13	2:B:516:HOH:O	2.10	0.51
1:A:422:SER:HB2	1:A:423:GLY:HA3	1.93	0.51
1:A:425:ASN:HB2	1:A:426:PRO:CD	2.41	0.50
1:A:421:GLY:HA2	1:A:424:LEU:HB2	1.93	0.49
1:A:419:ASN:O	1:A:429:GLN:NE2	2.46	0.48
1:B:342:GLU:O	1:B:353:LYS:HA	2.14	0.48
1:A:420:PHE:CG	1:A:421:GLY:HA3	2.50	0.46
1:A:387:SER:OG	1:A:477[B]:SER:OG	2.28	0.44
1:A:375:ARG:HD3	1:A:383:GLN:OE1	2.17	0.44
1:A:380:ARG:HD2	1:A:382:ILE:HD11	2.00	0.42
1:B:450:ILE:HG22	1:B:475:ARG:NH2	2.34	0.42
1:B:332:ARG:CZ	1:B:336:SER:HA	2.50	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	156/183 (85%)	148 (95%)	7 (4%)	1 (1%)	25	31
1	B	154/183 (84%)	148 (96%)	5 (3%)	1 (1%)	25	31
All	All	310/366 (85%)	296 (96%)	12 (4%)	2 (1%)	25	31

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	422	SER
1	A	419	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	136/158 (86%)	125 (92%)	11 (8%)	11	15
1	B	134/158 (85%)	126 (94%)	8 (6%)	19	26
All	All	270/316 (85%)	251 (93%)	19 (7%)	14	19

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	336	SER
1	A	376	LYS
1	A	386	VAL
1	A	389	ARG
1	A	420	PHE
1	A	422	SER
1	A	431	LEU
1	A	436	LYS
1	A	439	LYS
1	A	447	ILE
1	A	454	MET
1	B	336	SER
1	B	366	VAL
1	B	371	THR
1	B	376	LYS
1	B	386	VAL
1	B	436	LYS
1	B	439	LYS
1	B	447	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	443	GLN
1	B	357	HIS
1	B	429	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	157/183 (85%)	-0.12	5 (3%) 47 54	16, 30, 58, 99	0
1	B	156/183 (85%)	-0.13	3 (1%) 66 73	15, 32, 60, 98	0
All	All	313/366 (85%)	-0.13	8 (2%) 56 63	15, 31, 59, 99	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	421	GLY	4.7
1	B	420	PHE	4.7
1	A	420	PHE	4.5
1	A	422	SER	4.2
1	A	421	GLY	3.9
1	B	422	SER	3.6
1	A	423	GLY	2.7
1	A	454	MET	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.