



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 6, 2020 – 10:48 AM BST

PDB ID : 5EHN
Title : mAChE-syn TZ2PA5 complex
Authors : Bourne, Y.; Marchot, P.
Deposited on : 2015-10-28
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	543	
1	B	543	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8683 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetylcholinesterase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	536	Total	C	N	O	S	0	2	0
			4190	2689	727	760	14			
1	B	531	Total	C	N	O	S	0	1	0
			4148	2666	715	753	14			

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is 6-phenyl-5-[5-[3-[2-(1,2,3,4-tetrahydroacridin-9-ylamino)ethyl]-1,2,3-triazol-4-yl]pentyl]phenanthridin-5-ium-3,8-diamine (three-letter code: 5NZ) (formula: $C_{41}H_{43}N_8$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 49	C 41	N 8	0	0
3	B	1	Total 49	C 41	N 8	0	0

- Molecule 4 is HEXAETHYLENE GLYCOL (three-letter code: P6G) (formula: $\text{C}_{12}\text{H}_{26}\text{O}_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			19	12	7		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	126	Total 126	O 126	0	0
5	B	88	Total 88	O 88	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Acetylcholinesterase



• Molecule 1: Acetylcholinesterase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	79.75Å 112.45Å 227.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.00 – 2.60 45.95 – 2.69	Depositor EDS
% Data completeness (in resolution range)	98.5 (45.00-2.60) 98.1 (45.95-2.69)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.24 (at 2.69Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.182 , 0.203 0.234 , 0.224	Depositor DCC
R_{free} test set	1153 reflections (2.04%)	wwPDB-VP
Wilson B-factor (Å ²)	45.1	Xtriage
Anisotropy	0.784	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 49.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	8683	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 5NZ, NAG, P6G

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	0/4320	0.70	0/5903
1	B	0.50	0/4275	0.68	0/5846
All	All	0.50	0/8595	0.69	0/11749

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4190	0	4076	23	0
1	B	4148	0	4029	27	0
2	A	14	0	13	0	0
3	A	49	0	43	3	0
3	B	49	0	43	1	0
4	B	19	0	26	3	0
5	A	126	0	0	2	0
5	B	88	0	0	0	0
All	All	8683	0	8230	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:161:LEU:HD11	1:B:269:LEU:HD22	1.59	0.83
1:B:197:VAL:H	1:B:223:HIS:HD2	1.36	0.73
1:A:197:VAL:H	1:A:223:HIS:HD2	1.36	0.71
1:A:537:PRO:O	1:A:540:LEU:HG	1.92	0.69
1:B:211:MET:HG2	1:B:308:LEU:HD21	1.79	0.65
1:B:224:ARG:HG2	1:B:325:GLN:HB2	1.78	0.64
1:A:337:TYR:O	1:A:340:VAL:HG22	1.99	0.63
1:A:211:MET:HG2	1:A:308:LEU:HD21	1.80	0.63
1:B:197:VAL:H	1:B:223:HIS:CD2	2.19	0.60
1:A:197:VAL:H	1:A:223:HIS:CD2	2.18	0.59
1:A:296:ARG:HH21	1:A:369[B]:GLN:HE22	1.52	0.58
1:B:326:VAL:HG12	1:B:328:VAL:HG13	1.85	0.58
1:A:68:VAL:HG11	1:A:88:PRO:HB3	1.86	0.56
1:A:369[B]:GLN:H	1:A:369[B]:GLN:HE21	1.52	0.56
1:B:161:LEU:HD12	1:B:270:ILE:HD11	1.88	0.55
1:A:68:VAL:HG23	1:A:90:ARG:HB2	1.91	0.53
1:A:296:ARG:HH21	1:A:369[B]:GLN:NE2	2.07	0.53
1:B:68:VAL:HG11	1:B:88:PRO:HB3	1.89	0.53
1:B:68:VAL:HG23	1:B:90:ARG:HB2	1.92	0.52
1:B:166:GLU:HG2	1:B:270:ILE:HD13	1.91	0.51
1:A:376:GLU:O	1:A:380:LEU:HG	2.10	0.51
1:A:210:GLY:HA3	1:A:232:PRO:HD3	1.92	0.51
1:B:376:GLU:O	1:B:380:LEU:HG	2.11	0.50
1:A:380:LEU:HB2	4:B:1901:P6G:H142	1.93	0.50
3:A:1502:5NZ:H29	5:A:1622:HOH:O	2.11	0.49
1:B:210:GLY:HA3	1:B:232:PRO:HD3	1.93	0.49
1:B:381:HIS:HA	4:B:1901:P6G:H51	1.93	0.49
1:B:380:LEU:HB2	4:B:1901:P6G:H81	1.94	0.49
1:B:104:PRO:HD2	1:B:108:PRO:HD3	1.95	0.48
1:B:528:THR:O	1:B:531:PHE:HB3	2.14	0.48
1:B:328:VAL:O	1:B:427:ALA:HA	2.14	0.47
1:A:104:PRO:HD2	1:A:108:PRO:HD3	1.96	0.47
1:A:224:ARG:HG2	1:A:325:GLN:HB2	1.96	0.47
1:A:528:THR:O	1:A:531:PHE:HB3	2.15	0.47
1:A:369[B]:GLN:NE2	1:A:369[B]:GLN:H	2.13	0.46
1:A:74:ASP:CG	3:A:1502:5NZ:H8	2.35	0.46
1:B:340:VAL:HG11	1:B:443:MET:HE1	1.97	0.46
1:A:328:VAL:O	1:A:427:ALA:HA	2.16	0.46
1:A:374:ALA:HA	1:A:539:LEU:HD23	1.98	0.45
1:B:161:LEU:HD12	1:B:270:ILE:CD1	2.47	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:227:LEU:HB2	1:B:328:VAL:HG12	1.98	0.45
1:A:293:SER:HB3	1:A:368:PRO:HG3	1.99	0.45
1:A:227:LEU:HB2	1:A:328:VAL:HG12	2.00	0.43
1:B:103:THR:HG21	1:B:190:PHE:HB3	2.01	0.43
1:B:337:TYR:O	1:B:340:VAL:HG22	2.19	0.43
1:A:103:THR:HG21	1:A:190:PHE:HB3	2.01	0.42
1:B:257:CYS:HA	1:B:258:PRO:HA	1.85	0.42
1:B:293:SER:HB3	1:B:368:PRO:HG3	2.01	0.42
1:B:525:ARG:HG2	1:B:528:THR:HB	2.01	0.41
3:B:1902:5NZ:H30	3:B:1902:5NZ:C40	2.50	0.41
3:A:1502:5NZ:C29	5:A:1622:HOH:O	2.67	0.41
1:B:294:ILE:HG12	1:B:365:ILE:HG22	2.03	0.40
1:B:340:VAL:HG11	1:B:443:MET:CE	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	534/543 (98%)	522 (98%)	12 (2%)	0	100	100
1	B	528/543 (97%)	517 (98%)	11 (2%)	0	100	100
All	All	1062/1086 (98%)	1039 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	440/443 (99%)	434 (99%)	6 (1%)	67	85
1	B	436/443 (98%)	432 (99%)	4 (1%)	78	91
All	All	876/886 (99%)	866 (99%)	10 (1%)	73	88

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	132	VAL
1	A	146	LEU
1	A	161	LEU
1	A	181	GLN
1	A	220	SER
1	A	295	PHE
1	B	146	LEU
1	B	181	GLN
1	B	295	PHE
1	B	536	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	223	HIS
1	A	405	HIS
1	B	223	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	5NZ	B	1902	-	53,56,56	2.25	12 (22%)	66,79,79	1.66	12 (18%)
2	NAG	A	1501	1	14,14,15	0.40	0	17,19,21	0.83	1 (5%)
4	P6G	B	1901	-	18,18,18	2.31	6 (33%)	17,17,17	1.51	3 (17%)
3	5NZ	A	1502	-	53,56,56	2.19	12 (22%)	66,79,79	1.60	10 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	5NZ	B	1902	-	-	5/18/25/25	0/8/8/8
2	NAG	A	1501	1	-	2/6/23/26	0/1/1/1
4	P6G	B	1901	-	-	6/16/16/16	-
3	5NZ	A	1502	-	-	6/18/25/25	0/8/8/8

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1502	5NZ	C39-C37	7.36	1.51	1.40
3	B	1902	5NZ	C30-C39	7.31	1.51	1.38
3	B	1902	5NZ	C39-C37	7.00	1.51	1.40
3	A	1502	5NZ	C30-C39	6.76	1.50	1.38
3	B	1902	5NZ	C5-C18	5.23	1.51	1.41
3	A	1502	5NZ	C5-C18	4.90	1.51	1.41
4	B	1901	P6G	O10-C9	4.62	1.62	1.42
3	B	1902	5NZ	C10-C4	4.62	1.49	1.42
3	A	1502	5NZ	C10-C4	4.60	1.49	1.42
3	B	1902	5NZ	C31-C33	4.25	1.49	1.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1502	5NZ	C11-C10	4.18	1.50	1.43
4	B	1901	P6G	O13-C12	4.05	1.59	1.42
3	B	1902	5NZ	C11-C10	3.99	1.50	1.43
3	A	1502	5NZ	C31-C33	3.87	1.48	1.42
4	B	1901	P6G	O16-C15	3.78	1.58	1.42
3	A	1502	5NZ	N6-N5	3.67	1.40	1.34
4	B	1901	P6G	O19-C18	3.65	1.60	1.42
3	B	1902	5NZ	N6-N5	3.59	1.40	1.34
4	B	1901	P6G	O4-C3	3.49	1.57	1.42
3	A	1502	5NZ	C30-C31	3.28	1.49	1.43
3	B	1902	5NZ	C30-C31	3.24	1.49	1.43
3	B	1902	5NZ	N5-N4	3.12	1.40	1.34
3	A	1502	5NZ	C37-N8	3.05	1.36	1.32
3	B	1902	5NZ	C37-N8	3.02	1.36	1.32
4	B	1901	P6G	O7-C6	2.91	1.54	1.42
3	B	1902	5NZ	C18-N3	-2.69	1.36	1.40
3	A	1502	5NZ	N5-N4	2.30	1.39	1.34
3	B	1902	5NZ	C26-N4	-2.29	1.34	1.37
3	A	1502	5NZ	C26-N4	-2.22	1.34	1.37
3	A	1502	5NZ	C18-N3	-2.14	1.37	1.40

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1902	5NZ	C37-N8-C33	6.14	125.24	117.67
3	A	1502	5NZ	C37-N8-C33	5.40	124.34	117.67
3	A	1502	5NZ	C39-C30-N7	4.60	127.97	119.54
3	B	1902	5NZ	C39-C37-N8	-4.39	119.93	123.68
3	A	1502	5NZ	C39-C37-N8	-3.91	120.34	123.68
3	B	1902	5NZ	C39-C30-N7	3.65	126.22	119.54
3	A	1502	5NZ	C19-C18-C5	-3.49	116.18	120.61
4	B	1901	P6G	O1-C2-C3	3.37	131.38	111.81
3	B	1902	5NZ	C26-N4-N5	-3.04	110.31	113.04
3	B	1902	5NZ	C19-C18-C5	-2.97	116.85	120.61
3	A	1502	5NZ	C26-N4-N5	-2.91	110.43	113.04
3	B	1902	5NZ	C28-N4-C26	2.79	132.04	128.48
3	A	1502	5NZ	C28-N4-C26	2.74	131.97	128.48
4	B	1901	P6G	O13-C12-C11	2.58	122.04	110.39
2	A	1501	NAG	C1-O5-C5	2.45	115.51	112.19
3	B	1902	5NZ	C31-C33-N8	-2.44	120.22	122.81
3	B	1902	5NZ	C13-C12-C11	-2.35	116.77	120.24
3	B	1902	5NZ	C40-C39-C37	-2.32	118.91	121.08

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1502	5NZ	C31-C30-N7	-2.30	114.53	122.06
3	A	1502	5NZ	C40-C39-C37	-2.27	118.96	121.08
3	B	1902	5NZ	C10-C4-C5	-2.26	116.95	119.91
3	A	1502	5NZ	C32-C31-C30	-2.22	120.78	124.78
4	B	1901	P6G	O7-C6-C5	2.13	119.98	110.39
3	B	1902	5NZ	C17-C12-C13	2.02	121.62	117.59
3	B	1902	5NZ	C25-C26-N4	2.01	126.88	122.34
3	A	1502	5NZ	C10-C4-C5	-2.00	117.29	119.91

There are no chirality outliers.

All (19) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	1902	5NZ	C29-C28-N4-C26
3	B	1902	5NZ	N4-C28-C29-N7
3	A	1502	5NZ	C29-C28-N4-C26
3	A	1502	5NZ	N4-C28-C29-N7
4	B	1901	P6G	O1-C2-C3-O4
4	B	1901	P6G	O13-C14-C15-O16
4	B	1901	P6G	O16-C17-C18-O19
2	A	1501	NAG	C4-C5-C6-O6
3	A	1502	5NZ	C39-C30-N7-C29
3	A	1502	5NZ	C31-C30-N7-C29
4	B	1901	P6G	C11-C12-O13-C14
3	B	1902	5NZ	C39-C30-N7-C29
4	B	1901	P6G	O7-C8-C9-O10
3	B	1902	5NZ	C31-C30-N7-C29
2	A	1501	NAG	O5-C5-C6-O6
3	A	1502	5NZ	C20-C21-C22-C24
3	B	1902	5NZ	C29-C28-N4-N5
3	A	1502	5NZ	C29-C28-N4-N5
4	B	1901	P6G	C6-C5-O4-C3


There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1902	5NZ	1	0
4	B	1901	P6G	3	0
3	A	1502	5NZ	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.


6.3 Carbohydrates ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands ⓘ




Unable to reproduce the depositors R factor - this section is therefore empty.


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around P6G B 1901:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.