



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 21, 2020 – 07:14 PM BST

PDB ID : 5EHP
Title : Non-receptor Protein Tyrosine Phosphatase SHP2 in Complex with Allosteric Inhibitor SHP836
Authors : Stams, T.; Fodor, M.
Deposited on : 2015-10-28
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	526	<div> <div>6%</div> <div> <div></div> <div>89%</div> <div>5%</div> <div>6%</div> </div> </div>
1	B	526	<div> <div>4%</div> <div> <div></div> <div>83%</div> <div>6%</div> <div>10%</div> </div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8600 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein phosphatase non-receptor type 11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	494	Total	C	N	O	S	0	4	0
			4020	2539	713	749	19			
1	B	471	Total	C	N	O	S	0	3	0
			3840	2426	679	715	20			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP Q06124
B	0	SER	-	expression tag	UNP Q06124

- Molecule 2 is 5-[2,3-bis(chloranyl)phenyl]-2-[(3 {R},5 {S})-3,5-dimethylpiperazin-1-yl]pyrimidin-4-amine (three-letter code: 5OA) (formula: C₁₆H₁₉Cl₂N₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	Cl	N	0	0
			23	16	2	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	Cl	N	0	0
			23	16	2	5		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	349	Total 350	O 350	0	1
4	B	304	Total 304	O 304	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein phosphatase non-receptor type 11



- Molecule 1: Tyrosine-protein phosphatase non-receptor type 11



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	46.31 Å 213.73 Å 55.98 Å 90.00° 96.82° 90.00°	Depositor
Resolution (Å)	27.09 – 1.85 26.90 – 1.85	Depositor EDS
% Data completeness (in resolution range)	95.5 (27.09-1.85) 95.9 (26.90-1.85)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.95 (at 1.85 Å)	Xtriage
Refinement program	BUSTER BUSTER 2.11.5	Depositor
R, R_{free}	0.174 , 0.208 0.179 , 0.211	Depositor DCC
R_{free} test set	4388 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	26.0	Xtriage
Anisotropy	0.274	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8600	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, 5OA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/4113	0.61	0/5538
1	B	0.51	0/3923	0.62	0/5285
All	All	0.51	0/8036	0.62	0/10823

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4020	0	3972	9	0
1	B	3840	0	3805	9	0
2	A	23	0	0	0	0
2	B	23	0	0	0	0
3	A	25	0	0	0	0
3	B	15	0	0	0	0
4	A	350	0	0	0	0
4	B	304	0	0	0	0
All	All	8600	0	7777	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:132:HIS:HD1	1:A:154:GLY:H	1.48	0.61
1:B:133:GLY:HA3	1:B:213:LYS:HB2	1.87	0.56
1:A:125:LEU:HB3	1:A:216:LEU:HD21	1.88	0.55
1:B:125:LEU:HB3	1:B:216:LEU:HD21	1.89	0.55
1:A:198:LYS:HG3	1:A:212:LEU:HB2	1.93	0.50
1:B:290:VAL:HG11	1:B:344:MET:HG3	1.95	0.49
1:B:222:ASN:O	1:B:225:GLU:HG2	2.13	0.48
1:A:232:GLU:HA	1:A:235:LYS:HE2	1.94	0.48
1:A:90:GLU:HG3	1:A:96:ILE:HD11	1.97	0.46
1:A:309:ILE:HD13	1:A:328:ILE:HG12	1.99	0.45
1:A:129:LYS:HB2	1:A:216:LEU:HD11	1.99	0.44
1:B:432:PRO:HG3	1:B:516[A]:MET:HG2	2.00	0.43
1:B:177:LEU:HD13	1:B:177:LEU:HA	1.86	0.42
1:B:149:LEU:HD21	1:B:212:LEU:HD21	1.99	0.42
1:B:129:LYS:HB2	1:B:216:LEU:HD11	2.02	0.41
1:B:309:ILE:HD13	1:B:328:ILE:HG12	2.02	0.41
1:A:19:LEU:HD23	1:A:23:ARG:HD2	2.04	0.40
1:A:19:LEU:HD22	1:A:102:LEU:HD21	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	480/526 (91%)	471 (98%)	8 (2%)	1 (0%)	47	33
1	B	458/526 (87%)	448 (98%)	9 (2%)	1 (0%)	47	33
All	All	938/1052 (89%)	919 (98%)	17 (2%)	2 (0%)	47	33

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	505	VAL
1	A	505	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	443/468 (95%)	431 (97%)	12 (3%)	44	29
1	B	424/468 (91%)	406 (96%)	18 (4%)	30	13
All	All	867/936 (93%)	837 (96%)	30 (4%)	37	18

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	SER
1	A	47	ARG
1	A	94	ASP
1	A	198	LYS
1	A	294	ASP
1	A	298	ASN
1	A	408	GLN
1	A	413	ARG
1	A	481	GLU
1	A	484	VAL
1	A	508	GLU
1	A	523	GLU
1	B	11	ILE
1	B	18[A]	ASN
1	B	18[B]	ASN
1	B	47	ARG
1	B	89	LYS
1	B	120	LYS
1	B	131	LYS
1	B	152	ARG
1	B	177	LEU
1	B	211	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	233	LEU
1	B	245	GLN
1	B	275	ASN
1	B	325	LYS
1	B	413	ARG
1	B	484	VAL
1	B	508	GLU
1	B	525	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	211	GLN
1	A	256	GLN
1	A	298	ASN
1	B	37	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PO4	A	605	-	4,4,4	2.48	1 (25%)	6,6,6	0.70	0
2	5OA	B	601	-	25,25,25	0.77	0	34,36,36	2.42	12 (35%)
3	PO4	B	604	-	4,4,4	2.47	1 (25%)	6,6,6	0.51	0
3	PO4	A	606	-	4,4,4	1.97	1 (25%)	6,6,6	0.74	0
3	PO4	A	602	-	4,4,4	2.41	1 (25%)	6,6,6	0.88	0
3	PO4	A	604	-	4,4,4	2.06	1 (25%)	6,6,6	0.39	0
2	5OA	A	601	-	25,25,25	0.84	0	34,36,36	2.63	12 (35%)
3	PO4	A	603	-	4,4,4	2.52	2 (50%)	6,6,6	0.45	0
3	PO4	B	603	-	4,4,4	2.49	1 (25%)	6,6,6	0.48	0
3	PO4	B	602	-	4,4,4	2.38	1 (25%)	6,6,6	0.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	5OA	B	601	-	-	0/8/20/20	0/3/3/3
2	5OA	A	601	-	-	0/8/20/20	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	603	PO4	P-O1	4.20	1.60	1.50
3	A	605	PO4	P-O1	4.15	1.60	1.50
3	B	603	PO4	P-O1	4.12	1.60	1.50
3	B	604	PO4	P-O1	4.09	1.60	1.50
3	B	602	PO4	P-O1	3.98	1.60	1.50
3	A	602	PO4	P-O1	3.91	1.60	1.50
3	A	604	PO4	P-O1	2.35	1.56	1.50
3	A	603	PO4	P-O2	2.00	1.60	1.54
3	A	606	PO4	P-O1	2.00	1.55	1.50

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	5OA	C14-N15-C16	8.58	123.25	115.64
2	A	601	5OA	N15-C16-N17	-8.29	117.71	126.00
2	B	601	5OA	N15-C16-N17	-7.39	118.61	126.00
2	B	601	5OA	C14-N15-C16	7.25	122.07	115.64

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	5OA	N17-C16-N1	3.74	121.72	117.11
2	A	601	5OA	C13-C12-C7	-3.65	118.78	122.86
2	B	601	5OA	N15-C16-N1	3.52	120.71	116.90
2	A	601	5OA	N15-C16-N1	3.39	120.57	116.90
2	B	601	5OA	C12-C7-C8	3.11	122.34	120.19
2	A	601	5OA	C13-C14-N15	-3.07	119.66	124.49
2	B	601	5OA	C5-N4-C3	3.04	116.80	111.75
2	B	601	5OA	N17-C16-N1	2.84	120.62	117.11
2	A	601	5OA	C14-C13-C18	2.81	117.45	114.65
2	B	601	5OA	C13-C14-N15	-2.75	120.17	124.49
2	B	601	5OA	C14-C13-C18	2.65	117.29	114.65
2	A	601	5OA	C16-N17-C18	2.62	122.51	117.18
2	B	601	5OA	C16-N17-C18	2.55	122.36	117.18
2	A	601	5OA	C5-N4-C3	2.53	115.95	111.75
2	B	601	5OA	C13-C12-C7	-2.39	120.19	122.86
2	A	601	5OA	C6-N1-C2	2.30	119.81	113.42
2	B	601	5OA	C6-N1-C2	2.27	119.72	113.42
2	B	601	5OA	C13-C18-N17	-2.19	119.36	121.49
2	A	601	5OA	C13-C18-N17	-2.15	119.39	121.49
2	A	601	5OA	C7-C8-CL1	-2.05	118.53	120.52


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	494/526 (93%)	0.27	29 (5%)	22 22	15, 32, 58, 68	0
1	B	471/526 (89%)	0.25	22 (4%)	31 30	16, 32, 58, 77	0
All	All	965/1052 (91%)	0.26	51 (5%)	26 25	15, 32, 58, 77	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	236	LEU	5.3
1	B	426	HIS	4.9
1	A	177	LEU	4.7
1	B	301	VAL	4.7
1	A	298	ASN	4.3
1	A	93	GLY	4.1
1	B	177	LEU	4.0
1	A	92	ASN	3.7
1	A	206	LEU	3.6
1	B	235	LYS	3.6
1	A	457	VAL	3.4
1	B	95	VAL	3.4
1	A	300	PRO	3.2
1	A	94	ASP	3.2
1	A	207	GLY	3.1
1	A	144	PRO	3.1
1	A	313	GLU	3.1
1	B	121	GLU	3.1
1	A	469[A]	PHE	3.1
1	A	143	HIS	3.1
1	B	525	LEU	3.0
1	A	199	LYS	3.0
1	B	205	THR	3.0
1	B	409	GLY	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	237	ALA	2.9
1	B	120	LYS	2.8
1	B	199	LYS	2.8
1	A	394	HIS	2.8
1	A	299	GLU	2.8
1	B	450	MET	2.7
1	B	425	ASP	2.7
1	A	523	GLU	2.7
1	A	91	LYS	2.6
1	B	451	ASP	2.6
1	B	116	HIS	2.5
1	A	323	PRO	2.5
1	A	22	THR	2.4
1	B	140	SER	2.4
1	A	353	ILE	2.3
1	B	457	VAL	2.3
1	A	3	SER	2.3
1	A	314	PHE	2.2
1	A	35	LYS	2.2
1	A	116	HIS	2.2
1	B	410	ASN	2.2
1	B	136	LEU	2.2
1	B	353	ILE	2.1
1	B	118	SER	2.1
1	A	322	LYS	2.1
1	B	463	ILE	2.1
1	A	456	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,



median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PO4	B	604	5/5	0.73	0.28	101,101,102,102	0
3	PO4	B	603	5/5	0.81	0.24	79,79,79,80	0
3	PO4	A	602	5/5	0.85	0.35	66,68,69,71	0
3	PO4	A	604	5/5	0.85	0.21	81,83,83,84	0
3	PO4	A	606	5/5	0.85	0.19	73,73,74,77	0
3	PO4	A	603	5/5	0.86	0.20	69,70,71,72	0
3	PO4	A	605	5/5	0.94	0.10	49,53,54,54	0
2	5OA	B	601	23/23	0.94	0.12	25,31,38,39	0
2	5OA	A	601	23/23	0.96	0.11	23,27,34,36	0
3	PO4	B	602	5/5	0.99	0.11	31,35,39,42	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 5OA B 601:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.