



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 11:22 pm BST

PDB ID : 4EIP
Title : Native and K252c bound RebC-10x
Authors : Goldman, P.J.; Ryan, K.S.; Howard-Jones, A.R.; Hamill, M.J.; Elliott, S.J.;
Walsh, C.T.; Drennan, C.L.
Deposited on : 2012-04-05
Resolution : 2.33 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2096 (2.36-2.32)
Clashscore	141614	2193 (2.36-2.32)
Ramachandran outliers	138981	2159 (2.36-2.32)
Sidechain outliers	138945	2160 (2.36-2.32)
RSRZ outliers	127900	2067 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	549	<div> <div>5%</div> <div> <div></div> <div>81%</div> <div>11%</div> <div>• 5%</div> </div> </div>
1	B	549	<div> <div>5%</div> <div> <div></div> <div>81%</div> <div>11%</div> <div>• 7%</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8306 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative FAD-monooxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	521	Total	C	N	O	S	1	1	0
			3997	2504	757	726	10			
1	B	510	Total	C	N	O	S	0	0	0
			3875	2434	723	709	9			

There are 60 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	INITIATING METHIONINE	UNP Q8KI25
A	-18	GLY	-	EXPRESSION TAG	UNP Q8KI25
A	-17	SER	-	EXPRESSION TAG	UNP Q8KI25
A	-16	SER	-	EXPRESSION TAG	UNP Q8KI25
A	-15	HIS	-	EXPRESSION TAG	UNP Q8KI25
A	-14	HIS	-	EXPRESSION TAG	UNP Q8KI25
A	-13	HIS	-	EXPRESSION TAG	UNP Q8KI25
A	-12	HIS	-	EXPRESSION TAG	UNP Q8KI25
A	-11	HIS	-	EXPRESSION TAG	UNP Q8KI25
A	-10	HIS	-	EXPRESSION TAG	UNP Q8KI25
A	-9	SER	-	EXPRESSION TAG	UNP Q8KI25
A	-8	SER	-	EXPRESSION TAG	UNP Q8KI25
A	-7	GLY	-	EXPRESSION TAG	UNP Q8KI25
A	-6	LEU	-	EXPRESSION TAG	UNP Q8KI25
A	-5	VAL	-	EXPRESSION TAG	UNP Q8KI25
A	-4	PRO	-	EXPRESSION TAG	UNP Q8KI25
A	-3	ARG	-	EXPRESSION TAG	UNP Q8KI25
A	-2	GLY	-	EXPRESSION TAG	UNP Q8KI25
A	-1	SER	-	EXPRESSION TAG	UNP Q8KI25
A	0	HIS	-	EXPRESSION TAG	UNP Q8KI25
A	36	ASP	GLU	ENGINEERED MUTATION	UNP Q8KI25
A	37	ALA	GLN	ENGINEERED MUTATION	UNP Q8KI25
A	38	GLY	THR	ENGINEERED MUTATION	UNP Q8KI25
A	46	LYS	ARG	ENGINEERED MUTATION	UNP Q8KI25
A	48	SER	GLY	ENGINEERED MUTATION	UNP Q8KI25

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	117	ALA	GLN	ENGINEERED MUTATION	UNP Q8KI25
A	216	VAL	PHE	ENGINEERED MUTATION	UNP Q8KI25
A	231	SER	ALA	ENGINEERED MUTATION	UNP Q8KI25
A	239	ASN	ARG	ENGINEERED MUTATION	UNP Q8KI25
A	241	VAL	THR	ENGINEERED MUTATION	UNP Q8KI25
B	-19	MET	-	INITIATING METHIONINE	UNP Q8KI25
B	-18	GLY	-	EXPRESSION TAG	UNP Q8KI25
B	-17	SER	-	EXPRESSION TAG	UNP Q8KI25
B	-16	SER	-	EXPRESSION TAG	UNP Q8KI25
B	-15	HIS	-	EXPRESSION TAG	UNP Q8KI25
B	-14	HIS	-	EXPRESSION TAG	UNP Q8KI25
B	-13	HIS	-	EXPRESSION TAG	UNP Q8KI25
B	-12	HIS	-	EXPRESSION TAG	UNP Q8KI25
B	-11	HIS	-	EXPRESSION TAG	UNP Q8KI25
B	-10	HIS	-	EXPRESSION TAG	UNP Q8KI25
B	-9	SER	-	EXPRESSION TAG	UNP Q8KI25
B	-8	SER	-	EXPRESSION TAG	UNP Q8KI25
B	-7	GLY	-	EXPRESSION TAG	UNP Q8KI25
B	-6	LEU	-	EXPRESSION TAG	UNP Q8KI25
B	-5	VAL	-	EXPRESSION TAG	UNP Q8KI25
B	-4	PRO	-	EXPRESSION TAG	UNP Q8KI25
B	-3	ARG	-	EXPRESSION TAG	UNP Q8KI25
B	-2	GLY	-	EXPRESSION TAG	UNP Q8KI25
B	-1	SER	-	EXPRESSION TAG	UNP Q8KI25
B	0	HIS	-	EXPRESSION TAG	UNP Q8KI25
B	36	ASP	GLU	ENGINEERED MUTATION	UNP Q8KI25
B	37	ALA	GLN	ENGINEERED MUTATION	UNP Q8KI25
B	38	GLY	THR	ENGINEERED MUTATION	UNP Q8KI25
B	46	LYS	ARG	ENGINEERED MUTATION	UNP Q8KI25
B	48	SER	GLY	ENGINEERED MUTATION	UNP Q8KI25
B	117	ALA	GLN	ENGINEERED MUTATION	UNP Q8KI25
B	216	VAL	PHE	ENGINEERED MUTATION	UNP Q8KI25
B	231	SER	ALA	ENGINEERED MUTATION	UNP Q8KI25
B	239	ASN	ARG	ENGINEERED MUTATION	UNP Q8KI25
B	241	VAL	THR	ENGINEERED MUTATION	UNP Q8KI25

- Molecule 2 is 6,7,12,13-tetrahydro-5H-indolo[2,3-a]pyrrolo[3,4-c]carbazol-5-one (three-letter code: K2C) (formula: C₂₀H₁₃N₃O).




- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	170	Total 170	O 170	0	0
4	B	134	Total 134	O 134	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Putative FAD-monooxygenase



• Molecule 1: Putative FAD-monooxygenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	64.31Å 78.59Å 125.72Å 90.00° 99.87° 90.00°	Depositor
Resolution (Å)	41.37 – 2.33 49.32 – 2.33	Depositor EDS
% Data completeness (in resolution range)	97.7 (41.37-2.33) 97.7 (49.32-2.33)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.77 (at 2.34Å)	Xtriage
Refinement program	PHENIX 1.7.3 _928	Depositor
R, R_{free}	0.213 , 0.260 0.206 , 0.252	Depositor DCC
R_{free} test set	2616 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	35.0	Xtriage
Anisotropy	0.048	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 48.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8306	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: K2C, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/4096	0.47	1/5574 (0.0%)
1	B	0.24	0/3972	0.45	0/5410
All	All	0.26	0/8068	0.46	1/10984 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	47	VAL	CB-CA-C	-5.40	101.14	111.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3997	0	3936	42	0
1	B	3875	0	3795	32	0
2	A	24	0	13	1	0
3	A	53	0	31	0	0
3	B	53	0	31	3	0
4	A	170	0	0	5	0
4	B	134	0	0	4	0
All	All	8306	0	7806	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:416:GLU:HB3	1:A:417:PRO:HD2	1.41	1.00
1:B:46:LYS:HG3	1:B:47:VAL:HG23	1.48	0.95
1:B:108:THR:HG22	1:B:110:GLU:H	1.39	0.86
1:B:46:LYS:HE3	4:B:791:HOH:O	1.77	0.83
1:A:417:PRO:O	1:A:418:GLU:O	2.04	0.76
1:A:157:THR:HG22	1:A:159:ALA:H	1.51	0.75
1:B:130:GLU:O	1:B:133:ARG:NH1	2.22	0.72
1:A:278:LEU:HD11	1:A:302:SER:HA	1.72	0.72
1:B:39:ASP:HB3	1:B:135:ARG:HD3	1.73	0.70
1:B:372:ASP:O	1:B:381:ARG:NH2	2.25	0.69
1:B:46:LYS:CG	1:B:47:VAL:HG23	2.21	0.69
1:B:442:THR:HG23	1:B:444:THR:H	1.57	0.69
1:B:189:ARG:HG3	1:B:351:LEU:HD13	1.74	0.68
1:A:259:ARG:NH1	1:A:262:VAL:O	2.27	0.68
1:A:372:ASP:O	1:A:381:ARG:NH2	2.27	0.66
1:A:416:GLU:HB3	1:A:417:PRO:CD	2.22	0.63
1:B:85:ARG:HD3	1:B:263:ALA:HB2	1.81	0.62
1:A:39:ASP:OD1	1:A:39:ASP:N	2.34	0.60
1:A:373:ASP:HA	1:A:381:ARG:HH22	1.67	0.59
1:A:395:ARG:NH1	4:A:757:HOH:O	2.38	0.57
1:B:373:ASP:N	1:B:373:ASP:OD2	2.39	0.54
3:B:600:FAD:HM81	4:B:809:HOH:O	2.06	0.54
1:A:46:LYS:HD3	1:A:47:VAL:HG13	1.90	0.54
1:A:496:ARG:HD3	1:A:510:ARG:HD3	1.91	0.53
1:B:216:VAL:HG13	1:B:230:ARG:HG2	1.92	0.51
1:A:254:SER:OG	1:A:273:ASP:OD2	2.28	0.51
1:B:256:GLU:OE1	1:B:259:ARG:NH1	2.43	0.51
1:A:227:PHE:HB3	1:A:228:PRO:HD2	1.93	0.51
1:A:437:PRO:HB2	1:A:493:LEU:HD21	1.92	0.50
3:B:600:FAD:O5'	3:B:600:FAD:O1A	2.30	0.50
1:A:137:ARG:HD3	1:A:155:LEU:HD21	1.95	0.48
1:B:94:ILE:HD13	1:B:394:ARG:HB3	1.95	0.48
1:A:153:THR:HA	1:A:160:THR:HA	1.96	0.48
1:A:437:PRO:HD2	1:A:494:TYR:CZ	2.50	0.47
1:A:137:ARG:O	4:A:860:HOH:O	2.20	0.47
1:A:416:GLU:OE2	1:A:510:ARG:HD2	2.15	0.47
1:A:222:SER:O	1:A:226:ARG:HG2	2.16	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:56:GLU:OE2	1:B:108:THR:HG23	2.16	0.46
1:A:187:PRO:HA	1:A:188:PRO:HD3	1.80	0.46
1:A:154:ASP:OD1	1:A:156:ARG:HG2	2.15	0.46
1:A:72:TRP:HA	1:A:73:PRO:HD3	1.86	0.45
1:A:60:ARG:HD2	1:A:452:GLY:HA3	1.99	0.45
1:B:149:ARG:HG2	1:B:164:HIS:CE1	2.52	0.45
1:A:137:ARG:N	4:A:860:HOH:O	2.50	0.44
1:B:239:ASN:ND2	4:B:753:HOH:O	2.49	0.44
1:B:442:THR:HG22	1:B:445:THR:H	1.83	0.44
1:A:372:ASP:O	1:A:377:GLY:HA3	2.18	0.44
1:B:321:TRP:CD1	1:B:504:ASP:HB3	2.53	0.44
1:A:436:ALA:HA	1:A:437:PRO:HD3	1.75	0.43
1:B:121:ALA:HB3	1:B:122:PRO:HD3	2.00	0.43
1:B:201:ARG:NH1	1:B:267:GLU:OE2	2.51	0.43
1:A:476:ARG:NH2	4:A:855:HOH:O	2.52	0.43
1:B:254:SER:HB3	1:B:273:ASP:OD2	2.18	0.43
1:B:487:ALA:HA	1:B:488:PRO:HD3	1.88	0.43
1:A:190:HIS:HB2	1:A:280:HIS:CD2	2.54	0.42
1:B:212:ARG:HD2	4:B:834:HOH:O	2.19	0.42
1:A:130:GLU:H	1:A:130:GLU:CD	2.21	0.42
2:A:601:K2C:H4	2:A:601:K2C:O24	2.18	0.42
1:A:137:ARG:NH1	1:A:139:ASP:OD2	2.52	0.42
1:B:109:PRO:HD3	1:B:438:HIS:CG	2.55	0.42
1:B:46:LYS:HG2	1:B:47:VAL:N	2.33	0.42
3:B:600:FAD:H1'1	3:B:600:FAD:H9	1.80	0.42
1:A:502:ARG:HG3	1:A:508:ALA:HB2	2.03	0.41
1:A:109:PRO:HD3	1:A:438:HIS:CG	2.55	0.41
1:B:380:ILE:H	1:B:380:ILE:HG12	1.40	0.41
1:A:223:SER:O	1:A:226:ARG:HD2	2.21	0.41
1:B:139:ASP:OD2	1:B:153:THR:OG1	2.26	0.41
1:B:442:THR:CG2	1:B:444:THR:H	2.31	0.41
1:A:121:ALA:HB3	1:A:122:PRO:HD3	2.01	0.41
1:A:416:GLU:CB	1:A:417:PRO:CD	2.94	0.41
1:B:344:ARG:CZ	1:B:348:ILE:HD11	2.51	0.41
1:B:41:THR:HG22	1:B:43:THR:HG23	2.03	0.41
1:A:353:GLU:HG3	1:A:402:ILE:HD11	2.02	0.41
1:A:431:ARG:NH1	4:A:841:HOH:O	2.54	0.40
1:B:147:HIS:NE2	1:B:149:ARG:HD2	2.36	0.40
1:A:177:PRO:HA	1:A:180:LYS:HE3	2.02	0.40
1:A:47:VAL:CG2	1:A:47:VAL:O	2.66	0.40
1:A:7:THR:O	1:A:165:ALA:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	518/549 (94%)	499 (96%)	15 (3%)	4 (1%)	19	20
1	B	504/549 (92%)	486 (96%)	18 (4%)	0	100	100
All	All	1022/1098 (93%)	985 (96%)	33 (3%)	4 (0%)	34	38

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	417	PRO
1	A	418	GLU
1	A	375	PRO
1	A	304	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	407/432 (94%)	387 (95%)	20 (5%)	25	31
1	B	392/432 (91%)	369 (94%)	23 (6%)	19	22
All	All	799/864 (92%)	756 (95%)	43 (5%)	22	26

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	33	LEU
1	A	39	ASP
1	A	46	LYS
1	A	53	ARG
1	A	130	GLU
1	A	144	ARG
1	A	153	THR
1	A	168	LEU
1	A	225	LEU
1	A	226	ARG
1	A	230	ARG
1	A	237	LEU
1	A	277	HIS
1	A	373	ASP
1	A	394	ARG
1	A	441	LEU
1	A	471	ARG
1	A	491	HIS
1	A	496	ARG
1	A	510	ARG
1	B	12	LEU
1	B	21	LEU
1	B	46	LYS
1	B	66	GLN
1	B	101	THR
1	B	166	ARG
1	B	168	LEU
1	B	185	ASP
1	B	201	ARG
1	B	211	GLU
1	B	219	LEU
1	B	259	ARG
1	B	328	ARG
1	B	352	GLU
1	B	373	ASP
1	B	380	ILE
1	B	409	ARG
1	B	431	ARG
1	B	442	THR
1	B	448	LEU
1	B	458	LEU
1	B	491	HIS
1	B	517	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	371	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	FAD	A	602	-	51,58,58	1.85	5 (9%)	60,89,89	1.95	13 (21%)
3	FAD	B	600	-	51,58,58	1.84	5 (9%)	60,89,89	1.93	13 (21%)
2	K2C	A	601	-	22,29,29	1.83	3 (13%)	27,45,45	2.44	11 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	A	602	-	-	13/30/50/50	0/6/6/6
3	FAD	B	600	-	-	9/30/50/50	0/6/6/6
2	K2C	A	601	-	-	-	0/6/6/6

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	602	FAD	C4X-C10	9.61	1.48	1.38
3	B	600	FAD	C4X-C10	9.57	1.48	1.38
2	A	601	K2C	C7-C22	-5.39	1.46	1.50
3	A	602	FAD	C4-C4X	4.24	1.48	1.41
3	B	600	FAD	C4-C4X	4.22	1.48	1.41
3	A	602	FAD	C9A-C5X	3.50	1.49	1.42
3	B	600	FAD	C9A-C5X	3.46	1.49	1.42
3	B	600	FAD	C8-C7	3.29	1.49	1.40
3	A	602	FAD	C8-C7	3.25	1.49	1.40
2	A	601	K2C	C7-N6	2.97	1.48	1.45
2	A	601	K2C	O24-C5	2.96	1.29	1.23
3	A	602	FAD	C5A-C4A	2.43	1.47	1.40
3	B	600	FAD	C5A-C4A	2.39	1.47	1.40

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	600	FAD	C4-N3-C2	8.46	122.28	115.14
3	A	602	FAD	C4-N3-C2	8.45	122.28	115.14
2	A	601	K2C	C7-N6-C5	-6.47	107.63	113.85
3	A	602	FAD	C4-C4X-C10	-5.22	116.49	119.95
2	A	601	K2C	C23-C5-N6	5.21	111.64	106.37
3	B	600	FAD	C4-C4X-C10	-5.05	116.61	119.95
3	A	602	FAD	C4X-N5-C5X	4.00	120.77	116.77
3	B	600	FAD	C4X-N5-C5X	3.90	120.67	116.77
2	A	601	K2C	C23-C22-C21	-3.78	117.34	122.42
2	A	601	K2C	C8-C20-C19	3.67	123.03	118.17
3	B	600	FAD	C4X-C4-N3	-3.63	118.47	123.43
3	A	602	FAD	C4X-C4-N3	-3.58	118.53	123.43
2	A	601	K2C	C4-C15-C14	3.36	122.63	118.17
3	B	600	FAD	P-O3P-PA	-3.29	121.53	132.83
2	A	601	K2C	O24-C5-C23	-3.28	125.43	129.32
3	B	600	FAD	N3A-C2A-N1A	-3.20	123.68	128.68
3	A	602	FAD	N3A-C2A-N1A	-3.20	123.68	128.68
3	A	602	FAD	C4-C4X-N5	3.19	122.25	118.60
3	A	602	FAD	C3B-C2B-C1B	3.15	105.73	100.98

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	FAD	P-O3P-PA	-3.10	122.19	132.83
3	A	602	FAD	C1'-N10-C9A	3.04	120.68	118.29
3	B	600	FAD	C1'-N10-C10	3.01	121.10	118.41
3	B	600	FAD	C4-C4X-N5	3.00	122.02	118.60
3	B	600	FAD	C1'-N10-C9A	2.87	120.55	118.29
3	B	600	FAD	C4A-C5A-N7A	-2.78	106.50	109.40
3	B	600	FAD	C9A-N10-C10	-2.72	118.34	121.91
3	A	602	FAD	C1'-N10-C10	2.71	120.84	118.41
3	A	602	FAD	C4A-C5A-N7A	-2.69	106.59	109.40
2	A	601	K2C	C22-C7-N6	2.67	104.48	101.76
3	B	600	FAD	C5X-C9A-N10	2.65	119.64	117.72
3	A	602	FAD	C9A-N10-C10	-2.62	118.47	121.91
3	A	602	FAD	C5X-C9A-N10	2.49	119.52	117.72
2	A	601	K2C	C21-C20-C19	2.29	108.57	106.09
2	A	601	K2C	C8-C20-C21	-2.18	128.40	134.64
3	B	600	FAD	C3B-C2B-C1B	2.15	104.21	100.98
2	A	601	K2C	C9-C8-C20	-2.04	117.43	120.86
2	A	601	K2C	O24-C5-N6	-2.00	122.92	125.27

There are no chirality outliers.

All (22) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	602	FAD	C5B-O5B-PA-O2A
3	A	602	FAD	C3'-C4'-C5'-O5'
3	A	602	FAD	O4'-C4'-C5'-O5'
3	A	602	FAD	C5'-O5'-P-O1P
3	A	602	FAD	C5'-O5'-P-O2P
3	A	602	FAD	C5'-O5'-P-O3P
3	B	600	FAD	C3'-C4'-C5'-O5'
3	B	600	FAD	O4'-C4'-C5'-O5'
3	B	600	FAD	C5'-O5'-P-O1P
3	B	600	FAD	C5'-O5'-P-O2P
3	A	602	FAD	O4B-C4B-C5B-O5B
3	A	602	FAD	C3B-C4B-C5B-O5B
3	B	600	FAD	PA-O3P-P-O5'
3	A	602	FAD	C5B-O5B-PA-O3P
3	B	600	FAD	O4B-C4B-C5B-O5B
3	A	602	FAD	C5B-O5B-PA-O1A
3	A	602	FAD	C2'-C3'-C4'-O4'
3	A	602	FAD	O3'-C3'-C4'-C5'
3	B	600	FAD	C5'-O5'-P-O3P

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	602	FAD	O3'-C3'-C4'-O4'
3	B	600	FAD	P-O3P-PA-O1A
3	B	600	FAD	C3B-C4B-C5B-O5B

There are no ring outliers.


2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	600	FAD	3	0
2	A	601	K2C	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand FAD B 600




Bond lengths



Bond angles




Torsions




Rings


Ligand K2C A 601




Bond lengths



Bond angles



Torsions



Rings

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	521/549 (94%)	0.39	30 (5%) 23 32	17, 31, 56, 68	0
1	B	510/549 (92%)	0.47	26 (5%) 28 39	24, 40, 71, 92	0
All	All	1031/1098 (93%)	0.43	56 (5%) 25 36	17, 35, 64, 92	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	155	LEU	7.2
1	A	418	GLU	4.5
1	A	160	THR	4.4
1	B	423	THR	4.2
1	A	529	ALA	4.0
1	A	417	PRO	4.0
1	A	41	THR	4.0
1	B	529	ALA	3.9
1	B	365	GLU	3.9
1	A	137	ARG	3.8
1	A	139	ASP	3.7
1	B	43	THR	3.7
1	B	223	SER	3.6
1	B	248	SER	3.5
1	A	40	GLY	3.5
1	A	43	THR	3.4
1	B	221	LEU	3.4
1	A	369	GLY	3.1
1	A	420	GLU	3.1
1	B	247	ALA	3.1
1	A	246	ASP	3.1
1	A	39	ASP	3.1
1	A	151	THR	2.9
1	B	372	ASP	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	351	LEU	2.9
1	B	424	GLY	2.9
1	B	245	ASP	2.9
1	A	419	THR	2.8
1	B	349	THR	2.7
1	A	161	ARG	2.6
1	B	224	SER	2.6
1	B	516	ALA	2.6
1	B	379	ARG	2.6
1	A	252	MET	2.4
1	B	461	GLY	2.4
1	A	27[A]	HIS	2.4
1	A	5	ILE	2.3
1	A	152	ILE	2.3
1	A	409	ARG	2.3
1	A	259	ARG	2.2
1	B	260	ARG	2.2
1	B	201	ARG	2.2
1	B	250	SER	2.2
1	A	153	THR	2.2
1	B	41	THR	2.2
1	B	471	ARG	2.2
1	B	402	ILE	2.2
1	A	156	ARG	2.1
1	A	44	HIS	2.1
1	A	162	ALA	2.1
1	A	163	VAL	2.1
1	B	249	LYS	2.1
1	A	157	THR	2.1
1	A	135	ARG	2.1
1	B	106	GLU	2.0
1	B	369	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ


There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	FAD	A	602	53/53	0.87	0.23	34,42,54,56	0
2	K2C	A	601	24/24	0.91	0.16	29,30,34,35	0
3	FAD	B	600	53/53	0.95	0.15	28,34,43,47	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





Electron density around K2C A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD B 600:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.